SD11
A3
1973

Forest Service Research Accomplishments/1973

LIBRARY

SEP 30 1974

ROCKY MOUNTAIN STATION

UNITED STATES DEPARTMENT OF AGRICULTURE . FOREST SERVICE

FOREST SERVICE - U. S. DEPARTMENT OF AGRICULTURE
FOREST AND RANGE EXPERIMENT STATIONS,
FOREST PRODUCTS LABORATORY



Revised 4 - 72

UNITED STATES DEPARTMENT OF AGRICULTURE
FOREST SERVICE
Washington, D. C. 20250

1380(4000)

Mr. Robert W. Long
Assistant Secretary
U.S. Department of Agriculture
Washington, D. C. 20250

Dear Secretary Long:

I am pleased to send you this report of Forest Service Research accomplishments for 1973.

The report presents a broad overview of the Forest Service research program--a program aimed at solving the critical problems faced by public and private resource managers. Major steps in our continuing quest for knowledge to improve management of America's natural resources are highlighted.

A unique feature of this year's report is that it has been produced directly from computer output, without going through the usual steps of type composition. All of the references are now stored in a computerized data bank. Future accomplishment reports will be produced from field inputs to the computer system.

Sincerely,

John R. McGuire

JOHN R. McGUIRE
Chief

Use Pesticides Safely

FOLLOW THE LABEL

U.S. DEPARTMENT OF AGRICULTURE

NOTICE: The identification and description of commercial products in this publication are solely for information purposes. Endorsement of any commercial product is not intended and must not be inferred. Readers are cautioned to handle all pesticides, herbicides, and fungicides mentioned in this publication strictly in accordance with manufacturer's labels. These chemicals are harmful to people, farm animals, wildlife, and fish, and can contaminate water supplies.

FOREWORD

Forest Service research plans are coordinated through the Assistant Secretary for Conservation, Research and Education with research in other USDA agencies, with that conducted under the Hatch Act at land grant institutions, and with that conducted under the McIntire-Stennis Act at schools of forestry. Coordination of research with other educational institutions, private enterprises, nonprofit institutions, and other public agencies is maintained mostly through direct contact between people of these organizations and those of the Forest and Range Experiment Stations. Federal, State, industry, and university cooperation in solving mutual problems is achieved through cooperative agreements that provide for joint development and support of the research by the cooperators.

This report is arranged by research subject areas as a convenience to the reader. Each accomplishment is summarized in a single, short paragraph. Throughout this report an abbreviation signifies the Forest Service research unit best able to supply detailed information and copies of the publications listed. Abbreviations and addresses of the research units are given on the following page.

Issued June 1974

FOREST SERVICE RESEARCH UNITS

PNW - Director
Pacific Northwest Forest and Range Experiment Station
809 NE 6th Avenue, P.O. Box 3141
Portland, Oregon 97208

PSW - Director
Pacific Southwest Forest and Range Experiment Station
1960 Addison Street, P.O. Box 245
Berkeley, California 94701

INT - Director
Intermountain Forest and Range Experiment Station
507 25th Street
Ogden, Utah 84401

RM - Director
Rocky Mountain Forest and Range Experiment Station
240 West Prospect Street
Fort Collins, Colorado 80521

NC - Director
North Central Forest Experiment Station
Folwell Avenue
St. Paul, Minnesota 55101

NE - Director
Northeastern Forest Experiment Station
6816 Market Street
Upper Darby, Pennsylvania 19082

SE - Director
Southeastern Forest Experiment Station
Post Office Building, P.O. Box 2570
Asheville, North Carolina 28802

SO - Director
Southern Forest Experiment Station
T-10210 Federal Building
701 Loyola Avenue
New Orleans, Louisiana 70113

FPL - Director
Forest Products Laboratory
North Walnut Street
P.O. Box 5130
Madison, Wisconsin 53705

ITF - Director
Institute of Tropical Forestry
University of Puerto Rico
Agricultural Experiment Station Grounds
P.O. Box AQ
Rio Piedras, Puerto Rico 00928

WO - Deputy Chief for Research
Forest Service
U.S. Department of Agriculture
Washington, D.C. 20250

RESEARCH ACCOMPLISHMENTS - 1973

CONTENTS

	Paragraph Number	Page
I. IMPROVING ENVIRONMENTAL QUALITY, PRODUCTIVITY, AND USEFULNESS		
A. Reducing Pollution		
1. Controlling soil erosion.....	1- 10	1
2. Chemicals in the forest environment.....	11- 18	1
3. Air pollution.....	19- 25	2
B. Improving Wildland Animal Habitats		
1. Evaluating and improving animal habitat resources.....	26- 36	2
2. Characteristics and forage and browse values of plants.....	37- 41	3
3. Livestock and grazing management.....	42- 45	4
4. Integrating other uses and wildlife.....	46- 47	4
C. Improving Social and Amenity Values		
1. Environmental amenities--landscapes and open space.....	48- 52	4
2. Environmental amenities--wilderness.....	53- 57	4
3. Managing recreational opportunities.....	58- 75	5
D. Trees to Enhance the Environment.....	76- 80	6
E. Improving Environmental Quality Through Fire Management		
1. Fire prevention, hazard reduction, and prescribed burning.....	81- 97	7
2. Fire management methods and systems.....	98-108	8
3. Forest fire science.....	109-118	8
F. Improving Insect and Disease Control		
1. Detection and evaluation.....	119-133	9
2. Biology and understanding.....	134-227	10
3. Pest control techniques.....	228-273	15
4. Pest management strategies.....	274-277	18
G. Accelerating Forest Resource Surveys		
1. Forest inventory.....	278-283	18
2. Timber utilization and production.....	284-290	19
3. Survey techniques.....	291-296	19
H. Economics of Forest Management		
1. Methods of financial evaluation and planning.....	297-300	20
2. Timber growing economics.....	301-305	20
3. Multiple-use economics.....	306-308	20
4. Impacts on forest industry and regional economies.....	309-311	20

	Paragraph Number	Page
I. Improving Engineering Systems	312-317	21
J. Improving Water Quality and Yield		
1. Improving water yields	318-324	21
2. Managing, rehabilitating, and improving watersheds	325-343	21
3. Preventing watershed damage	344-347	23
4. Water quality	348-349	23
K. Bettering Silvicultural Systems		
1. Natural regeneration	350-357	23
2. Silvicultural methods	358-363	24
3. Ecological relationships	364-369	24
L. Remote Sensing Methods	370-374	25
M. Weather Modification and Weather Effects	375-382	25
II. IMPROVING UTILIZATION AND EXTENDING WOOD SUPPLIES		
A. Intensive Culture Methods		
1. Site evaluation and soil improvement	383-398	26
2. Artificial regeneration	399-431	27
3. Stand improvement	432-441	29
4. Animal damage	442-444	30
5. Growth requirements	445-462	30
6. Growth and yield	463-475	31
7. Forest measurements	476-487	32
8. Management planning	488-489	33
9. Maple syrup production	490-496	33
10. Naval stores	497	34
B. Breeding Improved Trees		
1. Distribution and classification of forest trees	498-500	34
2. Inherent variation	501-519	34
3. Insect-disease resistance	520-528	36
4. Tree breeding methodology	529-553	36
C. Improving Uses and Protection of Wood		
1. Utilization potential and processing of wood	554-594	38
2. Wood chemistry and fiber products	595-604	41
3. Wood engineering	605-615	42
4. Biological degradation	616-623	42
5. Prevention and control of wood-destroying organisms	624-631	43
6. Miscellaneous	632	43
D. Marketing Under-used Species and Residues	633-636	44
E. Supply, Demand, and Price Analysis	637-647	44
F. Marketing--General	648-651	44

RESEARCH ACCOMPLISHMENTS - 1973

PUBLICATION LIST WITH AUTHOR INDEX

	Publication Number	Page
FOREST ENVIRONMENT RESEARCH		
Watershed Management Research		
Controlling soil erosion	1- 15	47
Chemicals in the forest environment	16- 26	47
Improving water yields	27- 48	48
Managing, rehabilitating, and improving watersheds	49- 125	49
Preventing watershed damage	126- 139	51
Water quality	140- 147	52
Wildlife Habitat and Range Research		
Evaluating and improving animal habitat resources	148- 184	52
Characteristics and values of forage and browse plants	185- 212	53
Livestock and grazing management	213- 243	54
Integrating other uses and wildlife	244- 252	55
Forest Recreation Research		
Environmental amenities--landscapes and open space	253- 258	56
Environmental amenities--wilderness	259- 266	56
Managing recreational opportunities	267- 296	56
FOREST INSECT AND DISEASE RESEARCH		
Forest Insect Research		
Chemicals in the forest environment	297- 299	58
Detection and evaluation	300- 313	58
Biology and understanding	314- 392	58
Pest control techniques	393- 421	61
Pest management strategies	422	62
Remote sensing methods	423- 431	62
Biological degradation	432- 442	63
Prevention and control of wood-destroying organisms	443- 448	63
Forest Disease Research		
Air pollution	449- 459	63
Detection and evaluation	460- 472	64
Biology and understanding	473- 569	64
Pest control techniques	570- 603	67
Pest management strategies	604- 606	68
Biological degradation	607- 613	68
Prevention and control of wood-destroying organisms	614- 618	69

	Publication Number	Page
FOREST FIRE AND ATMOSPHERIC SCIENCES RESEARCH		
Air pollution	619- 621	70
Fire prevention, hazard reduction, and prescribed burning	622- 648	70
Fire management methods and systems	649- 662	71
Forest fire science	663- 679	71
Biology and understanding	680- 681	72
Water quality	682	72
Natural regeneration	683- 685	72
Ecological relationships	686	72
Weather modification and weather effects	687- 697	72
Miscellaneous	698	72
TIMBER MANAGEMENT RESEARCH		
Evaluating and improving animal habitat resources	699	73
Managing recreational opportunities	700- 701	73
Trees to enhance the environment	702- 712	73
Fire prevention, hazard reduction, and prescribed burning	713	73
Biology and understanding	714	73
Pest control techniques	715- 716	73
Forest inventory	717	73
Survey techniques	718	73
Natural regeneration	719- 733	73
Silvicultural methods	734- 748	74
Ecological relationships	749- 757	75
Site evaluation and soil improvement	758- 780	75
Artificial regeneration	781- 837	76
Stand improvement	838- 854	77
Animal damage	855- 858	78
Growth requirements	859- 888	78
Growth and yield	889- 908	79
Forest measurements	909- 925	80
Management planning	926- 933	80
Maple syrup production	934- 942	81
Naval stores	943- 947	81
Distribution and classification of forest trees	948- 951	81
Inherent variation	952- 980	81
Insect - disease resistance	981- 990	82
Tree breeding methodology	991-1023	83
Utilization potential and processing of wood	1024-1027	84
Miscellaneous	1028-1038	84

	Publication Number	Page
FOREST ECONOMICS AND MARKETING RESEARCH		
Forest Survey		
Forest inventory	1039-1057	85
Timber utilization and production	1058-1077	85
Survey techniques	1078-1097	86
Forest Economics Research		
Methods of financial evaluation and planning	1098-1102	87
Timber growing economics	1103-1122	87
Multiple-use economics	1123-1132	88
Impacts on forest industry and regional economies	1133-1139	88
Forest Products Marketing Research		
Marketing under-used species and residues	1140-1145	88
Supply, demand, and price analysis	1146-1176	88
Marketing -- general	1177-1193	89
FOREST PRODUCTS AND ENGINEERING RESEARCH		
Forest Engineering Research		
Improving engineering systems	1194-1218	91
Forest Products Utilization Research		
Utilization potential and processing of wood	1219-1312	92
Wood chemistry and fiber products	1313-1352	95
Wood engineering	1353-1395	96
Miscellaneous	1396-1407	97
AUTHORS		99
INDEX		107

I. IMPROVING ENVIRONMENTAL QUALITY, PRODUCTIVITY, AND USEFULNESS

A. Reducing Pollution

1. Controlling soil erosion

1. Grooming a ski area development will result in destruction of plant cover and soil disturbance. Revegetating these areas promptly to prevent erosion and restore soil-plant nutrient cycles is desirable. Field trials show that satisfactory vegetative cover could be obtained by fall seeding and applying fertilizers. Results also showed that covering the fertilizer and seed was at least as important as the choice of species to seed. PNW(5).

2. Basic stratification of landforms followed by careful analysis of factors contributing to unstable conditions can be used to classify unstable areas. Use of this method has resulted in a system to rate slopes as to whether they are in low, medium, or high hazard class. Management alternatives and developmental activities can then be designed to fit each stability situation encountered. PNW(11).

3. Added emphasis has been placed on development of improved logging systems which minimize soil disturbance, especially by road construction. Balloon logging caused substantially less soil disturbance than previous studies had shown for tractor, high-lead, and skyline logging methods. Deeply disturbed and compacted soil areas occupied 4.3 percent of the total area, and 15.8 percent of the area was classed as slightly disturbed. PNW(4).

4. Soil movement may decrease site productivity significantly either by removing the more productive surface soil or by covering the surface with less productive material. Clearcut logging and slash burning in a steep watershed resulted in increased rates of soil movement, especially on slopes unprotected by organic debris. The first growing season after burning, erosion occurred as dry ravel. Rapid invasion by vegetation during second year essentially halted soil movement on all slopes except extremely stony talus areas. PNW(9).

5. Wildfires induce numerous physical, chemical, and biological changes in soils. A water repellent soil is formed when a fire burns over a wildland area. The degree of water repellency that is developed depends upon soil physical properties, biological factors, and the soil temperatures during a wildfire. Chemical remedial treatments have only been partially successful in reducing water repellency and subsequent erosion on brushland. PSW(3).

6. Soil losses under simulated rainfall were used to test erodibility indices of soils from the Koolau and Waianae ranges on Oahu, Hawaii. Gross splash was related to soil erodibility index, bulk density, infiltration, and saturation moisture content; in contrast, maximum splash erosion variation was related to organic matter content as well as to an erodibility index and bulk density of the soil. Ash and basalt colluvium soils require more careful management than do basalt soils because of their higher splash rates. PSW(15).

7. The degree of weathering, fracturing density, and mineral grain size are important factors to assess in predicting slope stability. A key has been developed that provides a practical tool for classifying these factors for granitic rocks of the Idaho Batholith. This information is particularly helpful in evaluating the potential impact from road construction and for devising the most efficient and economical stabilization measures. INT(1).

8. Sediments deposited in reservoirs are important because they occupy usable water storage space. Volume weights collected in small sediment retention reservoirs in a number of forested watersheds are related to the organic content of the sediment. The relationship is a reliable basis for estimating sediment volume weight; better than the more conventional approach of relating volume weight to sediment particle size. The organic sediments are surprisingly high from the forested watersheds. These data have important implications to the functioning of the aquatic ecosystems found on forested watersheds. INT(6).

9. Temporary subsurface water tables are built up during periods of rapid water inflow due to a zone of lower permeability beneath the soil surface. Road construction incises subsurface flow. The water which previously ran through the soil is collected on the road surface. This can and often does create severe erosion and downstream sedimentation problems. Two years of data suggest that up to 8 area-inches of water can be intercepted by roads in steep mountainous areas, or about 40 percent of the total water yield. INT(7).

10. An analysis of sediment yields following road construction suggests that the largest proportion of sediment is produced the first year following construction. Sediment yields decrease rapidly with time to a minimum level within 3 or 4 years, which indicates that stabilization measures must be installed immediately after construction. Revegetation measures should be coupled with some procedure for providing mechanical stabilization during the period that vegetation is becoming established. INT(8).

2. Chemicals in the forest environment

11. Small quantities of DDT were leached through columns of DDT-treated forest floor material. Leaching of DDT was increased thirtyfold by the addition of urea fertilizer. In field tests, no detectable levels of DDT were in leachate from tension plate lysimeters installed 2 centimeters below the mineral soil surface. Lack of DDT recovery is attributed to extensive absorption on immobile material below the organic layer. These experiments indicate that humic substances may be important carriers of DDT in the organic layer of a forest soil, but they may be relatively ineffective in the mineral soil profile. PNW(16).

12. One month after aerial application of DDT (12 oz/acre) to an eastern Oregon forest, 3 oz/acre of DDT residues were detected in the forest floor; 3 years later, the DDT content had decreased by more than 50 percent, and had not leached into the surface mineral soil. No effect of the spraying was noted on soil microbial populations, nitrification rate, or amount of nitrate nitrogen in the soil. Approximately one-

third of the sprayed chemical reached the forest floor. PNW(26).

13. Herbicide applications are hazardous to fish when there is a probability fish will be exposed to herbicide concentrations which are toxic. General weed and brush control using herbicides becomes important to fish habitat when they are applied to lands adjacent to waterways. Ninety-eight herbicides were grouped into four classes based on median tolerance limits for fish. These classes may be used as an aid in selecting herbicides and formulations of least hazard to fish considering the toxicity of the herbicide and the potential for its entry into the water. PNW(21).

14. Prescribed fire is used as a tool for disposing of logging debris and for preparing the sites for planting. Logging and burning may temporarily impair watershed protection and increase overland flow and erosion of soils derived from Belt series rocks. Small nutrient losses occuring in the sediment and the overland flow during the denuded periods represent only a fraction of the available nutrients on these sites. Damage to soil and water resources and to the nutrient reservoir on forest sites as a result of clearcut logging and burning is not serious from the standpoint of future site stability and productivity. INT(19).

15. Surface mining exposes rock strata to weathering and erosion, and as a result some chemicals may reach streams and adversely affect water resources. Chemical quality of water in six small watersheds in Kentucky indicated surface mining for coal may result in chemical pollution of streams even in areas where acid is no problem. Greatest increases were in sulfate, calcium, and magnesium. Some elements continue to increase at least 2 years after mining; others peak quickly after land disturbance and soon return to premining levels. NE(18).

16. Insect mortality and aerial spray deposit often cannot be correlated when cards are used to collect droplets. Controlled experiments in a spray chamber with known emission rates and air movement indicated that under some conditions conventional spray cards may give an unrealistic picture of the spray deposit spectrum and prevent a precise correlation between deposit and insect kill. PSW(298).

17. The use of synergists can reduce the amount of insecticides needed to control and provide some understanding of the mode of pesticide detoxification. Screening of 40 compounds for synergism with one to five insecticides on western spruce budworm revealed that Zectran, pyrethrins, and SD-9077 were not synergized by any of the commonly known synergists. Because of larger losses of fluid from the insect body following poisoning with these insecticides, their primary mode of action may be exercised through increased secretions in this insect. PSW(299).

18. Previous methods for obtaining pure, acylated amitrole for use in studies of systemic pesticides have been unsatisfactory. Pacific Southwest Station chemists have now determined that N-acetylimidazole acylates amitrole to 2-acetyl-3-amino-1, 2, 4-triazole without the formation of other isomers. By using 3-amino-1, 2, 4-triazole-5-^{14}C, radiopurity was maintained. PSW(297).

3. Air pollution

19. Continuing chronic sources of phytotoxicants threaten trees over much of the country. Ozone, sulfur dioxide, and fluorides are currently the primary offenders, but others such as oxides of nitrogen, hydrogen chloride, and ammonia may become more destructive if they become more prevalent as predicted in long-range forecasts. A realistic appraisal of the magnitude of the impact of air pollution to forest trees can help in setting air quality standards. NE(450).

20. Photochemical smog from the Central Valley of California was shown to move to the Mineral King Valley in the Sierra Nevada. Measurements made at ground stations and by instrumented aircraft suggest that nitrogen oxides and hydrocarbons move upslope with afternoon winds and produce ozone in high mountain sites during following sunny days. It is suspected that the mixed conifer forest of the Sierra Nevada could be jeopardized by an increasing supply of smog in the Central Valley. PSW(456).

21. In the San Bernardino Mountains of California, oxidant air pollution damage to conifers decreases downwind from the Los Angeles-San Bernardino air basin. Over a 5-year period, the average durations of oxidant concentrations above 0.08 ppm ranged from 12.6 hours in the west to 5.7 hours 28 miles to the east. Nearly one-tenth of the ponderosa and Jeffrey pines in the western area died during this period and all portions of the coniferous forest in the 5,000- to 7,000-foot elevational zone suffered from significant exposure to pollutants. PSW(455,458).

22. An experimental chamber has been developed for identifying oxidant air pollution damage to vegetation in forest locations. Natural ventilation exhausts warm air through a stack and draws fresh air through activated carbon filters. Comparing plants in chambers with and without filters will improve the ability to identify acute and chronic oxidant air pollution damage to native vegetation. PSW(459).

23. Anatomical factors in pine needles indicate that sensitivity to ozone increases with an increasing capacity for gas exchange. Injury in the more tolerant pines is limited to cells in the near vicinity of stomates, whereas, generalized injury occurs in sensitive species. Understanding the mechanism of pollution tolerance and sensitivity is useful for predicting pollution impact and for developing new lines of tolerant trees. PSW(454).

24. Trees were shown to reduce the amount of particulate pollution in the air space beneath their canopies. Evaluation of suspended particulates and dust fall indicated that conifers were slightly more effective at removing particles than were hardwoods, but both significantly improved air quality as compared to open terrain. An improved understanding of the interplay of trees and air pollutants can lead to a more harmonious balance between man and his environment. NE(451).

25. Aircraft monotoring of photochemical smog in southern California has identified multilayers of pollution in the atmosphere with penetrations as far inland as Mineral King Valley from the Central Valley and up to 80 miles from the South Coast Basin. Severe damage to coniferous species has been documented. PSW(619-621).

B. Improving Wildland Animal Habitats

1. Evaluating and improving animal habitat resources

26. Considerable progress has been made on the application of camera-film systems to inventory and analyze the multiresources on rangelands. Scientists of Fort Collins, Colorado, are using 70-mm photographs to sample strips of landscape photographed at a small scale. They can determine the aerial extent of complex plant communities, density, cover and distribution of various plant species, or amount of bare soil; herbage yield; and animal populations (especially effective to estimate overwinter death losses of mule deer). RM(157,158).

27. Mice to accomodate the biological control of sagebrush? A possibility. Long-tailed voles kill and damage sagebrush and other shrubs over large areas during cyclic population peaks in Utah. Damage is greatest when a dense ungrazed herbaceous understory is present and when the snowpack persists throughout the winter. Population peaks are spaced 4 or 5 years apart which would fit a grazing cycle for livestock that rests a given unit of range from grazing every 4 or 5 years. A carefully coordinated management plan would promote sagebrush control as well as maintain and improve rangeland conditions. Sheep will also control big sagebrush if allowed to graze the bush in late fall. INT(162,163).

28. Livestock grazing capacities, needed yearly adjustments in stocking, and range readiness can be predicted for mountain grasslands in Montana. These predictions are made through the use of May precipitation and the air and soil temperature data during the month of June. Results indicate that range readiness can vary from year to year by as much as 26 days and readiness lags 12 days per every 1,000-foot rise in elevation. Readiness on northeastern exposures is one week later than on the southwest exposures. This information is essential to the perfection of guides for resource managers. INT(170).

29. Deer mice and pocket mice in Utah greatly increase the first 2 years following pinyon-juniper type conversion to herbaceous species. This was especially true for all small mammals taken collectively where slash was windrowed versus slash dispersed throughout open areas. Complete slash removal—burning or physical removal--resulted in a lower small mammal population. Numbers declined sharply after 2 years but still remained higher than before treatment. INT(152).

30. Saw-palmetto, the major shrub on 18 million acres of land in Florida, reduces the availability of forage and makes handling of cattle difficult. It also contributes to fire hazards. Crosschopping and root plowing treatments have now been developed by the Southeastern Forest Experiment Station to eliminate or greatly reduce the difficulties associated with saw-palmetto. The end result is a greatly increased forage yield, and also a reduction of fire hazard to pine stands. SE(169).

31. "Wildlife in your backyard" is the desire of many residents of the eastern megalopolis. Scientists at the Pinchot Institute of the Northeastern Forest Experiment Station, Amherst, Massachusetts, have devised schemes to produce wildlife in suburban yards. Alternative planing schemes were developed for various areas of the United States. This effort also explains the increased wildlife diversity as the plantings mature. NE(183).

32. Deer populations remain low over much of the forested parts of the Ozarks compared to most eastern forest. Research data by the Southern Forest Experiment Station indicates that the scarcity of high-quality winter browse, especially the more nutritious evergreen species, is probably one of the most important factors responsible for the low populations. Japanese honeysuckle, one of several shrubs studied, appears to be the almost ideal plant to provide increased production of a highly nutritious winter food. SO(179,206).

33. Time-lapse photography is a valuable technique to replace observers in evaluating animal uses of wildlife habitat. The research staff of the Rocky Mountain Forest and Range Experiment Station at Tempe, Arizona, developed a time-lapse camera for use in biological research. The apparatus was constructed from a super 8-mm movie camera activated through a solenoid controlled by an electronic timer and a photo cell for daylight operation. Timing intervals can be varied from 1/2 second to 60 minutes. Film lasts 25 days when operated at a 5-minute interval and 12 hours of daylight. Cost of the apparatus is approximately $165. RM(171).

34. Livestock ponds provide water for livestock and also provide waterfowl habitat which greatly supplements the diminishing wetlands of the Northern Great Plains. Over a 5-year period, 53 species of waterfowl and shorebirds and 47 other species of birds were found to have used 12 ponds in western South Dakota. The average pond produced 35 young puddle ducks. Extending the sample data to the 90,000 ponds in western South Dakota, a potential production of over 300,000 puddle ducks a year can be projected from this area alone. RM(160).

35. Three-fourths of a shrub plant in the cold desert of western Utah is below ground the remaining fourth is above ground. However, the ratio of below ground to above ground annual production is considerably different in that nine-tenths is below ground and one-tenth is above ground. Consequently, grazing practices that result in a reduction of below ground parts has an exponentially reductive effect on above ground parts. INT(166).

36. Dense pole stands of northern hardwoods do not produce much food and cover for wildlife, particularly game animals. Small clearcut openings throughout these pole stands will provide abundance and variety of herbs, grasses, and shrubs that supply vital nutrition for game animals and desirable niches for nesting and brood-rearing sites of ruffed grouse and woodcock. Careful location of openings on shallow soils, in frost pockets, and in areas of poor or excessive drainage, can delay their return to tree cover for 20 to 30 years. NC(699).

2. Characteristics and forage and browse values of plants

37. Fertilizer, according to laboratory tests at Ft. Collins, Colorado, not only increases the amount of Sherman big bluegrass foliage, but also develops the plants resistance to being pulled up by grazing livestock. This increased anchorage is due to increased root development when fertilizer is applied to the plants. Livestock commonly pull up the entire Sherman big bluegrass plant when grazing and reduce subsequent forage yields and grazing values. RM(194).

38. Herbaceous plants that grow profusely when moisture is available and provide low-quality roughage when needed for emergencies are highly desired in the Southwest to reduce gully erosion and also provide emergency feed. Tanglehead (*Heteropogon contortus*), according to the scientists at Tucson, Arizona, is a warm-season perennial bunchgrass native to the warm arid Southwest proven to be such a dual purpose plant. RM(210).

39. Carpetgrass, common to the pine forests in central Louisiana, is highly preferred for cover in firebreaks to help prevent the spread of wildfire. Carpetgrass tends to form pure stands with increased utilization; forage yields decrease, however, when utilization exceeds that known as moderate use. Research results indicate that carpetgrass stands must be utilized or they become weakened allowing other plants to invade and reduce the effectiveness of the firebreak. Consequently, cattle grazing can be used to protect the pine stand. SO(211).

40. The International Shrub Symposium, an interagency effort largely financed and administered by USDA, Forest Service Intermountain Forest and Range Experiment Station, Ogden, Utah, was the first endeavor to confer on genetic improvement of shrubs. Shrubs offer tremendous potentials for man's benefit in the arid and semiarid lands. The symposium

focused on the use and biology of shrubs. The 500-page proceedings of the symposium, published by the Forest Service, covers the "Continental Aspects of Shrub Distribution, Present Utilization and Potential for the Future" with special emphasis on "Genetics" as probably one of the most neglected areas in the biology of shrubs. INT(202).

41. Topping of shrubs on three deer winter ranges in Utah increased twig growth over a 4-year period providing increased browse for deer. The results of the Intermountain Forest and Range Experiment Station indicate there were no detrimental effects on shrub longevity or increases in damage from insect attack or disease. Shrub topping can be done at costs varying from $9 to $21 per acre depending on density of the shrub stand. INT(190).

3. Livestock and grazing management

42. Sheep in southwestern Montana trample from 50 to 60 percent of the herbage used while grazing. Resource managers must allow for that destruction when determining sheep grazing capacities or overutilization will occur. The degree of trampling varies with season and plant species. Consequently, management can alleviate losses due to trampling by grazing the various plants when the plants are least susceptible to trampling damage. INT(222).

43. Mule deer, common to the open woodlands, do well on a year-round diet made up almost exclusively of woody material. Scientists at Tempe, Arizona, also learned that the diet dominated by browse species was highly related to the availability of the various woody plants. The selection was dominated by birchleaf mountain mahogany and oaks which were adequate nutritionally to sustain deer during all seasons. Hence, habitat management should strive to maintain these two important shrub species and to prevent their overutilization. RM(215).

44. Beef production was considerably higher under a continuous yearlong grazing management system than under rotated-seasonal grazing management in the annual-plant type rangelands of California. The addition of sulfur-fertilizer only slightly improved livestock performance but considerably increased grazing capacity. Beef calf and cow weight performance on the perennial bunchgrass rangelands of California were slightly better under rest-rotation grazing management compared to yearlong grazing management systems. This rotated-seasonal grazing management required shifting of use to various times during a year. PSW(218,234).

45. Cost reductions can be achieved by thrice-weekly rather than the more common daily feeding of a protein supplement for cow-calf operations on slash pine-bluestem ranges in Louisiana between November 1 and May 31. This is the conclusion from research on cattle running yearlong from 1964 to 1970 on the Palustris Experimental Forest in central Louisiana. SO(231).

4. Integrating other uses and wildlife

46. Rock-raked windrows, commonly associated with land manipulation in northern Minnesota, are inviting to black bears as winter den sites but are conducive to almost twice the heat loss compared to earthen dens. The heat loss could jeopardize winter survival. Overwinter survival is dependent upon (1) the food supply available in summer habitats, and (2) the degree to which the bear must reduce its metabolic rate and weight loss. The insulating effect of the den greatly influences the latter. NC(246).

47. Sportsmen must realize that good wildlife management will be practiced on private lands only if the landowner develops an interest in wildlife and cooperates in wildlife habitat management. Landowners must expect an income from the use of their wildlife habitat before they will be willing to consider that resource in their ranch management plans. But hunter-landowner interests and attitudes differ, as was detected in comparing the problems connected to "commercial" hunting in South Dakota to those in Texas. Public hunting areas, that is, National Forests, greatly decrease the demand for commercial hunting as associated with private land unless the private landowner provides extra services. However, the implementation of a federal land use fee would greatly favor "commercial" hunting in South Dakota. RM(249).

C. Improving Social and Amenity Values

1. Environmental amenities—landscapes and open space

48. Attractive and pleasing landscapes are an important amenity resource. Americans are more interested than ever before in protecting and managing their visual resources. However, there are not good methods for inventorying, analyzing, or communicating about landscape resources. Research in landscape design is producing some answers. Six factors of recognition and six compositional types have been identified. A method for inventorying landscapes has been developed, and it has been tried out in the Lake Tahoe area. An approach to evaluating landscapes, including three esthetic criteria, also has been proposed. These developments will help resource managers do a better job of considering impacts on the visual resource of various land management alternatives. PSW(257).

49. An aid to landscape resource analysis and management is the use of computerized drawings of ground forms and vegetation. Two computer programs have been developed for this use. They generate perspective views of landscapes that can be used to predict effects that proposed changes would have on the visual resource. PNW(256).

50. Preference for urban scenery versus nature scenery has been tested in carefully controlled laboratory tests. University students were tested with selected color photo-slides. Scenes of nature were greatly preferred over urban scenes. The research also showed that people generally prefer scenery that includes variety and visual complexity. NC(255).

51. Related laboratory studies showed that, in addition to complexity, mystery and coherence contribute to people's preference for a scene. Coherent scenery "makes sense" to people. A landscape has a quality of mystery if it promises to reveal something new or interesting further on, over a hill or around a bend for example. NC(253).

52. Cemeteries can be valuable open space environmental resources—especially in urban areas where open spaces are scarce. A study of cemeteries in the Boston area showed that they are being used for bird study, squirrel feeding, picnicking, ball playing, kite and model plane flying, driving, bicycling, walking, jogging, photography, and other recreational pursuits. With planning and careful management these areas can be used as open space areas for the living without losing sight of the original purpose, respectful interment of the dead. NE(258).

2. Environmental amenities—wilderness

53. Wilderness provides unspoiled natural environments and opportunities for a unique kind of primitive recreational ex-

perience. Visits to wilderness areas are growing rapidly and, in certain places, threaten the integrity of some wilderness areas. Data collected from visitors to four areas have been used to identify important elements of wilderness management. Visitor satisfaction is related to (1) intensity of use, (2) types of encounters with other visitors, (3) how visitors are spread over space and time, and (4) destructive behavior by visitors. Solitude is important; "crowding" reduces visitor satisfaction. Large groups cause loss of satisfaction for others. Privacy at campsites also is important. Littering is very objectionable. There are conflicts among those using different means of travel (hikers versus horsemen for example). This research provides a basis for defining wilderness carrying capacities and for developing wilderness management techniques. INT(265).

54. Research has led to a proposal that special mathematical models be developed to predict the travels of individuals and groups in wilderness areas. These models, operated in computers, will be used to test alternative approaches to wilderness management. NC(261).

55. In some cases the Forest Service and The National Park Service require that visitors obtain permits before entering wilderness areas. Data on visitors and visits, obtained from the permits, are useful in producing a factual record of visitation. These data can help improve wilderness management. A computerized system is available for analyzing the data. PSW(260).

56. Fire always has been an important ecological factor in many wilderness areas. Considerable meticulous ecological research has revealed the history of natural fires in the Boundary Waters Canoe Area of Minnesota. Current studies are tracing the course of natural plant succession after the Little Sioux fire of 1972. In the second year after the fire, hardwood trees were regenerating very well and the area also had a large crop of blueberries and abundant populations of small mammals such as voles, mice, and chipmunks. NC(259,266).

57. Research on visitors to the Boundary Waters Canoe Area (BWCA), a unique part of the National Wilderness Preservation System, has shown that large groups of visitors have significant impacts on the wilderness. Large groups accounted for much of the total use in the BWCA. The impact of the large groups on the environment and on the experience and satisfaction of other visitors is greater than that of an equal number of people traveling in small groups. Based on the results of this research the Superior National Forest recently changed the regulation on maximum party size down from 15 to 10 persons. NC(263).

3. Managing recreational opportunities

58. Camping is one popular form of outdoor recreation. In recent years there have been great increases in family camping and in investments in campgrounds. Better information on the family camping market benefits private campground businesses and governments that operate public facilities. A nationwide survey has shown that there are 12.6 million active camping families in the U.S.A. and 9 million inactive. The inactive ones include 3 million who have temporarily dropped out of camping and 6 million former campers who are permanently out of the market. There are 10 million families who do not camp but are potential campers. Most of the families who are potential are in the Northeast and North Central regions of the country and earn $7,000 to $10,000 a year. NE(280,281).

59. Privately owned commercial campgrounds have become a major supplier of family camping opportunities in recent years. Research is helping to identify criteria for successful development and operation of commercial campgrounds. Much has been learned from an analysis of 182 campgrounds in New Hampshire between 1964 and 1971. Detailed cost and return records were available for 46 of the businesses. Only 78 of the 182 campgrounds were financially successful. The successful businesses were larger than the average, had over two-thirds of their sites with utility connections for travel trailers, had higher average operating costs per site, had been in business for several years, had customers who stayed more days per visit, served about four times as many customers as the unsuccessful campgrounds, and generally had more businesslike management. NE(283).

60. Skiing is an important and rapidly growing recreation industry. In the 11 Western States, for example, skiing grew from 1.4 to 4.3 million visits between 1956 and 1964. Pressure for additional ski developments is felt by both public resource agencies and private investors. A statistical study of skiing in California and western Nevada provides some clues to decisionmakers. Use of individual ski areas decreased as the price of day tickets was increased. Use increased as lift capacity increased. Nearby ski areas tend to complement one another. That is, all of them have more business because there is more than one development within a reasonable area. The statistical model used in this study is one tool for analyzing the market opportunities for proposed new ski areas. PSW(279).

61. Many Western areas that offer recreational opportunities, such as fishing, boating, and hunting, do not have enough trees and shrubs to provide shade, screening, windbreaks, traffic barriers, or pleasing landscapes for visitors. Research in California is finding better ways of establishing and growing trees and shrubs in difficult areas, such as at reservoir sites. Methods include irrigation, fertilization, mulching, selection of adapted species, and horticultural care of trees and shrubs. PSW(286).

62. In the southern Appalachian forests of the Southeast, tests have been made of visitors' preferences for features of developed campsites. The location of the comfort station was the most important factor people considered in selecting a campsite. Shading was next most important with moderate shade preferred over either full shade or little shade. Accessibility of drinking water also is important. Most campers want to be 50 to 100 feet away from community hydrants. Research findings also provide clues about management of grass and shrubs, arrangement of facilities, and the importance of fairly level but well-drained campsites. SE(272).

63. Canoeing has become an important recreational use of many rivers. Eroding streambanks on Michigan's Pine River, being considered for stabilization to reduce silting and improve fish habitat, generally were considered a natural part of the scenery by canoeists. Canoeists, saw a need for preserving the river in a nonresidential and noncommercial natural state and for providing more well-planned facilities, such as picnic and camping areas and toilets. Delicate ecological sites probably will have to be protected from canoeists. Present users of the river objected to litter and to the feeling of "crowding" on the popular river. NC(292).

64. Data on forest visitors' preferences, opinions, and attitudes can improve planning and management of recreational opportunities. When collected and analyzed according to scientifically developed procedures, this information can help managers do a better job of serving the public as well as protecting the resources. Factual data can be substituted for inaccurate preconceived notions. For example, analysis of users of campgrounds on the Superior National Forest showed that some campgrounds were used six times as much as others. Campers strongly prefer sites near water and with a scenic

547-679 O - 74 - 2

view. Other preferences revealed by the research were: Fishing opportunities nearby, uncrowded conditions, being away from main roads and towns, and small campgrounds (fewer than 15 campsites). NC(284).

65. Nearly a thousand references have been brought together in an annotated bibliography of articles on the human aspects of fish and wildlife conservation. PNW(289).

66. Interviews with fisherman in New York State showed that the most important factors of successful fishing trips were: Water quality, natural beauty, and privacy or solitude while fishing. Number and size of fish caught were less important. Facilities were not important at all. Fisheries management should include attention to the total environment and not to fish stocking only. NE(287).

67. Litter is a major environmental problem. Litter cleanup costs are high and are a burden on the public. Scientific study has led to development of an approach to litter control that (1) capitalizes on normal forest visitor behavior, (2) reduces litter levels greatly, and (3) cuts costs dramatically. The research showed that the traditional approaches, such as appeals and litterbags or threats are not very effective. The new system uses the voluntary help of children and givens them small rewards, such as Smokey Bear symbols, as incentives. It is a very effective, low cost, cleanup method. Indications are that it helps teach the children antilitter attitudes that continue after leaving that forest site. PNW(269-271).

68. Recreational use of forests is not easy to measure or to sample. Yet managers need accurate counts or estimates of use so that they can plan and manage more effectively to serve the public. Prior research has made substantial contributions to scientific methods of estimating various kinds of recreational use in the forests. Recent studies have developed an easy and inexpensive way of collecting and analyzing data on the home or origin of visitors. This method uses self-registration, ZIP codes, and computer data processing. SE(278).

69. Recreation research scientists have evaluated an electric eye counter for sampling traffic on a wilderness trail. These counters show promise for cutting costs of sampling trail use, but the equipment needs to be refined and tested further before being widely used. SE(277).

70. Estimating use of waysides and observation sites is difficult because of the short time spent by most visitors. Methods useful for other kinds of sites, such as campgrounds, don't work. A pilot study on the Pike National Forest resulted in a technique for solving this difficult problem by using an observer for sampling the visits and correlating that sample with automatically recorded road traffic data. SE(276).

71. Results of some years of research on methods of sampling and estimating recreational use have been made more readily available to users in a manual that provides four computer programs in Fortran IV language. SE(294).

72. Scientific study of forest visitors' preferences, attitudes, and opinions usually requires the use of questionnaires. These tools of social science research must be carefully constructed and used. Forest Service scientists have developed an annotated bibliography of nearly 200 references on the design and use of questionnaires. PNW(290).

73. Recreation planners feel the need for estimates and projections of demands for recreational opportunities. Twelve demand surveys, made in connection with state recreation plans, have been analyzed. Ten guidelines for strengthening demand estimates were formulated and outlined. There is a need for establishing uniform methods and sharing data among States. PNW(268).

74. Some current recreation policy issues have been analyzed. The results show that there is an urgent need for development of wide variety of recreational opportunities along the full continuum from concentrated to dispersed typed of activities. Too little attention is being paid to the many needs in the middle of the continuum—such as for semideveloped hiking and camping trails or off-road vehicle trails. The two ends of the continuum, pure wilderness and car camping, have received most of the attention. There is an urgent need for more, and a wider variety of, dispersed recreation opportunities outside formally designated wilderness areas. WO(285).

75. The huckleberry fields that provide economic and recreation benefits to thousands of residents of Oregon and Washington are dwindling in size and productivity, and effective management techniques for conserving and developing the huckleberry resource are urgently needed. A summary of available information has been prepared that includes descriptions and an identification key for the 12 northwestern huckleberry species. It also reviews current knowledge on blueberry management in eastern North America. This information is useful for planning research and developing management practices for the protection, culture, and use of the wild huckleberry resource in the Pacific Northwest. PNW(700,701).

D. Trees to Enhance the Environment

76. One of today's serious environmental problems is how to deal with excessive noise created by more and larger vehicles on highways and city streets. Studies in the past 5 years conducted by the University of Nebraska and the Forest Service have demonstrated that 20- to 50-foot-wide belts of trees and shrubs, with the edge of the belt from 20 to 50 feet from the center of the nearest traffic lane, can reduce the noise level by one-half. It is best to use shrubs 6 to 8 feet tall next to the traffic lane with backup rows of trees 15 to 30 feet tall. RM(702,703).

77. Trees and shrubs when properly placed in relation to the source of the noise can be effective in reducing outdoor noise level. Recent studies jointly conducted by the University of Nebraska and the Forest Service have shown that tree covered land-forms are considerably more effective than are land-forms alone or trees alone. To reduce noise from high speed car and truck traffic, the land-form should screen the noise source from view at the location to be protected. Trees, shrubs, and grass should be planted on and around the land-form to extend the range of protection and further reduce the noise level. RM(704,707).

78. The climatic and edaphic extremes characteristic of the Great Plains severely restrict the kinds of trees that can be used for environmental protection purposes in this region. Seven different seed sources of Japanese larch were grown at 13 locations in 8 States including eastern Nebraska. After 13 years Japanese larch of high elevation origins grew the best in the Great Plains. This finding is important because it adds a new species to the rather limited list of adapted conifers for environmental plantings in the Great Plains. RM(705).

79. Many older shelterbelts in the Great Plains have serious management problems resulting from mistakes in spacing and arrangement of the trees when they were planted. To avoid these problems, a proposed check list for planning windbreaks and shelterbelts was developed which includes consideration of the purpose of the planting, the best designs, and how to select the species to be planted. RM(706).

80. There is an increasing interest in the proper utilization of medicinal plants in West Pakistan since drugs of plant origin represent an important export to the United States. Over 635 flowering plants of pharmacological value have been identified recently in the Dir and Chitral forest ranges of West Pakistan by a PL-480 supported research program. A concurrent quantitative survey of the availability of medicinal plants suggests that for certain plants it is now feasible to make commercial collections for export to the United States, as well as to other nations. WO(708-712).

E. Improving Environmental Quality Through Fire Management

1. Fire prevention, hazard reduction, and prescribed burning

81. A specially designed series of fire prevention lessons were prepared for preschool children in the Headstart Project in Riverside County, California. The effects of this effort were most successful in meeting cognitive goals but positive changes in curiosity about fire and attitudes about fire were less apparent. It is important that children learn to cope with and avoid dangerous fire situations as well as to learn the prevention of wildfires. PSW(632,634).

82. Efforts are being made to involve other social scientists with the problem of man-caused forest fires. The intricate interrelationships with man, fire, and man's natural and social environments need to be studied in more detail. These efforts should increase the opportunity for involving competent scientists in the solution of man-caused fires. PSW(633).

83. Differences in organizational structure and approaches to community relationships effect management success. Basic principles must be followed in planning and executing forest fire prevention programs or the processes themselves become barriers. Six guidelines may assist the communicator to overcome some of the barriers. They are: (1) audience analysis, (2) selection of objectives, (3) message design, (4) media selection, (5) selection of communicator, and (6) evaluation. Strategies of organization and work accomplishment must be identified if we are to achieve forest fire reduction goals. SO(644), PNW(631).

84. The Southwest Interagency Fire Council has emphasized the importance of considering forest fuels and fire planning in the early stages of land-use and multiple-use planning. Such specialized operations as smoke management must also be a part of the early land-use decisions if they are to be effective because there are tradeoffs between the smoke of planned fires and wildfires. INT(622,624).

85. The weight of shrub fuel varies from a few hundred pounds to 50 tons per acre, but living material larger than 1/2 inch in diameter is seldom consumed in wildfires. Dead fuels vary from below 25 percent of the total volume to nearly 50 percent. Fuel arrangement is more important than fuel weight in determining fire behavior characteristics such as rate of spread, burning time, and intensity. Fuel moisture is rated heavily for fire-danger rating because it controls the current flammabilities of fuel. Ash content of shrub species also has effects on the pyrolysis process and affects the way a shrub stand will burn. INT(641).

86. Ten of 18 National Forests in the Intermountain Region have applied a fuel break concept recently developed on the Wasatch front. Atrazine and simazine are used to control the flammability of cheatgrass. Up to 50 percent reductions in rate of fire spread are achieved. PSW(635).

87. Fire spreads nearly 5 times as rapidly through dry annual grasses and forbes than on plots where these same plants are growing in combination with creeping sage stands. Because of its low stature, fuel volume, low fire spread, low resistance to fire control, and adaptability to site, creeping sage appears to be a good candidate for planning in areas where hazard reduction is a goal. PSW(642).

88. Perennial shrubs survive adverse conditions by maintaining a reservoir of dormant meristems or buds. The origin, development, dormancies, and photoperiodism of these buds are important in the management and maintenance of fuel breaks or other shrub treatments for grazing or watershed purposes. INT(623).

89. Fenuron, tandex, and picloram are herbicides used to maintain control over scrub oak regrowth. Fenuron and tandex are nonselective sterilants which eliminate most plants while picloram has little apparent effect on grass cover. After a 30-month test, good stands of grass developed. This is important in the maintenance of fuel breaks or in range management considerations. INT(643).

90. A random orientation line intersect sample line is best for measuring residue in cutover areas where topography and logging create orientation patterns. Areas where terrain is more gentle can be sampled by a unidirectional pattern. Estimates of volume within plus or minus 15 to 20 percent are possible. Higher intensities can improve precision. PNW(638).

91. Five fuel types in the Lake States Region have been described and displayed in graphical ADP generated form. Crown weights and crown volumes are shown. This method will help managers describe changes in fuel complexes. NC(647).

92. A set of alignment charts developed for the Lake States estimates fuel loading of downed dead wood using a planer intersect technique. Fuel inventories can be improved. NC(645).

93. A preattack area planning guide developed for the Pacific Northwest tells how to assemble, store, retrieve, and update preattack data within the total resource inventory system. PNW(630).

94. Surveys in 13 southern States indicate that over 2 million acres were prescribe burned annually during the past 8 years. Prescribed burning is a vital forest management tool in the South. It plays many roles such as, fuel hazard reduction, seedbed and site preparation, control of undesirable species, wildlife habitat and grazing improvement, and certain disease control. SE(628,636,640,646).

95. A new device for safely, electrically firing prescribed fires has been developed. Design, operation cost, and diagrams are available. Reliability improves efficiency and safety. NC(639).

96. Smoke from wildfires in the Florida Everglades restricted visability 68 percent of the time for 17 days of April 1971. Aircraft operations were reduced by 12 percent. However, maximum concentrations of particulate were well below national standards for air quality. This indicates that smoke from large fires has less effect on air quality than it does on certain of man's activities such as aircraft operations. SE(648).

97. The formation and diffusion of particulate matter from prescribed fire are being thoroughly investigated, but to date, there is no evidence to indicate that air quality has deteriorated more in areas where prescribed fire is used extensively than it has where prescribed fire is rarely used. Smoke from prescribed fire can be traded against the consequence of wildfires and large smoke plumes. SE(625-627,629,637).

2. Fire management methods and systems

98. Rigorous testing of the CL-215 airplane, the first aircraft built specifically for air tanker work, revealed that (1) the efficiency (feet of adequate fire line built per gallon of retardant delivered) of the CL-215 is approximately the same as that of other air tankers tested, (2) the capacity of the CL-215 is intermediate in the range of sizes available, (3) the CL-215 is less flexible than air tankers having multiple gating or trail gating, and (4) the CL-215 has the advantage of a water scooping mode and can use an onboard water thickening system. Gum thickened retardants show a 17 percent superiority in drop characteristics over water. The onboard mixing system using gelgard can increase the drop characteristics by approximately 10 percent. Thickened retardants allow drops from higher altitudes, thereby increasing safety. The test of the CL-215 could set precedence for testing new candidate air tankers. WO(661).

99. Florida Division of Forestry achieved successful drops with the DeHaviland beaver aircraft used as a water tanker in Canada. Forest Service studies showed that drops had good conformation comparable to aircraft such as the TBM. These small aircraft provide specific capabilities for unique situations. SE(656).

100. Increasing the amount of either ammonium sulphate or ammonium phosphate fire retardant on ponderosa pine needles or aspen excelsior fuel beds causes a reduction in rate of spread and rate of weight loss and an increased amount of residue. At all treatment levels ammonium phosphate is more effective in decreasing the rate of weight loss and increasing residue than ammonium sulphate which has little effect on residue. Weight loss and especially residue quantify glowing combustion. These relationships serve as standards for comparing the effects of other chemicals or proposed forest fire retardant additives. INT(655).

101. A commercially available hydraulic seeder used regularly by the Oregon State Highway Department for roadside maintenance gave satisfactory results for mixing and spreading powdered and liquid retardants. A two-man crew was able to apply large volumes of retardant quickly to aerial and ground fuel in "hard to reach" locations. This development makes it more efficient to apply retardant with ground tankers for fire control or for treating hazardous areas and borders of prescribed fires. PNW(662).

102. Helicopters and helibuckets are proving useful, additional tools to the Alaska fire program. Tactical and logistical problems of controlling remote fires in Alaska are tied to the vastness of the country. Consequently, fixed-wing aircraft have traditionally played an important role. Now helicopters are adding to their capability particularly since there are readily available sources of water in many of the fire fighting areas. PNW(659,660).

103. A computer program (FIRDAT), written in Fortran-4 developed for use in Regional data processing offices, will compute components and indexes of the National Fire-Danger Rating System for each day of weather. This permits the examination of historical weather records and an analysis of fire danger. RM(654,657).

104. A computer method of displaying information in the form of a 3-dimensional histogram was developed. This method can be used for a variety of data and provides the user sufficient information to examine three related variables. PSW(658).

105. A theoretical study of precipitation effects on dead cylindrical fuels indicates that rainfall rate and amount have little effect on changes in fuel moisture but duration of precipitation is very important. This information, when confirmed by field tests, will aid development and refinement of national fire-danger rating. RM(653).

106. A revised Fire Weather Observers Handbook was issued and distributed nationally. The handbook gives instructions for location layout of fire weather measuring stations. It is important that all fire weather stations have acceptable observations stations and some degree of uniformity. INT(652).

107. Moisture content of tobosa grass can be predicted using wind speed, relative humidity, and air temperature. Accurate predictions of moisture content are necessary for judging fire danger and predicting expected fire behavior. PSW(651).

108. An inexpensive 31-day battery-operated weather station has been developed to use in lieu of more expensive presently available commercial models. Inexpensive radiation shields for thermisters and thermocouples were also developed. These are helpful devices for recording weather in remote stations where frequent observations are difficult and for instrumenting many biological research and management activities. PNW(649,650).

3. Forest fire science

109. Minimum temperatures for pilot ignition were established as 340°C with a pilot flame using ponderosa pine needle samples. Other practical tests using lighted matches dropped into southern slash pine litter identified ignition probability for southern fuels. Ignition delay times and temperatures are necessary inputs for fire modeling. SE(663), INT(678), PSW(664).

110. Central temperature history and mass loss curves are correlated with initial diameter and are practically independent of environmental conditions. Temperature histories of fuel elements are important in the understanding of ignition and fire spread. SE(665).

111. A newly discovered dehydration product of levoglucosan has been shown to be important in the tar formation process when both the basic and acidic fire retardants are used. Both types drastically lower the levoglucosan yield but basic retardants have a much greater effect on the tar yield and on the flammability of cellulose. With acid retardants, the decrease in levoglucosan yield is accompanied by an increase in the dehydration product identified. PSW(670).

112. Progress is being made in new technology for studying the chemical breakdown of hemicellulose, a major plant material. Identification of products produced in the presence of different flame retardants allows scientists to describe the mechanism of fire suppressants. INT(676).

113. Through new computer techniques it is now possible to simulate the combustion of cellulose and other C-H-N-O fuels. The technique will compute combustion products for arbitrary fuels for specified combustion conditions. PSW(677).

114. The intensity of a combustion wave moving through a forest homogenous fuel array is an important but poorly defined concept of fire behavior. A concept of reaction intensity, which is related to the rate of fuel consumption per unit of basal area within the combustion zone has been developed. This knowledge is necessary for the characterization of fire intensity in terms that are useful for predicting the behavior of a spreading fire. INT(668), SE(673).

115. The National Fire-Danger Rating System uses a new fire spread model developed specifically for a wide range of

wildland fuels. This model is persented from conceptual stage through evaluation and demonstration and applied to hypothetical fuel models. INT(674).

116. None of five experimental and theoretical fire spread models tested in Arizona oak chaparral gave adequate, absolute predictions. However, a statistical model using esentially the same inputs, but weighted with chemical coefficients, accounted for 81 percent of the variation in rate of spread. The revised model provides useable guides for predicting absolute values of fire spread in Arizona oak chaparral. RM(672).

117. Fireline supervisors and firefighters have been given new aids to help understand the concept of fire environment and the specific processes that influence the formation of fire whirls. Fire whirls are dangerous to firefighters and make fire control difficult. PSW(666,667).

118. Fire managers have gained additional useful information from the analysis of wildfires. A study of the Air Force Bomb Range Fire in North Carolina indicated that 14 miles of fire spread in 7 hours was a direct result from gusty winds, adverse wind profile, and passage of a cold front, along with heavy uniform loading of fine fuels. Byram's equation for predicting fire blowup was useful in the analysis. The importance of fast initial attack was demonstrated by a study of the 1971 northern Minnesota fire which burned almost 15,000 acres in a 3-day period. The fire reached 60 percent of its final size in the first 6 hours. The studies made on two state protection districts on the Clark Forest in Missouri identified factors that contribute to fires, particularly on multiple fire days. SE(671,679), NC(669,675).

F. Improving Insect and Disease Control

1. Detection and evaluation

119. Aerial color and infrared photography are becoming increasingly valuable for detecting and evaluating forest diseases. Recent studies have identified "signatures" that distinguish trees affected by air pollution, Dutch elm disease, oak wilt, ash dieback, white pine basal canker disease, annosus root rot, and *Poria* root rot. Combined with comfirmation by direct ground sampling, aerial photography promises to be a powerful tool for disease survey and a useful aid in research on disease epidemiology. NE(462,463), PSW(461).

120. Aerial photography provides a reliable means of monitoring the trend and spread of bark beetle infestations. Analysis of large- and small-scale photography now permits choice of the optimal photoscale to minimize interpretation error for damage surveys. A scale of 1:32,000, for example, is generally best for assessing tree mortality due to the mountain pine beetle in the Black Hills. PSW(312).

121. A number of remote sensing techniques are available for use in forestry. Some are presently too costly or do not provide satisfactory resource data. Applications include side-looking airborne radar (SLAR), multispectral scanning, and color and color infrared photography. The latter techniques have been most used by forest managers and protection specialists during the past 5 years. PSW(304).

122. The effect of gypsy moth defoliation on tree condition and mortality has not been fully evaluated. A series of preliminary predictive tables have been prepared by the Northeastern Station following a comprehensive analysis of a large mass of historical data. These give both annual and cumulative tree mortality rates and the condition classes of surviving trees for particular tree species according to the intensity and duration of defoliation. These tables require confirmation in current outbreaks, but they will provide initial means for predicting the potential impacts of gypsy moth defoliation in northeastern forests. NE(301).

123. Through the years, the airborne dispersal of insects such as the gypsy moth and beech scale has been regarded as a natural phenomenon which is beyond our control. It is not a purely random process, however. Recently developed atmospheric dispersion models for air pollutants can be integrated with a gypsy moth larval dispersal model to increase the accuracy of predictions of rate of spread, insect numbers, and resultant defoliation. A predictive model for larval dispersal is an essential tool in population assessment. NE(306,307).

124. A fast procedure was needed to permit individual examinations of large numbers of diseased Douglas-fir tussock moth larvae infected with either or both of two types of nucleopolyhedral viruses. A method was developed which utilizes an alkali solution containing nonionic hydrocolloid, hydroxyethel cellulose. This procedure permits counting of inclusion bodies in mixed samples and the determination of the density and ratios of the two viruses. PNW(308).

125. A description of distinguishing features of important needle diseases of Scotch pine Christmas trees has been developed. Christmas tree growers now will be able to detect these troublesome diseases early, thereby allowing effective control measures. NC(468,469).

126. A new technique for the direct isolation from soil of the black root rot fungus was developed. A combination of wet sieving and culturing on a selective medium allows estimation of the propagule population of the casual fungus. This technique can be used in forest nurseries to determine the need for control prior to seeding. PSW(466).

127. Fruiting bodies of the endotrophic mycorrhizal fungus, *Endogone*, were described and a technique permitting detection of this fungus in the soil was developed. Sporocarps can be identified in soil samples by their form and bright red color after soaking and washing in an acidified acid fuchsin solution. This procedure worked well with *Endogone* associated with sugar maple and can aid in future biological studies of these beneficial symbionts. NC(464).

128. Improved survey methods are needed for predicting defoliation of Douglas-fir by the western spruce budworm in Oregon. Optimum size and allocation of egg and larval samples have been determined. Ranges of egg mass and larval densities have been related to categories of defoliation. These guidelines will improve the precision and reliability of budworm evaluation in the West. PNW(302).

129. Spruce-fir stands defoliated by eastern spruce budworm were compared for 10 years with stands protected by insecticides to determine the effect of defoliation on growth and mortality of balsam fir. Defoliation reduced growing stock by an average of more than 1,000 cu. ft. per acre. Most mortality and growth reduction occurred during the fourth and fifth years of heavy infestation. Suppression decisions should therefore be made prior to this point. An index has been developed to assist the manager in assessing the stage of damage in individual stands and the need for action. NC(300).

130. Outbreaks of the western spruce budworm in the central and southern Rocky Mountains are eventually terminated by natural causes. Analyses of life tables compiled for two generations at three locations revealed that considerable mortality occurred during fall and spring dispersal periods and during the winter months. To improve the accuracy of biological evaluations, eggmass surveys in late summer or fall should be followed by spring surveys when most of the larvae are in the fourth instar. This will permit determination of overwinter survival and improve suppression decisions. RM(305).

131. Determination of parasitism of western budworm can be a tedious task when it involves dissection or rearing. Newly developed radiographic techniques will permit rapid differentiation of normal and parasitized budworm pupae. INT(310).

132. Research is leading to the application of nondestructive techniques for detecting internal defects in trees and wood products. A mobile X-ray unit developed for field use looks promising for determining wood quality prior to harvest, for evaluating tree hazard in municipal and recreational areas, and for inspecting utility poles, pilings, and other wood products. Cooperative research with the University of New Hampshire indicates that wood decay can also be detected by measuring changes in resistance to a pulsed electric current. These techniques promise to be inexpensive and safe alternatives for nondestructive detection of defect. NE(460,470-472).

133. Light traps may be used to determine the seasonal occurrence, relative abundance, and flight activity of insects affecting cones and seeds. By this means, the flight habits of two species of pine-feeding loopers, *Nepyta semiclusaria*, and *Lambdina pellucidaria*, in the Georgia Piedmont region have been determined. This information will aid in the timing of control of these pests. SE(313).

2. Biology and understanding

134. *Ips grandicollis* aggregation behavior is influenced by volatiles produced by the host and the insect. The combination of host terpenes and frass from male beetles is one of the primary sources of attraction involved in mass attack and aggregation. Various environmental and physiological factors influence the response of male and female beetles to the attractant complex, however. These findings will aid in developing attractants for surveys and suppression of this insect. SE(383, 384).

135. Response to natural attractants varies among *Ips pini* populations occurring in different geographic regions. Tests conducted in California, Idaho, and New York revealed differences both in pheromone production and behavioral responses. These results dramatize the necessity for considering pheromonal variability in survey and control programs involving the use of attractants of widely distributed pests. INT(351).

136. The pathogenicity of entomogenous fungi varies with temperature and relative humidity. Laboratory tests with adult southern pine beetles indicated that *Beauveria bassiana*, *Aspergillus flavus*, and *Fusarium solani* were most pathogenic at temperatures of 15-25° C and relative humidities of 55-94 percent. SE(360).

137. Tests of 10 strains of naturally occurring bacteria against southern pine beetle larvae revealed that *Bacillus cereus*, *B. thuringiensis* var. *thuringiensis*, *B. thuringiensis* var. *kenyae*, *Pseudomonas aeruginosa*, *P. fluorescens*, and *Serratia marcescens* were all pathogenic to the insect. Virulence differed between strains of *S. marcescens*. SE(359).

138. Fungal symbionts in a glandular structure (mycangium) in female adult southern pine beetles are thought to aid nutrition of larvae developing in the phloem tissue. An undescribed *Sporothrix* and a basidiomycete are the predominant fungi, although other microorganisms occasionally occur in the mycangium, also. SO(321).

139. Concentrations of inorganic constituents, water, and ash in the inner bark of loblolly pine vary seasonably and on different soil types. In North Carolina, ash, water, calcium, aluminum, manganese, and iron differed significantly among trees on three soil types. Most elements reached maximum concentrations in the fall, minimum in the spring. These findings may provide a partial explanation for the late season buildup of such pests as the southern pine beetle in the southern Piedmont-Appalachian Mountain region. SE(385).

140. Prey consumption and rate of development of the predator *Thanasimus undatulus* vary with the size and availability of mountain pine beetle brood. The larval period lasts from 56 to 68 days. In laboratory studies, 18 to 43 host larvae were consumed during this period, depending upon their size and numbers. This information will be useful in evaluating and predicting the effectiveness of known numbers of this predator under natural conditions. INT(316).

141. A model has been developed to predict the effects of three species of woodpeckers on endemic, epidemic, and pan-epidemic populations of spruce beetle larvae. Resultant predictions compared favorably with estimates made by measuring relative survival of larvae inside and outside of woodpecker exclosures. The combined predatory impact of the woodpeckers was least effective on epidemic populations of spruce beetles. RM(349).

142. Xylem resin of pines is closely linked with their resistance to tree-killing bark beetles. Three resistance mechanisms—preference through attraction, repellency, and synergism; antibiosis by both chemical and physical properties; and tolerance by healing and secondary resinosis--are active in this relationship. The level of resistance to bark beetle attacks can be increased through tree selection or breeding. PSW(374).

143. An epidemic of the spruce beetle in north central Colorado declined after 2 years. Reduced fecundity, resin flow, interspecific and intraspecific competition for food, predation, parasitism, dry inner bark conditions following woodpecker drilling, and subzero temperatures all contributed to the decline. RM(354).

144. Detailed investigations have been completed on the mating, egg-laying behavior, and brood development of *Ips pini* under controlled laboratory conditions. This information will be used in field and laboratory studies of the population dynamics and control of this pest. INT(373).

145. *Ips cribricollis* and *I. calligraphus* infestations are currently limited to felled trees in young Caribbean pine stands in northeastern Nicaragua. When these stands become older and more dense, it is anticipated that *Ips* and southern pine beetles may cause significant tree mortality. Managers should be able to identify these pests and be alert to the need for direct control if the problem arises. SE (391,392).

146. The identification of economic insect species and their life histories and behavior are important to understanding the structure and functioning of forest ecosystems in interior Alaska. A key has been developed and flight periods determined for major bark beetles associated with white spruce stands. This information will aid studies of insect populations and development of stand management techniques. PNW(323, 324).

147. Dwarf mistletoes reduce root growth as well as stem and crown growth of ponderosa pine. Roots of seedlings were considerably shorter with only about one-half the mass of roots of paired healthy seedlings when measured 2 years after experimental inoculation. Young pines infected with dwarf mistletoes apparently suffer a competitive disadvantage for soil moisture and nutrients which helps explain why infected pines grow so poorly. PNW(501,502).

148. Dwarf mistletoe in a single, isolated, overstory of ponderosa or Jeffrey pine can infest about an acre of pine

reproduction. The maximum distance of spread from isolated trees averaged 120 feet and was generally greatest in the direction of prevailing winds. Prompt removal of infected residual trees, left after fire or logging, is advised to avoid losses in the succeeding generation. PSW(520).

149. Light stimulates infection of dwarf mistletoe in Digger pine. Radicles of germinating dwarf mistletoe seeds failed to penetrate and become established in pine tissues when light was excluded. Light and photosynthesis of the young mistletoe plant appear to condition normal trophic responses of the expanding radicle. This is another step toward understanding the physiology of infection which ultimately will lead to new techniques for managing these destructive parasitic plants. PSW(538).

150. Basic biological experimentation clarified the requirements for infection of junipers by *Phomopsis* blight. In controlled environments, infection was limited to new foliage and occurred when wet, mild conditions prevailed. These findings will be useful in investigations of the epidemiology of this blight which is recognized as the most damaging nursery disease of conifers used for Great Plains' shelterbelts. RM(522).

151. New evidence indicates that the beech bark disease, caused by fungi infecting feeding wounds of scale insects, is well established and threatening most of the beech resource in the eastern United States. Beech is an important component of our forests for wood, recreation, and wildlife habitat. Research is needed to better understand and, eventually, control this disease. NE(548).

152. Brown spot needle cast is causing major defoliation of Scotch pine Christmas trees in north central United States. Control studies following epidemiological research show that one properly timed application of Bordeaux fungicide will protect most conifers from infection for an entire season. Red pine also was discovered to be susceptible to this disease. Research now underway will determine how best to prevent serious outbreaks in this important pulp and timber species. NC(516,517,550).

153. Differences between northern and southern isolates of the brown spot needle cast fungus have been found. Isolates from Scotch pine in the Lake States have different physiological requirements for growth in culture than isolates from longleaf pine in Mississippi. Furthermore, the northern isolates are most pathogenic to Scotch pine while the southern isolates are more pathogenic to southern pines. These indications of separate races of the fungus have important implications in tree improvement programs and on the development of control strategies. SO(497).

154. The cause of an aggressive pine needle cast of the Pacific Coast, *Lophodermella morbida*, has been identified and described. Evidence suggests that this fungus is endemic and limited to moist, moderate climates. It is particularly damaging in ponderosa pine planted in the westside Douglas-fir zone of Oregon. RM(556).

155. Information on anthracnose disease of eastern hardwood trees, especially American sycamore, black walnut, and white oak, was summarized in a new pest leaflet. This foliage disease is enhanced by moist situations. Use of resistant species or varieties is recommended. Removal of infected foliage and spring applications of Ferbam or Bordeaux fungicides are used for nurseries and ornamental trees. NE(481).

156. Identification of the orange-spored needle rusts of true firs in the Rocky Mountains by morphological means is now possible. The fir-huckleberry rust occurs on both current and previous season's needles and has rougher spores than the fir-fireweed rust. Proper identification is a necessary prerequisite to understanding these diseases. INT(507).

157. A *Coleosporium* needle rust known previously only on limber pine in New Mexico and on pinyon pines throughout the Southwest was found in epidemic proportions in high-elevation forests of Mexican white pine in Oaxaca, Mexico. This and other new information on the geographic and host range will aid in evaluating the potential hazard of this rust to pine forests. INT(523).

158. In sparse gypsy moth populations, the resting place of larvae greatly affects their survival. It was found that late instar larvae follow silk trails down the trees at dawn and relocate in protective niches at or near the base of the trees. They then ascend the same trees at night and resume feeding. This behavior protects them from several natural predators. NE(357).

159. Eighty-six aerobic bacterial pathogens have been isolated, characterized, and identified from gypsy moth larvae collected in sparse and dense populations. The most common pathogens were *Streptoccoccus faecalis*, *Bacillus cereus*, *B. thuringiensis*, Group C. *Enterobacter* types, and *Pseudomonas* spp. Only the first two were found to cause appreciable mortality in dense populations. Continuing studies will determine the significance of the bacterial complex in regulating gypsy moth population. NE(370).

160. The western spruce budworm normally passes through an obligate diapause, or dormant period. Pacific Southwest Station entomologists have been able to prevent diapause by altering the physical conditions in which first-stage larvae are reared. This will greatly increase the number of generations in a year and permit higher production of the insect for research purposes. PSW(353).

161. Fatty acids and oils affect the glucose metabolism and survival of western spruce budworm. In laboratory tests, oleic and linoleic acids and sweet almond oil prevented glucose metabolism in pupae but not in larvae. These acids were 15 and 3 times more toxic to larvae and pupae, respectively, than almond oil or mineral oil. This knowledge of biochemical changes induced by fatty acids may lead to new, selective methods of insect control. PSW(318).

162. Many parasitic wasps are highly selective in their choice of host insects. Specially designed experiments have shown that *Monodontomerus dentipes*, for example, differentiates between larch and red-headed pine sawfly cocoons. This confirms previous observations on the host specificity of this parasite. SE(331).

163. The lodgepole needle miner, *Coleotechnites milleri*, attacks and kills lodgepole pines in the western United States and Canada. Results of studies in Yosemite National Park on the biology and ecology of this insect, its natural enemies, environmental influences that affect population fluctuations, and field tests of insecticides have been summarized in a USDA Technical Bulletin. PSW(379).

164. Stands heavily defoliated in 1936-37 by the Douglas-fir tussock moth at Mammoth Lake, California, were studied to determine the incidence and extent of subsequent decay in top-damaged trees. Few decay organisms were isolated. However, old-top damage and wetwood were common. It was concluded that in eastside Sierra Nevada white fir stands defect is not economically serious where larger trees will be logged within 35-40 years after heavy defoliation and top damage. PNW(386).

165. Insects influence the flow of energy in aspen forests through their activities in defoliating, boring, girdling, gall making, and sucking plant juices. Their net impact is not well understood but in some cases may be beneficial. Greater knowledge of the role of insects in the energetics of an aspen

stand will improve management decisions aimed at increasing productivity and other resource values. NC(322).

166. Outbreak populations of the lodgepole needle miner in central Oregon occur primarily in topographic basins where lodgepole pine grows in pure stands and is the "climax" species. The insect is virtually absent in sites with high seasonal water tables, deeply developed soil profiles, and dense tree stocking. Similarly, populations usually decline abruptly in stands on well-drained slopes where lodgepole pine is seral or is growing with ponderosa pine. Degree of infestation thus appears to be influenced by a combination of environmental and tree physiological factors that vary significantly with different forest-site conditions. PNW(358).

167. The life history, behavior, and recent outbreaks of an oak leafroller, *Archips semiferanus*, have been described. This insect defoliates red oaks in northeastern Lower Michigan and Pennsylvania. The killing of scrub oak in Michigan favors the release of underplanted red pines. However, in urban, scenic, wildlife, and oak timber-producing areas, defoliation and mortality cause serious economic, social, and ecological losses. NC(387).

168. At least 40 species of weevils feed on trees and shrubs in North Dakota windbreaks. Of these, the red elm bark weevil, white pine weevil, poplar- and willow-borer, an ash seed weevil, the strawberry rooot weevil, and an acorn weevil are the most important. Knowledge of the life histories and behavior of these pests has been summarized in a Station Research Note. RM(356).

169. In the laboratory, elm spanworm eggs do not hatch at temperatures lower than 5° C. Optimum hatch apparently occurs at 19-20° C and 95-100 percent relative humidity. Mathematical models have been developed to predict the time of hatch of the spanworm in the field. SE(332).

170. Field and laboratory tests in Washington and Oregon revealed that high evaporative conditions reduced egg numbers and survival of the European pine shoot moth. This factor could limit the distribution of the insect in the semiarid western pine region. PNW(330).

171. Hybrid poplars planted on Ohio spoilbanks in 1951 are suffering decline and mortality from *Fusarium* canker. Isolation and inoculation trials confirmed the pathogenic nature of the fungus on 16 hybrid combinations. Adverse soils and too close spacing seemingly predispose the trees to *Fusarium* attack. *Fusarium* canker must be overcome if poplars are to be used extensively in reclamation projects. NE(484).

172. The *Sirococcus* pine tip blight was discovered on Coulter pines in forest nurseries in coastal northern California. This disease, found last year on Jeffrey pine, is a new problem in western United States. It has proved to be periodically troublesome to spruce and pines in Europe and eastern United States. Studies in California indicate that this fungus is inactive during winter, but development of fungicide treatments for warmer season application in nurseries is warranted. PSW(553).

173. To reduce the harmful impact of *Sclerroderris* canker disease to red pine plantations in the Lake States, planting stock must be disease-free and not planted near residual infected trees. Research on inoculum production and dissemination emphasized the hazard of local intensification within plantations. Prescribed burning appears to be a promising tool to remove infected trees before problem sites are replanted. Development of effective controls for this damaging canker disease is requisite to large-scale reforestation with red pine. NC(549,551).

174. Research at the North Central Experiment Station shows that shade favors growth of *Hypoxylon* cankers on aspen. Cankers receiving the smallest amount of available light in controlled experiments developed the most rapidly while those receiving full light grew the least. Variations in canker growth rate must be considered when evaluating impact from disease incidence surveys. NC(474).

175. An outbreak of *Diaporthe* canker in 1958 killed many yellow birch seedlings in the Upper Peninsula of Michigan. Infection and mortality were most severe on poorly drained sites and among the less vigorous seedlings. Although the mortality is a part of the natural thinning process, any mechanical thinning should favor disease-free dominants. NC(498).

176. Research on spore dissemination of the western gall rust fungus showed that dispersal of inoculum is limited to springtime in the Bessey forest nursery of Nebraska. Control of this pine disease in the nursery could be achieved with protective fungicides applied only in May and June. RM(521).

177. Tissue cultures of slash pine when subjected to attack by fusiform rust provide an ideal test system for study of host-parasite relations. Physiological, cytological, and biochemical studies now permit precise control over pathogenic interactions. The tissue culture technique could lead to more efficient and economical screening of potential chemical, biological, and genetical controls damaging diseases of southern pines. SO(488,510,511,560).

178. Evidence now implicates the existence of genetic variation within geographically distinct populations of the fusiform rust fungus. Differential responses were found among several pine progenies exposed to different geographic sources of the causal fungus. This information will aid in developing resistance to fusiform rust in pines for use in particular regions of the South. SO(554), SE(525).

179. Limb rust was observed on several pine species in northern Mexico. This rust appears to be caused by a fungus also endemic in ponderosa pine forests of our Southwest. Consequently, our research on silvicultural control should be directly applicable to forests in northern Mexico. INT(524).

180. Teliospore induction by the white pine blister rust fungus was determined to be regulated by host factors that can be indexed by winter bud formation in *Ribes*, the alternate host. Prevailing cool temperatures are required, but temperature controls induction less than previously suspected. These findings permit greater temporal flexibility when experimenting with the short-lived teliospores. Identification of the specific physiological changes responsible for sporulation could serve as targets for artificial control of the parasite. INT(489).

181. PL-480 research in India identified a common stem rust of Chir pine to be the cause of mortality to plantings of Canary Island pine in the western Himalayas. The alternate host, *Swertia*, is not known within the Canary Islands, but the rust should be considered as a potential threat to the culture of Canary Islan(pine on the continents where *Swertia* occurs. WO(476).

182. New evidence suggests that black locust witches'-broom might be caused by a mycoplasma. Electron microscope studies of ultrastructure showed mycoplasmalike bodies in phloem cells of broomed locust. Such bodies were not found in healthy tissues. This provides new hope that the disease might be susceptible to treatment with systemic antibiotics. NE(542).

183. Techniques for mass-rearing *Dendrosoter protuberans* and improved methods for studying its biology are needed before the effect of this parasite on elm bark beetle populations can be ascertained. Recent studies showed that *D. protuberans*

can oviposit and develop on smaller European elm bark beetle larvae developing in artificial media. This permits continuous rearing and detailed studies under controlled conditions. NE(348).

184. Two diets have been developed that permit completion of larval development of the smaller European elm bark beetle in the laboratory. Though unsuitable for oviposition, the two diets represent an advance in developing artificial means for rearing elm bark beetles. NE(342).

185. A mite, *Pyemotes scolyti*, is an important predator of the smaller European elm bark beetle. Cross-breeding with a mite which preys upon the fir engraver beetle (*Scolytus ventralis*) revealed that the mites are the same species. Thus, the mite strain from either beetle species may have control potential for both insects. SO(362).

186. Cooperative research with Yale University clarified the effect of temperature on growth of several *Ceratocystis* fungus species. A constant temperature of 25° C was optimum for most species and temperature fluctuations generally reduced growth. An understanding of physiological responses to environmental conditions will aid further research on these fungi which include the causal organism of Dutch elm disease. NE(528).

187. A polysaccharide isolated from filtrate of oak wilt fungus cultures induces wilt symptoms in oak cuttings. The compound was chemically described as a mannan. This phytotoxin explains part of the oak wilt disease syndrome. NE(515).

188. Control of the insect vectors of oak wilt disease requires knowledge of the biology and behavior of the oak bark beetles, *Pseudopityophthorus minutissimus* and *P. pruinosis* . The overwintering stages and period of adult emergence of these species have been determined. This information will be of value in determining the timing of control actions. NE(371).

189. An outbreak of sapstreak disease of sugar maple was discovered in Wisconsin. The disease was traced to root and trunk scars created by previous logging activity. To prevent further disease intensification, harvesting should cease for a few years in areas contiguous to diseased trees. NC(499,500).

190. Prevention of insect damage to seed crops requires an intimate knowledge of insect-plant interactions. The interrelationships between insects and seed production have been described in a reference book on seed biology. This publication will be a valuable reference to seed producers. PSW(326).

191. Various investigators have sought to establish and prove that certain glands produce sex pheromones in insects. It has been determined that such a gland is located in the intersegmental membrane between abdominal segments 8 and 9 in female *Dioryctria abietella* adults. The function of the gland has been confirmed by bioassays with male moths. SE(334).

192. *Eucosma cocana* has been found to cause destruction of second-year cones of loblolly and shortleaf pines in Georgia. New information has been obtained on the larval biology, parasites, and predators of this pest. SE(333).

193. The pathogenic nature of several nematodes to southern pine seedlings has been clarified. The lance nematode proved damaging to sand pine, while lance, lesion, stunt, and pine cystoid nematodes had deleterious effects on longleaf pine. SE(536,537).

194. The influence of nematodes on forest trees was summarized as part of a new book "Economic Nematology." Some nematodes such as those parasitizing destructive forest insects are beneficial, but many others directly harm tree roots and increase their susceptibility to pathogenic fungi. Although estimates of impact are lacking, nematodes are known to cause serious problems in forest nurseries and are suspected to cause widescale growth loss in plantations and natural woodlands. SE (535).

195. Favorable responses of Douglas-fir to nitrate fertilization were attributed to mycorrhizal fungi. Nitrate reductase provided by symbiotic fungi allows reduction of nitrate to forms that can be assimilated and used in tree growth. PNW(512).

196. Several mycorrhizal and other root-associated fungi were identified as new species of *Endogone*, *Elaphomyces*, and *Phaeocollybia*. Proper recognition of mycorrhizal fungi is a necessary prerequisite for developing an understanding of natural biological systems for root disease control. PNW(552,558,559).

197. Root numbers and their distribution and condition differ significantly for loblolly pines growing on flat sites and on low, pimple mounds in central Louisiana. A 2-year study revealed that trees growing on the wet, flat sites were deficient in fine and mycorrhizal roots compared with their neighbors on the mounds. Response of flat site trees to intermittent, severe moisture stress probably contributed to premature tree decline and susceptibility to bark beetle attack. SO(352).

198. Sand pine was shown to be susceptible to the root disease fungus, *Phytophthora cinnamomi*. High mortality occurred in groups of experimental seedlings within 2 weeks following planting in infested soil. It is suspected that *P. cinnamomi* alone, or in combination with the root fungus *Clitocybe tabescens*, limits successful establishment of sand pine in poorly drained sites outside its natural range. SE(533).

199. The black root rot pathogen which is a common pest in forest nurseries in warm areas might also be a potential hazard in cooler situations. Cooperative research with Yale University indicates that the pathogen may vary in temperature response and that natural selection processes could allow adaptation to lower temperatures. NE(529).

200. *Armillaria* root rot of sugar maple is an especially serious problem to trees under stress. Stress such as that following premature defoliation by gypsy moth was found to increase susceptibility of sugar maples to severe root rot damage. Artificial defoliation reduced starch reserves and increased levels of simple sugars and amino acids. These and other chemical changes in roots favor the development of the root rot fungus. Artificial regulation of biochemical changes resulting from stress could reduce harmful effects. NE(562-564).

201. A highly virulent root disease of forest plantations was identified in Brazil. The causal fungus, described as a new species, *Cylindrocladium clavatum*, is lethal to both native *Araucaria* and exotic pines. Although currently known in only a few locations, this disease poses a serious potential threat to reforestation of cleared tropical hardwood sites. SE(494).

202. A root rot fungus, *Phellinus torulosus*, known previously to occur only in Europe and Asia, has been found in two locations in southeastern Arizona. It occurs at elevations of above 8,000 feet, mostly on southwestern white pine in which it causes a white pocket root rot. The fungus probably is not a new introduction but rather an overlooked problem common in high elevation southwestern white pine stands. FPL(485).

203. The complex chemical-microbiological system of wood decay in living trees in being elucidated. Early responses of stems to wounding include a decrease in enzyme activity, reduction of starch, occlusion of vascular elements, and an ac-

companying succession of microorganisms. Some bacteria that occur in decaying wood have a nitrogen-fixing ability that probably aids in the nutrition and activity of wood-destroying fungi. Complete understanding of the decay processes can lead to improved means of preventing defect and to management of forest residues. NE(531,543,545,547), PNW(541).

204. White-rotting wood decay fungi produce unique forms of peroxidase which might be of commercial value. Four isozymes were detected from a *Poria* fungus, and three were distinct from those commercially extracted from horseradish. Because of their unique composition, fungus peroxidase could find use in industrial synthesis of lignin-related compounds. SE(504).

205. Extensive losses of usable wood in oaks due to decay occurs in eastern United States. Decay generally increases with age and is especially prominent in the black (red) oak group. Fire scars, large dead branches, and various wounds are common entrance points for the numerous fungi involved in the decay process. The incidence of decay can be reduced by better fire control and by silvicultural procedures such as controlling density to reduce branch size and damage by wind and ice, by reducing harvest rotation ages, and by minimizing injuries during logging or thinning operations. NE(480,491), SO(513).

206. Decay causes little economic loss to hickory in the Central States although individual trees do suffer severely. Fire scars and other wounds are entrance points for several fungi. Estimates based on destructive sampling indicate that overall losses account for only 1.6 percent of the gross volume. Consequently, decay will not limit utilization of the hickory resource. NE(479).

207. Ring and ray shakes greatly reduce the value of wood for many products. Dissections of numerous trees showed conclusively that shakes were associated with wounds. These findings clarify the basic understanding of shake formation and promise for eventual reduction in losses caused by shakes. NE(546).

208. Female carpenterworm moths produce a potent sex pheromone that lures the male for mating. Bioassays indicate an acetate of an unsaturated alcohol with 14-16 carbon atoms is the active component. SO(376).

209. Field studies in Mississippi hardwood forests revealed that emergence of carpenterworm moths begins in April, peaks in late May-early June, and ends in late June-early July. Timing of female and male emergence differs as do daily emergence patterns. These findings will help pest managers to predict seasonal and daily occurrence and patterns of attack in the South. SO(377).

210. Tree age influences trunk borer infestations in cottonwood plantations. Studies of plantations representing 8 age classes and two sites revealed that the cottonwood clearwing borer is a problem during the first 3 years following establishment. The poplar borer is a problem thereafter. Site quality has no descernible effect on the level of borer infestation. SO(315).

211. The living beech borer, *Goes pulverulentus*, severely injures individual hardwoods in the United States and southern Canada. Its biology and habits in red oak and American beech were studied in Arkansas and Mississippi. Though not a major pest, this borer does cause significant damage, particularly in trees that are small and occupy subordinate positions in the stand. SO(375).

212. PL-480-sponsored studies in India indicate that gummosis of eucalyptus is caused by wounding. Gummosis is a natural reaction in trunks wounded by animal chewing, attack by insects or fungi, and various physical disturbances such as result from fire, frost, and wind. It is essential, especially in eucalyptus with smooth bark, to reduce trunk wounding in order to produce high-quality wood. WO(475).

213. Very few natural enemies are known to affect wood borers. In a 3-year study of woodpecker predation in central Ohio, it was determined that first-year red oak borer larval populations in the inner bark were reduced by as much as 33 percent. Thus, woodpeckers may have a significant effect in regulating borer populations and their damage. NE(345).

214. The cerambycids *Saperda inornata* and *Oberea schaumii* infest and damage the stems of small trembling aspens and the root suckers and twigs of large trees. Their distribution within the crowns of larger trees, the importance of their galleries as infection courts for *Hypoxylon* cankers, and the relationship of infestation to site quality has been determined for Michigan aspen stands. This information contributes to a better understanding of these insects in the Lake States. SE(365-368).

215. The black twig borer, *Xylosandrus compactus*, has spread to many host plants in the Hawaiian Islands since its first discovery in Hawaii in 1961. Beetle infestations have caused heavy tree damage, and recent attacks have been associated with the death of apparently health forest trees. To date, mortality has occurred in five tree species growing in plantations on the island of Oahu. PSW(363).

216. Hawaiian forests of native ohia and koa are suffering from an acute decline disease. Recent aerial and ground surveys indicate a steadily deteriorating condition, especially on the slopes of Mauna Kea. Studies were initiated to determine the cause and ecological consequences of this severe disorder. PSW(509).

217. Physiobiochemical research on cellular protoplasm is providing new explanations of the mechanisms by which plants resist extremes of heat, cold, and drought. Changes in amounts and kinds of protoplasmic chemicals, such as amino acids and sugars, affect protoplasmic viscosity, membrane plasticicity and permeability, and water-ice relations. These are associated with seasonal and species variations in resistance to stress. An adequate understanding of stress resistance could lead to amelioration through plant breeding and chemical regulation. NE(518,519).

218. A widespread decline of ponderosa pine in the Denver area, previously thought to be due to air pollution, was found instead to be caused by excessive soil salinity. In cooperative studies with Colorado State University, typical tipburn symptoms were induced by experimental applications of salts. Total salts and chloride levels in soils with injured pines exceeded levels around healthy pines. Improving soil drainage shows promise for ameliorating this problem in ornamental trees. RM(555).

219. An information retrieval system (SOLAR), designed cooperatively with Washington State University, to facilitate searches of plant pathological literature is operational. The present data base of 10,000 abstracts was prepared primarily for white pine blister rust. It is expected to grow to 40,000 abstracts by 1975, with rather complete coverage of the physiology and biochemistry of plant diseases. A compatible remote terminal and telephone can provide worldwide access to SOLAR, thereby improving the efficiency of pathological research. INT(492,530).

220. The status of aspen in the Lake States has been reviewed. Aspen is subject to several debilitating diseases. Some stands are too decadent for economical harvest and will

eventually deteriorate and be replaced by other species. Unless the rate of harvest is increased or the pathological rotation age is raised by reducing the impact of disease, large acreages of aspen will be lost. NC(473).

221. Aspen on the Gros Ventre elk winter range in Wyoming are dying because of a combination of "barking" by big game, injurious insects, and diseases. The outlook for aspen regeneration is especially critical because of the heavy impact of browsing, pests, and competition on aspen sprouts. If the trend continues, about a two-thirds reduction in overstory aspen stems is predicted by the year 2000. The Forest Service is now studying the possibility of using fire to regenerate the deteriorating aspen type. INT(506).

222. A monograph on forest diseases in India climaxes 5 years of intensive forest disease research carried out under the PL-480 program. The treatment contains a comprehensive guide to the numerous parasitic organisms that attack over 100 species of forest trees. Silvicultural measures to reduce damage are suggested for each of the more serious diseases. This monograph is a major step toward the growing of healthy, highly productive stands of forest trees in India. WO(477).

223. Scientists sponsored by PL-480 carried out quantitative forest disease surveys in West Pakistan. Prominent diseases included diebacks of mulberry and *Dalbergia*, leaf rust of *Acacia*, blights of juniper and deodar cedar, and heart rots of various conifers and hardwoods. This marks a major step in forest pathological research in West Pakistan where healthy forests are in great demand because of an inadequate supply of high-quality wood. WO(527,569).

224. A recent analysis suggests that the adverse impact of diseases and insects to wildland shrubs is comparable to that experienced with forest trees. Numerous diseases and pests are especially troublesome to important browse shrubs on big game winter ranges in western United States. Increased emphasis on pathology and entomology is necessary to insure the success of programs to improve wildlife habitat and produce genetically superior shrubs. INT(341).

225. Damaging forest diseases and insects may increase in significance as intensively cultured forests are created to meet the growing demand for wood for an exploding world population. The trend toward highly mechanized operations results in a biotic simplification that could easily lead to catastrophic forest damage from diseases and insects. To prevent such losses, broadly based research programs need to accompany the development of intensive forest culture. WO(544).

226. A key to adult bark beetles commonly associated with white spruce damage in interior Alaska is available for scientists and managers. PNW (680,681).

227. The mahogany shoot borer (*Hypsipyla grandella*), a serious pest of the Meliaceae (mahoganies and related species) in the Western Hemisphere, has made it almost impossible to raise plantations of these valuable tropical hardwoods. Results of field and laboratory studies in Puerto Rico indicate that the various Meliaceae species produce chemical substances poisonous to *Hypsipyla* species, but the specific shoot borer which has evolved with each particular tree species has adapted to these substances and uses them as an attractant. This information will be helpful in devising methods for controlling the borer or reducing its damage. ITF(714).

3. Pest control techniques

228. Biological control of dwarf mistletoes appears possible but costs are uncertain. An analysis of current knowledge suggests that the best approaches lie in the introduction of foreign insects that attack only dwarf mistletoes and in the encouragement of certain fungal diseases, such as resin disease, which destroy the mistletoe root systems along with foliar shoots. RM(575).

229. Dwarf mistletoe causes extensive damage to hemlock in west coastal North America. In just Washington and Oregon, annual timber losses are estimated at over 40 million cubic feet. In addition, associated witches'-brooms and decay increase danger from falling trees in recreation areas. Effective control is possible by complete harvest of infested stands, or in recreation sites by pruning of brooms and removal of hazardous trees. PNW(596).

230. Silvicultural control practices have reduced the level of dwarf mistletoe and protected the ponderosa pine recreation forest on the South Rim of Grand Canyon. Plots studied for 20 years document that periodic sanitation is an effective control for dwarf mistletoe in high-use recreational sites. Sanitation has forestalled mortality and associated vegetative changes along the South Rim, thereby sustaining scenic and esthetic values provided by ponderosa pine. RM(582).

231. Combinations of three synthetic attractants, host monoterpenes, and infested bark were tested in sticky traps and on live trees to determine their effectiveness in inducing attack or disrupting aggregation of Douglas-fir beetles. Frontalin-treated trees were attacked but most of these and adjacent attacked trees survived. Methylcyclohexenone (MCH) disrupted or blocked attraction to frontalin-treated trees or traps and the associated "spillover" into adjacent trees. Transverbenol had no conclusive effect on beetle aggregation behavior. It is concluded that MCH offers a viable alternative for regulating beetle populations and protecting susceptible trees. INT(403,415).

232. Reduced concentrations of pesticides or the use of an oil carrier alone promise to reduce costs and environmental contamination in western bark beetle control programs. Field tests in Wyoming using 0.25 pound of ethylene dibromide in 5 gallons of fuel oil or fuel oil alone killed at least 90 percent of the spruce beetle larvae and pupae in downed trees. Large-scale tests are now needed to determine the feasibility of altering current operational control recommendations for this and other bark beetle species. RM(416).

233. Better guidelines are needed for determining treatment height on mountain pine beetle-infested ponderosa pines in Colorado. Systematic analyses of brood development in relation to height of attack, presence of pitch tubes, and blue stain have shown that trees need to be treated only to within 5 feet of the highest beetle pitch tubes. This will prevent overtreatment, reduce costs, and reduce effects of the pesticides on natural control organisms occupying the upper trunk. RM(408).

234. Small-diameter lodgepole pines were baited with the attractant trans-verbenol and the host terpene, alpha pinene, to determine if mountain pine beetles could be attracted to such trees and away from nearby, larger trees. Although baited trees less than 8.5 inches in diameter were generally attacked, a greater number of nearby, unbaited, larger-diameter trees were also killed. thus, further refinements are needed before we can effectively regulate mountain pine beetle populations with attractants. INT(410).

235. The fungicide benomyl shows great promise for control of Dutch elm disease and oak wilt. A water-soluble derivative, MBC-HCl, was synthesized and is being extensivly tested for efficacy. This form of benomyl is soluble even in high concentrations, it has a low viscosity which favors uptake and vascular distribution, it is chemically stable, and it retains fungitox-

icity for longer periods with no phytotoxicity at working concentrations. Preliminary results from field trials, using pressure injection, suggest that it is a promising candidate for disease prevention and therapy. NE(574,585).

236. Tests were conducted to determine if the smaller European elm bark beetle—the principal insect vector--would be affected adversely by feeding on twigs containing benomyl. The disease was suppressed, but beetle feeding, longevity, and reproduction were unaffected by the fungicide. NE(404).

237. Bioassays of elm bark beetle feeding must be correlated with assays of insecticide residues by gas liquid chromatography to provide a basis for predicting beetle feeding on treated, living elms. Twig crotches collected from trees sprayed by helicopter and by mist blower revealed low and variable correlations for helicopter-sprayed trees. Correlations were high and uniform for mistblower-sprayed trees. Thus, mistblower applications provide more reliable protection from elm bark beetle attacks than do helicopter applications. NE(393).

238. Forest tree seedlings can now be protected during storage. Benomyl fungicide, applied either as a prelifting field spray in the nursery or as a seedling dip, reduced losses of 50 percent in giant Sequoia nursery stock to a negligible amount. With benomyl treatment, gray mold should no longer be a detriment to successful cold storage. PSW(600).

239. The proceedings of an international meeting on resistance of forest trees to rust diseases held in 1969 in Moscow, Idaho, was published. More than 140 specialists from all over the world discussed white pine blister rust and new research developments in hard pine and cereal rusts. Considerable progress toward production of rust-resistant white pines has been made, along with new ways of increasing the efficiency of tree breeding programs. INT(571).

240. Resistance to western white pine to blister rust is a composite of independent mechanisms inherited in a simple Mendelian manner. These mechanisms include exclusion or abortion of needle infections and walling-off of stem cankers. It now appears feasible to develop a balanced system of resistance in western white pines. Such resistance would maintain a gene pool within forest populations that could guard against changes in virulence by the blister rust pathogen. INT(576,577,584).

241. Methods for large-scale inoculation of western white pine progeny used in developing resistance to white pine blister rust were evaluated. Inoculations are made in covered nursery beds with inoculum from *Ribes* leaves. The technique results in heavy but spotty infection. Lack of uniformity continues to be a costly problem that is overcome by computerized statistical analysis. INT(570,583).

242. A method was developed for uniform inoculation of large numbers of pines or oaks with fusiform rust. Water suspensions of basidiospores or aeciospores are applied in an aerosol spray on greenhouse-grown slash and loblolly pine or oak seedlings. This method was increase efficiency and accuracy of an operational program for selecting rust-resistant southern pines. SE(573,589).

243. Microscopic examination revealed an apparent anatomical mechanism in southern pines for resistance to the fusiform rust fungus. The mechanism involves the death of cells in and around the infection point in young stem tissues. Reaction areas apparently wall-off the pathogen and the new tissues develop disease-free. Knowledge of how rust resistance operates can lead to more efficient programs of genetic improvement of valuable southern pines. SO(578).

244. Intensive site preparation, including burning and plowing before planting, significantly increased the incidence of fusiform rust in slash pine plantations in southeastern Georgia. When examined 7 years later, plantations established on treated sites had more fusiform rust than did those in which planting was done without prior site preparation. However, site preparation also increased overall tree growth. Caution in site preparation should be exercised in high rust hazard sites. SE(590).

245. Ferbam, carboxin, and oxycarboxin provided better control for fusiform rust of loblolly pine in nursery seedbeds than did benomyl or chloroneb. Apparently none of the test fungicides were systemic even with the addition of DMSO. Satisfactory control of fusiform rust in southern forest nurseries still requires application of foliar sprays before, during, and after periods of wet weather. SE(594).

246. In field trials, sex pheromone perception by male European pine shoot moths was inhibited by the use of synthetic acetates. Cis-7, cis-8, and cis-9 dodecenyl acetates were particularly effective. One or more of these materials may have potential as a behavioral control agent for manipulating populations of this insect. PNW(396).

247. Case histories of five Douglas-fir tussock moth outbreaks in California and Oregon revealed that all infestations followed a 3-year cycle. Minimal defoliation occurred during the first year, severe foliage loss the second year, and minor defoliation and ultimate collapse of the populations by the end of the third year. Although other factors were involved, a virus disease appeared to be the principal natural cause of insect mortality during the collapse. To prevent serious damage, outbreaks must be detected early and controls applied selectively in high moth population centers. PNW(420).

248. Use of nucleopolyhedrosis viruses in controlling forest and agricultural insects requires that microbial preparations be essentially free of hazardous contaminants. Oak Ridge National Laboratory and Forest Service scientists have developed a procedure for purifying specific virus formulations for the Douglas-fir tussock moth, European pine sawfly, and the gypsy moth in quantities sufficient for operational use. NE(395).

249. A hymenopterous parasite, *Agathis pumila*, was released at five locations near St. Maries, Idaho, in 1960, to control the larch casebearer. Although the parasite became established in three of the areas and caused up to 68 percent mortality, its inability to spread has limited its usefulness. Consequently, the casebearer has spread through more than 90 percent of the range of western larch. Researchers have therefore altered their approach to include introductions of multiple native and/or introduced parasite species and other means of control. INT(398).

250. Alternative, environmentally acceptable toxicants are needed for control of many forest defoliators. Several insecticides have shown promise in controlling *Malacosoma disstria*, *Hemerocampa vetusta*, *Symmerista canicosta*, and *Acleris gloverana*. Field tests are needed to establish minimum effective dosage rates and optimal timing of applications. PSW(407,409,413,414).

251. Juvenile hormone analogs effect the fecundity, development, and survival of insects. Several chemicals were tested on 6th instar larvae of the western spruce budworm. Low dosages tended to decrease egg production and hatch. Higher dosages increased the degree of deformity, disrupted growth and development, and increased mortality. More potent compounds caused additional moulting. PSW(412).

252. Preparations of Thuricide HPC were applied with a mistblower at rates of $16x10^9$ and $8x10^9$ International Units per acre, or 1 gallon and 0.5 gallon per acre, respectively, for gypsy moth control. These treatments, applied at 10-day intervals, provided partial protection of foliage and significant larval mortality. Unsprayed areas were nearly completely defoliated. Further study is needed to determine optimal dosage rate, number of applications, and timing of treatments. NE(421).

253. The construction details, costs, and calibration data for a microbial insecticide spray tower have been described. This device permits precise dosage and formulation evaluations at low cost prior to actual aerial field testing. NE(406).

254. The gypsy moth is currently spreading southwestward into new environments where its natural enemies are sparse or lacking. Rearing techniques and biologies are being determined for five promising parasites by the USDA for possible mass release and colonization in these new areas. NE(411).

255. Mistblower applications of malathion are registered for use in suppressing populations of the red-headed pine sawfly. Further tests with aerial and mistblower applications of this pesticide applied at the rate of 10 ounces per acre and 2 gallons per acre, respectively, provided 84-100 percent control. These data would support the registration of aerial application of malathion for control of this sawfly. NE(401).

256. The major volatile trail-marking pheromone of the Texas leaf-cutting ant, *Atta texana*, has been isolated, identified as methyl-4-methylpyrrole-2-carboxylate, and synthesized. The synthesized pheromone and several of its analogs elicited strong trail-following responses from worker ants in the laboratory and the field. The synthetic pheromone offers potential in trapping ants or in leading them away from food sources. SO(417,419).

257. A defoliating weevil, *Magdalis gentilis*, is attracted to thinned stands of lodgepole where it may destroy up to 100 percent of the foliage on crop trees. Though damage to forests has been limited to date, activity of this insect can be minimized by delaying thinning until late July. INT(400).

258. A pictorial guide has been prepared for the insects and diseases affecting southern seed orchards. Hosts, distribution, damage symptoms, biology, and cultural control treatments are presented for each species or species group. This publication will be used by orchard managers and others interested in seed production. SE(405).

259. Soil applications of 1 lb. of the systemic insecticide phorate per tree in Alabama, Georgia, and Florida seed orchards reduced infestation or damage by *Dioryctria* spp., *Laspeyresia* spp., thrips, and seed bugs by more than 80 percent. The treatment also increased the percentage of sound seed harvested. Lower dosages were less effective, suggesting that the 1 lb. rate was the most efficacious. SE(397).

260. Twenty-two different species of predators were introduced from Europe, Asia, and Australia during the period 1959 through 1966 to control the balsam woolly aphid on Fraser fir in the Mt. Mitchell area of North Carolina. From observations in 1968, *Larcobius erichsonii* is the only species established. Three species of native, predatory mites were found to be abundant and, possibly, more important as natural enemies. However, there is no tangible evidence as yet that the native or foreign predators have significantly altered the course of this outbreak. SE(399).

261. Damping-off, nematodes, root rots, diebacks, and several leaf diseases periodically reduce nursery production. Sanitation, soil amelioration and fumigation, and protective foliar fungicides are recommended to meet production demands for disease-free planting stock. SE, SO(595).

262. Basic copper sulfate fungicide can partially control needle blight of Arizona cypress in southeastern nurseries. Tests were made of 30 materials in search of a replacement for restricted mercury compounds. Basic copper sulfate provides reasonable control except during severe disease epidemics. SE(593).

263. Control of the *Poria* root rot fungus by biological means is being evaluated in the Pacific Northwest. Nitrogen and organic compounds of red alder influence soil chemistry and prevailing microflora in ways that inhibit *Poria*. On *Poria*-diseased sites, short, merchantable rotations of pure alder or mixtures of alder with *Poria*-susceptible conifers are promising for control of the disease. PNW(601-603).

264. Reasons were found for reduced colonization of pine stumps by the annosus root rot fungus during late spring and summer in the Southeast. The explanation lies in a complex combination of reduced inoculum, high temperatures, and competing organisms such as *Peniophora gigantea*. This confirms an earlier suggestion that plantations in southeastern United States south of approximately 34° N. latitude can be thinned in late spring and summer with little danger of increasing annosus root rot in residual trees. SE(592).

265. Progeny of loblolly pines used in a southeastern cooperative tree improvement program were shown to be susceptible to annosus root rot. Mortality was uniformly high in inoculation tests among the more than 10,000 seedlings from controlled crosses of 151 "superior" parents. This nullifies an earlier first-year report of suspected resistance. If resistance is needed to combat this disease, it probably will be necessary to select specifically for this trait. SE(579).

266. Research has verified the importance of mycorrhizae as agents for biological control of rootlet pathogens. The mechanisms whereby mycorrhizae induce resistance include the creation of physical barriers to invasion by pathogens, production of antibiotics that inhibit pathogens, and chemical stimulation of favorable rhizosphere microorganisms that successfully compete with pathogens. Research now underway will assess the potential of several mycorrhizal fungi for inoculation into forest nursery soils. SE(586,587).

267. Mycorrhizae protect slash pine from dehabilitating attack by *Phytophthora* root rot. Controlled experiments indicated that slash pine seedlings bearing ectomycorrhizae formed by *Pisolithus tinctorius* or *Cenococcum graniforme* can grow well in soil infested with *P. cinnamomi*, whereas nonmycorrhizal seedlings in similar infested soil become chlorotic and stunted. This is important new evidence useful in developing strategies for protecting the rootlets of forest trees from pathogens. SE(588).

268. Standard treatments with 1 percent aldrin solutions dispensed from planting machines have provided erratic control of white grubs in the Lake States. Reevaluation of this method, as well as applications of 1 percent aldrin solutions with a backpack sprayer and granules dispersed from a planting machine revealed that all three methods properly carried out would provide adequate protection for at least two seasons. NC(402).

269. The composition of resin may have a prime role in determining succession of fungi invading pine sapwood. Variation in the growth of several sapwood fungi occurred in laboratory trials containing pine terpene additives. Early colonizers were generally more tolerant than their successors and variation was expressed within species isolates as well. Screening of genetic variants for possible use in hastening

degradation of logging residues might be feasible by this pure-culture technique. SO(572).

270. Placing paraformaldehyde pills in sugar maple tap holes to increase sap flow can lead to heart rot. Experiments in the Northeast determined that the frequency and extent of decay is significantly greater with this practice than with untreated tap holes. Losses due to decay more than offset the gains in maple sap yields. It is advised that paraformaldehyde not be used. NE(598).

271. Preliminary results suggest that dressings applied to tree wounds are ineffective in preventing invasion of microorganisms. Fresh axe wounds on sugar maple and American elm were either coated with asphalt dressing, orange shellac, polyurethane varnish, or left untreated. No differences in invasion of microorganisms and wood discoloration were noted one year later. Improved wound dressings appear necessary to prevent defect in valuable shade and ornamental trees. NE(597,599).

272. Snow movement causes severe stem defects in conifers on steep, mountainous terrain. Studies in Idaho defined six classes of deformities of which "butt sweep" is the most frequent, whereas, "dog leg" and "stem failure" are the most injurious. Deformed stems reduce mechantability and together with other economic and environmental costs make commercial forest operations on steep slopes difficult. Where harvesting on steep slopes is feasible, partial cutting is recommended to minimize stem deformatives since residual trees impede snow movement. INT(581).

273. Salvage harvesting in northern Idaho of pole blight-affected western white pines had no effect on disease development in residual white pines, but it did influence associated conifers. Pole blight continued to develop in remaining white pines. Western redcedar responded well to release, but the Douglas-fir component suffered heavy mortality from *Poria* root rot. These findings will aid in predicting future consequences of silvicultural practices on similar sites in the Inland Empire. INT(580).

4. Pest management strategies

274. A new cost-optimizing procedure allows managers to formulate goals and safety standards for control of hazardous trees in recreational sites. A selected goal and a uniform hazard-rating procedure can be established and a balance between hazard, control costs, and attractiveness of the forest site attained. PSW(604).

275. Common diseases of shade forest trees in urban areas of the South can be reduced. Selection of disease-resistant trees, special cultural techniques, and fungicidal controls are integrated into recommendations to promote a healthier, more satisfying forestlike environment in southern cities and suburbs. SE(605).

276. The effects of mountain pine beetle, the need for control, and the effectiveness of certain suppression methods are critical questions facing managers of lodgepole pine stands. An evaluation of 10 treated and untreated areas in Wyoming and Idaho showed that chemical control treatments slowed the rate of tree mortality but did not affect the beetle outbreak period or the final number of trees surviving. Generally, harvesting of large-diameter, thick phloem pines at mid- and low-elevations in stands composed of 64 percent or more lodgepole pine was suggested for reducing the potential for severe outbreaks. INT(422).

277. *Phytophthora* root rot continues to threaten survival of the unique and highly valuable Port-Orford-cedar. The disease spreads through movement of contaminated soil and water. Exclusion of people, equipment, and animals from drainages that still harbor healthy stands is the only means of prevention. Practical ways to restrict movement of contaminated soil are advised for salvage logging of infected stands. Losses can be minimized, but there appears no hope of raising another merchantable rotation of Port-Orford-cedar under existing conditions of disease and land use. PNW(606).

G. Accelerating Forest Resource Surveys

1. Forest inventory

In a continuing effort to provide current information for future timber supply analysis, field work was conducted in Louisiana, Alabama, Vermont, New Hampshire, Wyoming, Utah, South Dakota, New Mexico, Missouri, Tennessee, Georgia, Oregon, and California. Reports appraising the forest resource condition were issued for five States or parts of States.

278. Wisconsin's growing stock volume increased over 30 percent between 1956 and 1968 to a level of 11.0 billion cubic feet, and sawtimber volume increased from 13.8 to 21.8 billion board feet. These gains were made despite a small loss of commercial forest area—from 14.9 million acres in 1956 to 14.5 million acres in 1968. The aspen-birch type (4.2 million acres) and the maple-beech-birch type (3.5 million acres) led other types in area. Growing stock growth in Wisconsin in 1967 exceeded removals by over 100 percent, 500 million cubic feet to 233 million cubic feet. NC(1053).

279. Preliminary statistics on Delaware's forest resources indicate that forest area has remained about the same but inventory volume has increased. Growth exceeded removals for hardwoods but growth only equalled half of removals for softwoods. Thus, the softwood proportion of inventory volume has dropped sharply. NE(1039).

280. Tennessee's commercial forest area declined 5 percent between 1961 and 1971, with a diversion to pasture dominating the loss. Softwood timber volume increased 26 percent and hardwoods gained 11 percent in spite of this area loss because of the net growth excess over removals. Most of the volume increase accrued to trees less than 18 inches in diameter. The same pattern of constant or declining commercial forest area but increasing timber inventory was also observed in the 1972 reinventory of Alabama. SO(1049,1050), SE(1043-1045).

281. The first complete inventory of timber resources in Mendocino and Sonoma Counties, California, indicates that there is 19.1 billion board feet of sawtimber on 1.56 million acres of unreserved forest area. Almost half the area is stocked with low-quality hardwoods and scattered conifers resulting from logging, wildfires, or unsuccessful attempts to convert timberland to grazing land. In recent years forest industries of the counties have been utilizing predominantly old growth from their own lands. In the next 10 to 15 years they will have to rely increasingly upon young growth redwood sawtimber, much of it from farm and miscellaneous private lands. PNW(1052).

282. Because of the rapidly changing nature of the Southern forest resource and industrial structure, more detailed and up-to-date information than is available in published reports is often requested. In response to this, both the Southern and Southeastern survey units have developed custom data retrieval capacities. For example, data on more than 100 specified areas were compiled in the Southeast in FY 1973. SO(1042).

283. A broader scope of forest inventory data is needed to enable land managers to evaluate the feasibility of meeting in-

creasing demands for nontimber uses of the forest. A transitional sampling design was developed that permits comparisons between stands for size, accessibility, productivity, and restrictions imposed by competing uses. Trial use of this design in Montana and Idaho in 1970 was the first step in establishing complete timberland records. INT(717).

2. Timber utilization and production

284. The trend of similar sawmills dropping from production and larger or medium sized mills replacing them has continued into the mid and late 1960's in four Eastern States. Pennsylvania, Kentucky, Missouri, and Alabama all experienced a 30-45 percent decline since the early 1960's in the number of active sawmills according to recent industry surveys. The same States, however, had constant or increasing saw-log production levels. NC(1065), NE(1067, 1068), SO(1062).

285. Pulpwood production increased roughly 85 percent in Alabama from 1962 to 1971 and the pulp industry recently entered Kentucky with three mills. Pulpwood production dropped slightly in the Lake States and the South from 1970 to 1971, however. NC(1064), SE(1058).

286. Southern pulpwood prices rose slightly in 1971. Softwood pulpwood price was $21.15 per cord in the Southeast, $19.49 per cord in the Midsouth east of the Mississippi, and $18.73 per cord in the Midsouth west of the Mississippi. SE(1072), SO(1063).

287. Midsouth pulpwood hauling distances from the stump to the mill are about what they were in 1960. Forests within 50 airline miles of pulpmills have consistently supplied about half of the wood requirements. SO(1059).

288. The percentage of primary plant residues used for other products varies greatly by State. Roughly 85 percent of the residue was used in Alabama in 1971 but only 40 percent was used in Missouri in 1969. Pennsylvnia (1969) and Kentucky (1969) fell between these States. the dominant residue market is for pulping material. Alabama's excess residue was mostly fine particles which generally have too short fibers for pulping, however. Most of the unused Missouri residue was hardwood and the State only has two pulpmills. A 1968 survey of mill residue flows in Washington, Oregon, and California showed that 88 percent went to domestic plants and 12 percent was exported. Eight-three percent of the domestic residue shipments was used by pulpmills. NC(1065), NE(1067,1068), SO(1062), PNW(1069).

289. The 1971 Oregon log harvest was up 12 percent from its 1970 level but Washington harvest was unchanged. Log harvest levels in Western Oregon have been steady since the early 1950's with the increase in public harvests offsetting the decline in private harvests. Western Washington harvests have been increasing on both public and private lands, reflecting to some extent the increased availability of second-growth stands. Eastern Washington and Eastern Oregon harvests have increased since the early 1950's. Pacfic Northwest log production has been counter cyclical with reference to the business cycle since World War II. Housing starts lagged one year, and National Forest stumpage prices were significantly related to log production. PNW(1074-1077).

290. Estimates of logging residue in western South Dakota, Wyoming, Utah, and Colorado for 1969 ranged from 4.72 percent of the cubic volume to 10.43 percent of total growing stock. Most of the residue is the result of damage from tree felling and is wood from trees from which saw logs are harvested. Knowledge of residue volumes is especially important where there is an active wood chip market since the residue volume is usually too small or low quality for sawn products. An on-site chipping study in Wyoming of logging residues in lodgepole pine stands was found to increase fiber yields by 35 percent. INT(1070,1073).

3. Survey techniques

291. Analysis of two-stage 3P sampling, indicates that this approach may be an efficient means of obtaining intermediate volume estimates between full scale forest surveys. A simulated 3P sample predicted the total cubic volume for five counties in southern Alabama to within 0.7 percent of that predicted by the latest forest survey. The 3P sample only required measurement of 4 percent of the trees measured in the full survey. SO (1093,1094).

292. Evaluation of photo inventory plot location has demonstrated that not only do irregularities of ground elevation cause problems but a plot bias may also exist on flat areas under some conditions. It is essential that the photo plots are uniformly distributed over the earth's surface if a bias is to be avoided. Analysis of inventory data in the 12 Pacific Northwest counties revealed that double sampling for stratification, using a combination of photo plots and field plots, is about twice as efficient as simple field plot sampling for timber volume estimation. PNW(1090,1095).

293. The possible use of high altitude and satellite imagery for land resource classification and measurement has induced considerable research work. High altitude color infrared photographs of a 1:120,000 scale covering eight 41-square-kilometer study areas over a period of 20 months were used to identify 13 forest and nonforest land classes. Over 97 percent of the forest points were interpreted correctly regardless of season. The study also revealed the likelihood of specification error for particular land use classes. Computerized techniques for land use classification on 1:120,000 scale color infrared photographs were also analyzed using an automatic scanning micro-densitometer with a red, a blue, a green, and a clear filter. Results were compared with a map derived from 1:32,000 scale color infrared film and ground checks. PSW(1078,1091).

294. Analysis of medium scale (1:32,000) color infrared photographs successfullly monitored the spread of mountain pine beetle infestations in the Black Hills of South Dakota from May to September of 1972. New research permits the optimization of the choice of photoscale against interpretation error. Airborne multispectral scanner equipment was also used for detection of forest stress in the ponderosa pine ecosystem in the Black Hills. Preliminary results indicate the best channel ratios for stress detection. PSW(1096,1097).

295. Normal color and color infrared photos are becoming increasingly valuable tools for detecting symptoms of forest diseases by remote sensing. Recent studies show these films can distinguish between healthy and disease-stressed vegetation in air pollution injury, Dutch elm disease, oak wilt, dwarf mistletoe, ash die-back, *Fomes annosus*, and other diseases. PSW(1085).

296. The field of geographic information systems is a rapidly evolving one in which there has been no commonly accepted vocabulary. To meet this need, a partial glossary of 225 terms has been assembled and is now available as a guide to standard usage. PSW(718).

H. Economics of Forest Management

1. Methods of financial evaluation and planning

297. An improved timber inventory projection model uses exogenous estimates of (1) supply and demand to determine annual cut, (2) changes in acreage which reflect changes in forest investment, and (3) net growth, including changes in investment in cultural practices. This model has several advantages over many other available methods. First, it estimates market equilibrium quantities each time period rather than postulating their intersection at some point far in the future. Second, it focuses attention on an increasingly important forest management problem—changes in land use. Third, it can be adapted to a variety of inventory and supply problems. NC(1098).

298. A method for projecting value growth of hardwood timber stands uses tree-development (changes in diameter, height, and quality) probabilities based on continuous forest inventory (CFI) data. This is the first step in building a more precise tool for forest land managers desiring to recognize the value production potential in their stands and forests. NE(1100).

299. A zero-one integer programming procedure has been developed for systematic planning of commercial thinning operations in young-growth Douglas-fir. This new approach is demonstrated, tested, and compared to existing rule-of-thumb methods of a case study area in western Oregon. PNW(1101).

300. The allowable cut effect, which may be a factor in investment analyses where harvest volumes are dependent upon the physical rate of growth of a forest, is described and illustrated. Some of the physical and economic ramifications of the concept may be important determinants in developing management plans. PNW(1102).

2. Timber growing economics

301. Projections of aspen inventory and cut in the Lake States indicate that today's apparent surplus of aspen cannot continue indefinitely. The analyses assumed three sets of assumptions and historical trends of cut, growth, and utilization. The projections indicate that Minnesota may be able to support more than the projected cut. Forest industries drawing their wood from traditional sources in Michigan and Wisconsin must plan to procure their wood elsewhere within these States, substitute other species of wood for aspen, or cut less than the projected amount during the next 15 to 25 years. NC(1110-1112).

302. The fourth in a series of determinations of financial maturity for the important timber species in the northeast, establishes the rates at which sugar maple trees increase in value due to diameter and height growth and improvement in quality over a 10-year period. Detailed rates of value increase to be expected over a 10-year period are exhibited. The effects of different conversion costs and changing costs and prices over a 10-year period are analyzed. NE(1115).

303. An analysis of the economic desirability of accelerated roadbuilding for access to old-growth timber on a unit of the Umpqua National Forest in Oregon found four accelerated roadbuilding alternatives inferior to the current rate of construction. Only in the case of substantial, continuing inflation were projected rates of return on investment for accelerated roadbuilding above 6 percent. The inclusion of nontimber benefits and costs, changes in price and cost levels since 1966, and recent improvements in logging technology do not appear to alter results appreciably. PNW(1117).

304. The behavior patterns of small woodlot owners revealed by a study in Pennsylvania suggest that we should provide information about the management and marketing of forest products to more owners of large wooded tracts (50 acres or more). Owners who have sold forest products and practiced some form of forest management on their woodlands in the recent past should prove most receptive to sales promotion. NE(1109).

305. Life insurance companies have accepted forest land and timber as loan security since 1950; six firms now have timber loans in 25 States, chiefly in the South and Pacific Northwest. Lending policies vary, but all firms require that the woodland be managed soundly. Two-thirds of the borrowers are individuals. SO(1122).

3. Multiple-use economics

306. A comprehensive model of the U.S. range resource evaluated potential programs of range land development. The system uses linear programming to analyze alternative plans for the management and protection of range land in 36 ecosystems. It shows how each proposed plan affects the output and nine physical products including annual unit months and water yield and 13 environmental values such as soil stability and rare and endangered species. WO(1126).

307. A typical dense chaparral area on the Tonto National Forest was analyzed to explore the economics of chaparral management. Costs and benefits were developed for 12 alternative management plans, ranging from complete conversion of chaparral to grass to leaving the chaparral untouched. Labor, equipment, materials, and transportation costs were identified for each alternative over a 50-year period. Products considered for each alternative include water, sediment, range, recreation, wildlife habitat, esthetics, and fire protection. Primary emphasis was on the development of values for the products affected. This study showed that the method can give adequate information to the land manager for management planning. RM(1130).

308. Economic returns from planting forage in a shelterwood management regime in ponderosa pine-grass areas of eastern Washington are summarized for various assumed planting costs, forage values, and growth data. The approach allows a forest manager to rank forage-planting alternatives according to their rates of return, when better forage yield for domestic grazing is the primary objective or when noneconomic benefits appear equally desirable. PNW(1132).

4. Impacts on forest industry and regional economies

309. An analysis of the economics of producing both timber and pulp chips showed that if pulp and paper firms installed chipping headrigs at pulpwood yards, they could earn greater profits, timber growers could receive higher prices, and society could benefit from improved utilization of the timber resource. SO(1139).

310. Competition in the timber harvesting industry and legal obstacles have thwarted recent collective bargaining efforts by pulpwood producers' associations. A study indicated that cooperatives appear to offer the best prospects for producers to legally bargain collectively with wood buyers. SO(1135).

311. A national forest economy input-output model describes the flows of values in this major economic sector dealing only with industries that process logs and bolts or utilize the wood products immediately after this stage. The model was successfully implemented with data from the Bureau of the Census and other standard sources. SO(1138).

I. Improving Engineering Systems

312. One of the most pressing problems facing forest managers is finding suitable means for utilizing forest residues. The engineering component of this gross problem includes the development and testing of creative handling and on-site processing of these residues. Preliminary trials of on-site chipping of lodgepole pine residues suggests a potential for adding as much as 35 percent greater total wood yield from clearcut areas where whole unmerchantable trees or residue portions of merchantable trees are yarded and chipped *in situ*. INT(1206).

313. Logging residues, chipped in their entirety in the field, contain whole-tree material, including bark. An essential engineering component for residue utilization in paper-making or for most reconstituted wood products necessitates a practical, economic bark-chip separation and segregation technology. Such a technology is now ready for pilot testing. NC(1194-1196, 1202-1205, 1217, 1218).

314. A basic requirement for designing suitably engineered logging systems is a precise means of determining log or tree weights and centers of gravity. Recently completed studies report on estimated log or tree weights for typical northwestern species and for weights and centers of gravity for typical Lake States species. PNW(1212), NC(1216).

315. Proper design and construction of logging road systems is an important engineering element for successful logging, especially where critical environmental stewardship issues are at stake. One critical analysis of a road in sensitive Idaho batholith soils indicates that with proper attention to design and construction, suitable and environmentally secure logging roads can be built. 0For Pacific Northwest conditions, recent studies show how yarding distances for aerial logging systems can be predicted from map data and how road and landing criteria can be subsequently engineered into the logging plan. PNW(1200, 1213), INT(1201).

316. Advanced logging concepts are moving from the laboratory to operational situations. Recent trials of balloon logging in Idaho provide the initial data for substantial modification and improvement to suit conditions heretofore untried. 0Running skyline systems developed through research over the last 5 years are now operational and are proving themselves under a variety of field conditions. These initial systems suggest a need for an even broader "family" of skylines that would be useful in locations other than the Northwest. INT(1207), PNW(1211).

317. Simulation and other computer-based modelling systems provide an excellent means for examining numerous possible ways for solving complex logging problems. Recently completed studies, under a wide variety of assumptions and conditions, focuses attention on the potential for bringing about substantial cost reductions for commercial loggers. NE(1198, 1209), NC(1199), INT(1208).

J. Improving Water Quality and Yield

1. Improving water yields

318. It is estimated that 2,000,000 acre-feet of water is lost annually to evaporation from snow in the Sierra Nevada of California. Suppression of evaporation from snow may be achieved with use of long-chain-fatty alcohol monolayers. Twelve pounds per acre are sufficient to achieve maximum evaporation reduction. The chemical is not a pollutant and does not cause increased snowmelt at this rate of application. It does not migrate from point of application. The treatment is recommended for reduction of evaporation from high loss sites. PSW(42).

319. Information on the flow of water and dissolved solutes in trees provides valuable insights into tree growth processes under changing environmental conditions. Flow of radioactive tracers in 6-year-old lodgepole pines and red firs ranged fro 12 to 84 cm per hour. No water moved in lodgepole pine during the winter, whereas red fir transpired a small amount during that period. The procedures permit precise determination of conditions favoring tree growth, and provide a new means of estimating water use per acre. PSW(36).

320. Seasonal water depletion from a 1.5 m soil mantle averaged 188 mm on uneven-aged plots, 87 mm on clearcut plots, and 57 mm on plots without vegetation in a Wisconsin hardwood forest. Total amount of depletion increased with mantle depth; thus cutting had a greater water-saving effect on deeper soils. NC(38).

321. Sublimation from blowing snow is estimated from the balance of heat and mass transfer on a volume of air and nonuniform in particles. The effect of turbulence on convection around the particle appears unimportant in view of the small particle size. Turbulent transfer determines the snow concentration profile, and the gradients of heat and water vapor necessary to balance the sublimation process. RM(41).

322. A mathematical model for the sublimation of windblown snow was developed which predicts the distance a particle of given size will travel before completely sublimating; critical variables are particle speed, relative humidity and temperature of the air, and total insolation. Measurements of these conditions during all drifting events over the 1970-71 winter, indicate average transport distances of 460 and 900 m for particle diameters of 0.010 and 0.015 cm. respectively. RM(46).

323. A dynamic model has produced tentative results regarding the probable effects of timber harvesting on snowmelt in a central Colorado watershed. Snowmelt rates in mature lodgepole pine and Engelmann spruce forest were compared against a hypothetical pattern of 5 tree-height diameter openings, cut into the watershed so that trees on 40 percent of the area were removed. The estimated effect showed increased melt rates early in the melt season and lower snowmelt later. Although harvest cutting affected timing by causing the bulk of the snowmelt to occur earlier, the duration of the snowmelt season was apparently not significantly changed. Early-season snowmelt was accelerated in the small openings, but in both natural forest and cut-over areas, the last snow melted at about the same time. RM(33).

324. The efficiency of loblolly pine in eliminating sediment production from severely eroded and gullied waste lands is shown by an analysis of hydrologic measurements from experimental watersheds. Sediment yields 25 years after planting loblolly have been reduced to 0.03 ton per acre per year. The yields are a direct function of stormflow volumes. While loblolly reduces sediment to amounts approaching the geologic norm, the effect of reforestation on stormflows is largely a function of soil characteristics. Stormflows can be completely eliminated from small catchments of well-drained soils. SO(47).

2. Managing, rehabilitating, and improving watersheds

325. An open ditch hastened the drainage of water from surface or near surface horizons of peat on a typical lake-filled organic soil. The ditch had little influence on the water table beyond 5 m from the ditch when the water table was in

547-679 O - 74 - 3

deeper horizons of moderately decomposed peat. A similar treatment influenced the water table to a distance of 50 m laterally in an organic soil consisting of less decomposed peat. NC(57).

326. Little is known about quantitative relationships of surface form and water behavior of small forested watersheds. Three small contiguous Black Hills watersheds were analyzed in terms of elementary length dimensions, nondimensional elements, and similitude concepts. Maximum length of watershed and average length of first order basins scale most nearly the same as volume yield and stormflow peaks. It is theorized that these parameters are (1) primary controls over volume yield and peak flows and (2) better indicators of watershed behavior than watershed area. Refinements of this type improve the basis for small watershed systems analysis and simulation. RM(125).

327. Water yield increase following a wildfire in north central Washington averaged about 3.5 inches. The increase came mainly during the spring snowmelt and during summer months. Since the protective canopy was destroyed, maximum daily stream temperature was increased as much as 10° F. The temperature increases were well within limits prescribed for trout, however. PNW(80).

328. Mesquite not only redistributes nutrient elements within desert grassland ecosystems, but also provides a microenvironment that enhances availability of nitrogen and phosphorus. Pot tests disclosed differential availability of N, S, and P for three native perennial grasses in soil from under mesquite trees compared with soil from adjacent open areas. These measured differences in physical and chemical properties between areas appears to be the reason for greater abundance and improved perennial grasses observed under mesquite. PNW(120).

329. Trembling aspen root suckers originate from newly initiated meristems, preexisting primordia, or suppressed short shoots, but the first two are probably the most important. Evidence was found of suppressed short shoots in aspen roots but not of suppressed buds buried in the periderm. Vascular trace shows that the developing sucker responds to polarity in the root; it grows distally as it extends inward and basipetally to the root cambium. INT(110).

330. Streams require an additional number of gravel bars to adjust to slope. Average step length between logs and gravel bars are strongly related to channel gradient and median bed material size. More bars formed when fewer numbers of logs were available. Implications for forest management are that sanitation cuttings (removal of dead and dying trees) would not be permissible where dynamic stream equilibrium exists and bed material movement should be minimized. RM(79).

331. Soil ripping in 1963 effectively reduced runoff on the San 0Luis watershed of the Rio Puerco, New Mexico, and caused a favorable shift in forage production from galleta to alkali sacaton. Ripping effects on runoff are shortlived but forage production patterns may persist for 10 years. RM(50).

332. Preliminary production functions are identified for alternative land treatments in northern Arizona. These treatments are designed to increase water yield within a multiple use framework. Even without price information, a range of "best" input combinations can be identified. Product-product functions were developed for water, timber, and herbage for five strip cutting alternatives. These functions indicate the supplementary, complementary, and competitive outputs obtained from a given expenditure. Outputs and costs were evaluated over a 90-year period to account for the dynamic nature of production. RM(97).

333. Worldwide experience indicates that alder contributes significantly to the supply of nitrogen in the ecosystem0 A definite potential exists for employing alder in forest management in much the same way that legumes are utilized in agriculture. Current research indicates also that *Alnus rubra* may play a significant role in controlling *Poria weirii*, a virulent root pathogen which causes extensive losses of commercial timber tree species in western North America and Japan. PNW(117).

334. Overlapping, vertical aerial photos provide sufficient information for determining extent of snow cover in a small open area of mountainous terrain where accurate ground control is not available. Index measurements can be extrapolated to a much larger drainage. RM(76).

335. Snowmelt floods cannot be reduced by forest management practices in the Northeast. Snowmelt floods are regional in nature so management would have to be applied over very large areas to have any effect. Management of snow to produce more summer streamflow is also not feasible in the Northeast, since snowmelt runoff cannot be prolonged beyond spring. However, removal of conifers from a watershed will increase spring streamflow by up to 2 or 3 inches due to elimination of the interception loss. NE(72).

336. It is desirable to establish herbaceous and woody species in combination to stabilize spoil banks. However, herbaceous vegetation competes with tree species for light, moisture, and nutrients. Herbaceous vegetation planted on coal-mine spoil did not significantly affect the survival of trees but greatly suppressed their growth. By the fifth growing season, tree growth was suppressed by grass alone and was greatest in plots dominated by legumes. Competition from herbaceous vegetation can be lessened by encouraging legumes and avoiding strongly competitive grasses. NE(14).

337. The high wall is the dominant physical feature of most surface mining in mountainous terrain. High walls represent only 15 percent of the horizontal area disturbed and can be reduced by blasting or covering with spoil, but these measures may be necessary only in areas of high esthetic value. However, certain grading practices and placement of trees can reduce the visual impact of high walls by partially screening them from view. NE(101).

338. Black locust is one of the more commonly planted species on surface-mine spoils with a pH of 4.5 or less. Seedlings fertilized with nitrogen and phosphate, and triple-superphosphate in combination with ammonium nitrate are equally effective. Black locust should be fertilized with nitrogen and phosphorus to achieve rapid site protection on acid surface-mine sites. NE(102).

339. Relationships between forest and floods are complex, and often not properly identified. Forests provide the maximum opportunity for controlling runoff from flood-producing rainfalls, but forests cannot prevent floods. The principal source of flood runoff from forests is subsurface flow that moves rapidly downslope through the forest floor and topsoil. Forests provide the best possible natural cover for minimizing overland flow, runoff, and erosion. NE(93).

340. Interception loss was measured in loblolly pine stands and in a mature hardwood-pine forest in South Carolina. Annual interception loss from the hardwood-pine stand was similar to that of the pine stands. Where extensive conversions of hardwood to loblolly pine occur, significant reductions in the amount of water available for streamflow or ground water may be expected. SE(113).

341. Soil freezing did not change the infiltration rate sharply on natural oak-hickory forest and abandoned field

plots near LaCrosse, Wisconsin, until late winter. However in a 25year-old conifer plantation, the infiltration rate was reduced to near zero by early winter due to an impermeable snow-ice layer on the ground caused by snowmelt drip. Thus, conifer plantation may contribute more surface runoff than either deciduous forest or fields during snowmelt and winter rain. NC(77).

342. Contour trenches have been used for emergency protection of denuded mountain watersheds. Six years of soil water measurements on and adjacent to a contour-trenched area shows that some redistribution of soil water occurs following trenching. Trenching did not alter soil water conditions enough to significantly change water yields from the watershed. A reduction in water use from trench bottoms was offset by an increased loss from the cut-bank and trench fills. No change in soil water was apparent between trenches. INT(66).

343. Water quality management relates to problems of ultimate social importance. The technology is available to control large-scale changes in sediment, temperature, and chemical and bacteriological quality of water from forest land. In many cases, water quality considerations are also economic in terms of maintaining site quality, minimizing road mileage and maintenance, and reducing treatment cost. PNW(107).

3. Preventing watershed damage

344. Erosion by mass-wasting processes is often underestimated because some processes, such as soil creep and dry creep, are unspectacular and landslides occur infrequently. These forms of erosion make up a large proportion of the total erosion process. Mass movements are especially sensitive to disturbances by man such as during road building, logging, and vegetative manipulation. Terrain features can indicate the likelihood of slope failure. Using these features, the costs of repair can be estimated fairly well in advance. PSW(137).

345. A linear array of detonating cord was used to simulate a sonic boom. The boom was directed toward the fracture zone of a small avalanche path where the snow was unstable, as indicated by natural avalanches in the area. In three of four tests, avalanches were released by a boom of 12 pounds per square foot overpressure after withstanding lesser booms. One of the avalanches had a fracture face 8 feet 11 inches deep. RM(131).

346. Net radiation measured over a black spruce stand and a clearcut strip in a bog showed little difference in daytime values the first summer after logging. The second summer net radiation was lower in the clearcut strip than over the black spruce stand by 14 percent in June and 20 percent in September. While more energy was potentially available for evapotranspiration in the black spruce stand more energy was converted into sensible heat than in the clearcut strip. NC(127).

347. Some of the techniques which have proven most useful in management of wildland resources are multivariate and factor analysis. These techniques can help identify the complex physical, social, and economic variables which make up the management system by characterizing the management process as discrete treatment-response phenomena. PSW(133).

4. Water quality

348. Annual loss of nitrogen, phosphorus, silica, and the cations sodium, potassium, calcium, and magnesium followed the same pattern as annual runoff. Water passing through this Douglas-fir ecosystem conserved nitrogen effectively as indicated by an average annual dissolved nitrogen outflow of 0.5 kg/ha from an annual average input of 1.0 kg/ha in precipitation. Loss of nutrients by soil erosion may become of major importance over a longer time scale due to widely spaced unsampled catastrophic erosion. PNW(142).

349. Insecticides, herbicides, and fertilizers are important chemical tools in intensive management. Aerial application of these chemicals will result in their initial distribution among the air, vegetation, forest floor, and surface water compartments of the forest environment. Evaluation of the toxicity characteristics of forest chemicals indicates carefully controlled forest chemical applications will not significantly impact on stream organisms. PNW(145).

K. Bettering Silvicultural Systems

1. Natural regeneration

350. Basic knowledge on the growth and maturation of fruits and seeds of southern hardwoods is inadequate to support improvements on seed technology and use. Recent studies at the Forest Tree Seed Laboratory in central Mississippi have shown that sweetgum and sycamore fruits reach full size by late June and are physiologically mature by mid-August and early September respectively. Chemical changes in the two species were similar except that the major food reserves were carbohydrates in sycamore and fats in sweetgum. These results expand our knowledge on seed maturity and confirm the value of present seed maturity indices. SO(719).

351. There are several methods available for the determination of seed moisture content, but most have not been calibrated against a standard reference method. A recent study at the Forest Tree Seed Laboratory in Mississippi has shown that oven drying methods give moisture contents within 0.5 to 1.0 percent of the standard toluene distillation method for sycamore, sweetgum, and green ash. An electronic meter was also found to be suitable for sycamore and sweetgum. These results will make determination of seed moisture content simpler and more reliable. SO(720).

352. Information on factors affecting germination of swamp tupelo seeds is needed to develop methods of regenerating this species after a harvest. Tests under controlled conditions in the laboratory revealed that germination was rapid in moist, drained soil at 21 ° C, but did not occur in flooded soil. Thus, seedling establishment requires rapid germination and growth during the brief periods when conditions are favorable. SE(723).

353. Because of very limited timber cutting in the past, little is known about the natural reforestation of Sitka spruce on Afognak Island in Alaska. Observations made in one 25-year cutover area indicate that much of it is dominated by grass and herbaceous vegetation; growth of Sitka spruce seedlings is slow as a result of this competition. If these conditions are typical, special measures will be necessary to ensure adequate regeneration after logging on this unique Alaskan island. PNW(724).

354. Increasing use of shelterwood cutting in forests of coastal Oregon provides seedbeds that contain much more organic material and receive more shade than previously thought desirable for regeneration. Recent studies with seven coastal species show that organic seedbeds in the Sitka spruce—hemlock type, although desirable when exposed to full sunlight, are suitable seedbeds under the partial shade of a shelterwood cutting. The duff provides extra nutrients for the short-rooted seedlings that develop under these low light conditions. PNW(727).

355. If viable seeds of the desired species accumulate in the forest floor, they may ensure an adequate quantity of reproduction when favorable germination conditions are created by harvest cuttings. In a West Virginia study, black cherry seeds stored in the forest floor retained their viability over 3 years. Fewer than 10 percent germinated after the first winter, 50 percent germinated after two winters, and 25 percent germinated after three winters. Forest managers need not rely on current seed crops in scheduling harvests in black cherry. NE(732).

356. Little is known about seed production of paper birch in interior Alaska. A 5-year study near Fairbanks revealed that total seed crops during the period 1958-63 varied between 542 and 72,805 seeds per square meter; viable seed per square meter varied between 42 and 27,520. Seed crops adequate for natural regeneration of 100-foot-wide clearcuts occur in at least 1 out of 4 years in this portion of the taiga. PNW(733).

357. Forest managers in Alaska have available to them new information for managing white spruce. Results of research and experience in western Canada and Alaska were brought together to suggest a number of silvicultural treatments known to provide adequate conditions for natural regeneration. Similar summaries have been prepared for Alaska paper birch, quaking aspen, balsam poplar, and black cottonwood. PNW(683-685).

2. Silvicultural methods

358. Forest managers in the central Rocky Mountains are faced with the problem of modifying established timber harvesting practices to meet the increased use of forests for many purposes. Guidelines are now available to aid the manager in developing alternatives to clearcutting old-growth lodgepole pine forests. Alternative partial cutting practices provide for maintenance of forest cover in recreation areas, travel zones, and scenic view areas; other patterns of cutting give options for increasing water yields, improving wildlife habitat, or integrating timber production with other key uses. RM(734, 735).

359. Foresters have been challenged to harvest wood more efficiently while maintain a forested landscape while seedlings planted on the cleared strips are growing up. Several variations of this harvesting method are suitable for areas where natural scenic beauty must be maintained. NC(736-739).

360. Control of competing herbaceous vegetation is essential for successful planting of hardwoods on upland sites in the mid-South. A preemergence spraying with 2.5 pounds per acre of simazine reduced both herbaceous and grass competition by more than 80 percent in a trial in Louisiana without damage to the hardwood seedlings. This treatment was also effective in weeding pines, and appears to be worth testing in bottomland planting as well. SO(740).

361. A comprehensive review of current knowledge about the effect of clearcutting eastern forests on various ecological factors has revealed that much of the concern about clearcutting is largely unfounded. Clearcuttings do produce a temporary unsightly appearance that make them unsuitable for use in some locations. However, the complex environmental changes that occur after clearcutting favor the establishment of certain tree species; improve habitats for certain types of wildlife; produce a small increase in water yield; and, if properly applied, incur little risk of long-term environmental damage. NE(745).

362. Past harvesting of aspen has not always resulted in satisfactory regeneration because competition from residual trees and shrubs in many areas has inhibited the development of aspen suckers. Consequently the areas of merchantable aspen in the Lake States have been reduced. Competition on some of these areas has been minimized by mechanical harvesting, prescribed burning, felling residual trees, or by applying herbicides. The resulting stocking was much greater than on areas where competition was not reduced. Hence, clearcutting is essential for satisfactory regeneration of aspen. Production of merchantable aspen was increased further by thinning to about 800 trees per acre at 10 to 15 years and to 200-400 trees per acre at 20 years. NC(746, 747).

363. In response to the surge of national interest in forest management, a handbook has been prepared synthesizing present knowledge regarding biologically feasible systems of forest culture for harvesting, renewing, and maintaining 37 major forest types in the Nation. The handbook, entitled "Silvicultural Systems for the Major Forest Types of the United States," recognizes the importance of forests for wildlife, recreation, water, and timber and discusses the various systems of forest culture that can help meet these needs. Some of the systems are most appropriate for application on lands best suited to meet the Nation's need for forest products. 0The book also covers limitations in the use of certain systems imposed by species requirements, natural hazards, and the makeup of existing stands. It should help dispel uncertainties about silvicultural options and their consequences in the major forest types of the United States. WO(748).

3. Ecological relationships

364. A guide to the use of Research Natural Areas (natural scientific preserves) on Federal lands in Oregon and Washington has been compiled which describes the physical and biological features for each of 45 reserved tracts. Maps and photographs are included. Capsule summaries for each area include establishment data; size; administering agency; location; availability of special maps; environmental, biological, historical features; and history of research. The guide will be a key reference for scientists and educators who wish to study the biota and ecological processes of selected natural environments. PNW(751).

365. Recent concern about endangered species of plants has prompted a reevaluation of the principle of competitive exclusion—the theory that two or more competing species cannot coexist indefinitely. In a New Hampshire study of simulated forest stand development, yellow birch steadily replaced paper birch, a phenomena that was confirmed by actual species representation in a virgin stand of northern hardwoods. The real possibility of extinction of paper birch in undisturbed stands illustrates the need for intensive management to assure a future supply of this highly valuable and aesthetically desirable species. NE(752).

366. Fine roots are often neglected in studies of forest biomass because of the difficulty of separating finely divided, widely dispersed roots from soil samples. In a white spruce plantation in Maine, the biomass of roots 3 mm in diameter and smaller represented as much as 20 percent of the total root biomass. This fact must be taken into account by ecologists when estimating total tree biomass or standing crop. NE(755).

367. Knowledge of differences in drought resistance among the oaks is needed to fully understand where the different species should be grown. Recent studies in Missouri show that leaf and root cells of post oak are better able to survive dehydration than those of black, white, or red oak. Red oak

was least successful at surviving dehydration, and it was not able to postpone dehydration as well as the other species. Thus, post oak is the most drought resistant, and red oak the least resistant of the species studied here. NC(756).

368. Environmentalists are concerned about the possible changes in productivity and composition of the earth's forests that could be brought about by the ever-increasing amounts of carbon dioxide, particulates, and gaseous pollutants in the atmosphere. Although the direct effect of a higher carbon dioxide concentration is an increase in vegetative productivity, this increase would be partially offset by reductions in sunlight intensity and air temperature associated with increasing particulates. Drift of particulates could change the amount and distribution of precipitation but the effect of changes in this factor on overall productivity cannot be predicted. In general, any climatic changes that might occur are likely to be deleterious because tree species are adapted to a specific climate. NC(757).

369. Knowledge about structure and functioning of natural ecosystems in the taiga of Alaska is meager. The distribution of selected nutrient elements in portions of 5-, 15-, and 20-year-old alder ecosystems developing on the Tanana River have been studied and described. Changes in nutrient distribution with time between soil and plant materials are essential items of information for understanding the function of an ecosystem. PNW(686).

L. Remote Sensing Methods

370. The linking of remote sensing data for wildland resources with appropriate computer programs will permit data analysis, storage, updating, and rapid retrieval by planners, policy-makers, and managers in many resource fields. Many information systems and automated procedures for land use classification are in early stages of development and should be more fully utilized in research and in practice. PSW(428-431).

371. New methods have been developed for handling the digital calibration data and sensitometric analysis of color films. This will permit computer analysis and improved classification and pattern recognition. PSW(425,427).

372. Spectral signature measurements using multiband camera systems require precise calibration of camera shutters. A technique has been developed for measuring the exposure times of focal plane shutters with special emphasis on aerial cameras. Any user of aerial cameras can now adjust or calibrate the exposure times without removing the camera lens. PSW(426).

373. There is increasing interest in forest and nonforest land classification for using imagery taken from high altitude aircraft and from space. Photographs taken over a 20-month period and covering eight 16-square-mile areas near Atlanta, Georgia, were correctly classified in more than 75 percent of the cases. Use of smaller scale, high-altitude color infrared photography significantly increased area coverage without sacrificing accuracy. Substantial savings in operational costs were also obtained. PSW(423).

374. A low-cost projection-viewer has been developed to enlarge parts of microscale aerial photographs. These can be used for assessment of insect- and disease-caused damage, type mapping, monitoring environmental changes, or other purposes. PSW(424).

M. Weather Modification and Weather Effects

375. Forest Service scientists are cooperating with the BLM and are evaluating their Alaskan lightning abatement program. This work will influence future USDA lightning modification efforts. INT(687).

376. Continued research strongly supports the hypothesis that forest fires are caused by a specific type of discharge. This work provides a sound physical basis for predicting the potential of fire ignition and is a major step in developing a natural risk index for the National Fire-Danger Rating System. INT(691).

377. Lightning fire statistics for southeastern Georgia indicate that while some thunderstorm days produce lightning-strike fires others do not. It was found that thunderstorm days with lightning-strike fires were characterized by higher values of buildup index and spread index than fire-free days. SE(694).

378. Studies of large outdoor propane fires have resulted in a spin off. Halon flame extinguishers can be used to suppress explosion hazards. This suggests a safety action that could be taken when spraying LP gas for fog seeding. PSW(693).

379. New mathematical and statistical techniques have been developed for generating families of probability distributions for analyzing weather patterns and for explaining and predicting wind flows. SE(688,695,696), RM(690).

380. Observations in the meso scale of the Cascade Mountains identified strong gale force winds, instability lines, and squalls that were not identified by routine synoptic analyses. These are important weather influences which effect ignition and spread of forest fires. More attention needs to be placed on meso scale systems for identifying and warning fire control officers. PNW(689).

381. Scientists from the Macon Laboratory report that passage of a cold front and the dynamic action of the high-level jet stream were responsible for most of the large fires that occurred in North Carolina on a major fire day, December 4, 1970. Eight southeastern States had 1167 fires which burned 43,294 acres. SE(697).

382. The effects of weather on seed production of loblolly pine were shown by a study from 1947-1968 on the Hitchiti Experimental Forest. Temperature and evaporation shortly after pollination in May were dominant critical variables. Rainfall in July appeared to be a significant factor. Cool moist weather in March hindered fertilization while warm, dry April weather favored it. SE(692).

II. IMPROVING UTILIZATION AND EXTENDING WOOD SUPPLIES

A. Intensive Culture Methods

1. Site evaluation and soil improvement

383. The reliability of plant analysis as a technique for predicting fertilizer needs of forest crops depends upon careful standardization of sampling procedures to control the many variables that affect nutrient composition. A study of 60-year-old even-aged oaks in West Virginia showed that easy-to-reach foliage from the lower crown is suitable for foliar nutrient composition and dry weight determinations. Application of these findings should reduce the high cost of leaf sampling and promote the effective use of foliar analysis for diagnosing the fertilizer requirements of forest stands. NE(758).

384. To achieve the maximum growth increase at minimum cost from fertilizer applications, it is essential that plant responses to various nutrient combinations be known. A recent study in West Virginia has shown that red oak seedlings will not respond to phosphorus applications unless nitrogen requirements are also satisfied. Likewise, potassium produces a small additional response only in the presence of high levels of both nitrogen and phosphorus. Red oak fertilization on soils similar to those studied here should always include nitrogen and phosphorus but additional potassium is often unnecessary. NE(759).

385. The growth of Ocala sand pine in some of the sandhill plantations in Florida is limited by deficient amounts of available soil phosphorus. A greenhouse test revealed that seedling growth was greatest in soil containing 1.0 to 1.5 ppm of available phosphorus. Phosphate fertilization, therefore, will increase the growth of planted Ocala sand pine only on soils containing less than 1 ppm of available phosphorus. SE(760).

386. Evaluation of fertilizer needs in forest trees requires a background of knowledge on nutrient requirements of the species involved. Several studies in Maine have shown that paper birch seedling growth is much more sensitive to nitrogen supply than to supply of phosphorus or potassium. Foliar nitrogen levels of 3 to 4 percent indicated adequate supplies of this nutrient. NE(761,762).

387. Silty old-field soils on the Mississippi River flood plain are often depleted of nutrients and are marginal for the production of cottonwood fiber. However, growth of a 6-year-old cottonwood plantation was increased by as much as 200 percent with nitrogen fertilization (168 kg/ha) in a recent study. Thus, fertilization should permit utilization of some of these marginal sites. SO(763).

388. Because rhododendrons generally thrive on acid soils, the application of lime may be detrimental to this plant. However, application of as much as 35,866 kg of agricultural limestone per hectare caused no visible reaction in a rosebay rhododendron stand in the southern Appalachians of North Carolina, even though pH in the surface 2.5 cm was increased to as much as 7.1. Thus, liming of soil for other purposes probably will not harm rhododendron, nor does it offer promise as a means of eradicating rhododendron in places where it is not wanted. SE(765).

389. Regeneration and growth of water tupelo and baldcypress in the swamp forests of the southeastern States have not been satisfactory after harvest cutting. In a greenhouse study of potted seedlings, growth was better in saturated soil than in unsaturated soil, but only when water in the unsaturated soil decreased substantially below field capacity between waterings. Growth was increased further by fertilization with solutions of urea. These results can be used in prescribing drainage and fertilizer practices to improve growth of trees planted in these swamps. NC(766, 767).

390. Land managers in the Ozark Highlands of Arkansas and Missouri need techniques to evaluate potential pine growth on sites where there are now no pines for growth measurements. A study of soil and site relationships showed that, in this area, site index for shortleaf pine can be predicted from slope convexity, aspect, elevation, and degree of oak and hickory competition. SO(768,769).

391. Yellow birch trees growing on acid, podzol soils in New England contain very high concentrations of manganese. A recent New Hampshire study shows that these concentrations, which range up to 4500 ppm, are much higher than those required to produce toxicity in seedlings grown in nutrient solutions. Although other elements present in forest-grown birch may alleviate the effects of high manganese to some extent, the possibility of toxicity must be considered in any future research on birch nutrition. NE(771).

392. Black walnut can be expected to survive well and grow rapidly only if planted on suitable soils. A recently published compilation of the soil series occuring in Illinois, with classification of the suitability of each, will assist in selecting the better sites for planting this valuable hardwood. NC(773).

393. Extensive flooding in the Mississippi Valley has raised concern over effects of flooding on forests. After floods deposited 2600 tons of silt on 2.8 acres of a black walnut plantation in north central Missouri, extensive mortality and growth reduction resulted. Damage was closely related to the thickness of the silt deposits and was more harmful after spring than fall deposition. This episode demonstrates the need for erosion control practices and for more careful site selection. NC(774).

394. Site index for young ponderosa pine stands in California cannot be estimated accurately from existing curves. But recent studies have shown that estimates can be improved either by adjusting the existing curves or by estimating site index based on total length of the first four internodes above breast height. These methods materially improve estimates of site index in young stands and improve forecasts of forest productivity. PSW(775,777).

395. Guidelines on when to fertilize black walnut plantings have been inadequate to ensure that maximum growth or nut production will be obtained. Research has now shown that foliage symptoms and foliage nutrient concentrations can serve as quick guides to nutrient status of black walnut. These guides and suggested procedures for using them have recently been published for use by walnut growers. NC(776).

396. Site index curves for California black oak, previously unavailable, have now been developed. The curves have been related to slope aspect and to site index of a close associate—ponderosa pine. For the first time, these curves provide a means of estimating the growth potential of California black oak in even-aged stands on the Sierra Nevada and southern Cascade Ranges. PSW(778).

397. Prediction of site quality would be greatly facilitated if it could be determined from soil types. However, analyses of soil-site plots in the Missouri Ozarks failed to reveal any relationship between soil types and oak site index. Furthermore, site index was not related to the presence or absence of a fragipan. The variability in site quality found here is apparently due to factors not delineated in current soil classifications. NC(779).

398. Information on nutrient cycling is essential for effective soil management. In the Piedmont area of North Carolina, total litterfall was 25 percent greater in a mixed hardwood stand than in a loblolly pine stand with a hardwood understory. Hardwood litter contained about twice as much K, Mg, Mn, and Cu as pine litter. Results indicate that the hardwood understory of the pine was quantitatively important in nutrient cycling. SE(780).

2. Artificial regeneration

399. Previous studies in India have shown that silver fir seedlings do not readily regenerate naturally in deep raw humus unless the roots reach mineral soil and mycorrhizae develop. To avoid these problems associated with natural regeneration, artificial regeneration methods are now being tried. Seedlings with mycorrhizae are first grown at low-elevation nurseries and then planted in the mountains. Although growth rate was not increased, the method of growing planting stock at low elevation in India is still of some benefit since survival was much better. WO(781).

400. Seed dormancy is a problem in both direct seeding and nursery culture of southern pines. Research in Louisiana has shown that in loblolly pine that seedcoat seems to delay germination more by mechanical constraint than by limiting water or gaseous exchange. Stratification acts to increase the capability of the embryo to rupture the seedcoat. This information helps to clarify the basic causes of dormancy, and will be useful in finding ways to improve the germination efficiency of pine seeds. SO(782).

401. Can loblolly pine seed that has been stratified to overcome dormancy be successfully returned to storage if it is not all used? An experiment in Louisiana showed that viability of vigorous seed lots remains high, but they gradually regain their dormancy, whereas lots of lower vigor store poorly and become more dormant than untreated seed. This information will be helpful in deciding on disposal of left-over stratified seed. SO(783).

402. A knowledge of pollen phenology and dispersal pattern is important for seed production management. In a natural stand of shortleaf pine, pollen collected on celluloid tape traps indicated significant differences in density of pollen for years, location, elevation, and trap orientation. Annual variation was extremely large and could be a major factor reducing seed yields in some years. SE(789).

403. Strobilus development in shortleaf pine was studied in detail to determine the cause for low seed set in the Virginia Piedmont. During a 6-year period strobilus survival ranged from 3 to 65 percent with an average of only 29 percent. The major factors causing strobilus mortality were found to be spring frosts, insects, and physiological abortion. These factors must be controlled in seed orchards in this region if high yields of seed are to be produced. SE(790).

404. A quantitative estimate of annual seed efficiency in shortleaf pine cones was needed to evaluate the low seed production of this species. An analysis of mature cones for 6 years indicated that seed losses are of three major types: (1) small first-year aborted ovules, (2) sunken and resin-filled second-year aborted ovules, and (3) unsound (empty) seed. Based on the potential number of seed per cone the seed efficiency was only 14 percent. There is an obvious need to reduce these various forms of ovule abortion if seed production for this species is to be greatly increased. SE(791).

405. When planting pines in the Piedmont—for timber production, Christmas trees, or highway beautification--land managers must choose among the many species available. In a South Carolina study, loblolly pine grew faster than five other species tested, while slash pine grew nearly as well and can probably be substituted for loblolly with little growth loss. Shortleaf pine may also be suitable for timber production but had slightly poorer growth and survival than the other two speices. Virginia, longleaf, and eastern white pine performed so poorly that they would be uneconomical for timber production but should be suitable for other purposes. SE(792).

406. Unmerchantable scrub oaks and wiregrass dominate millions of acres of sandhills in the Southeast. Conversion to pine, however, probably will be profitable. A new USDA Handbook brings together the results of 45 years of research on methods of reducing competition for soil moisture and nutrients that will enable planted pines to survive and grow under the severe stress of the sandhill environment. Practical site preparation methods, both mechanical and chemical, are described. Loblolly pine and slash pine have proven suitable for planting on the more moist sites, but Choctawhatchee sand pine grew better on the drier, more infertile sites. SE(793).

407. Nursery production of the large stock required for successful reforestation of difficult Douglas-fir sites is expensive. A recent study in Washington has shown that 3-year-old seedlings are cheaper to produce than the usual transplants of the same age, even though the seedlings require more bed space. For optimum results, production of 3-year seedlings at a density of 8 to 10 seedlings per square foot is recommended. PNW(795).

408. Cutover jack pine stands on sandy soils (Grayling Sands) in the northern half of lower Michigan have not regenerated satisfactorily. In trials of various site preparation methods, best stocking was obtained on plots that were scarified with an Athens disk 2 weeks after burning and then seeded at the rate of 20,000 jack pine seeds per acre. NC(796).

409. A natural hybrid between West Indian (*Swietenia mohogani*) and Honduras (*S. macrophylla*) mahoganies in the United States Virgin Islands reputedly combines the drought resistance and excellent wood quality of West Indiana mohogany with the faster growth of the Honduras species. Ten years after planting, in a replicated trial on three sites in St. Croix, the hybrid had survived better and grown faster than either parent species. The hybrid mahogany is therefore recommended for plantations of this valuable hardwood species in the Virgin Islands, except on very dry sites. ITF(800).

410. Valuable tropical timber trees from other countries have great potential for supplementing the native flora of Puerto Rico and similar areas. Extensive adaptability trials in the sandy well-drained uplands show that the following introduced species have the most promise: Honduras pine for most landowners; mahoe (*Hibiscus elatus*) for a cabinet wood;

kadam for a general utility wood or for areas with severe weed problems; and eucalpytus or casuarina for post or pole crops. ITF(797).

411. Reliable methods are needed to reforest Allegheny hardwood areas that fail to restock naturally after cutting. Studies of direct seeding of black cherry in northwestern Pennsylvania show that fall sowing of fresh seed (before October 1) and spring sowing of stratified seed are equally effective. Seedbed cultivation is unnecessary, but proper seed handling and covering, control of weeds and grasses, and protection from browsing in areas of high deer populations are critical. NE(803).

412. Less expensive techniques of establishing stands are always desirable, and will be especially important for short-rotation cultural schemes where very dense stands are needed. Recent studies in Georgia with sycamore and in Mississippi with green ash have shown that cuttings will sprout when planted horizontally in furrows, providing a technique of establishing new stands that should be economical because it can easily be mechanized and requires a minimum of plant material. SE(808), SO(805)0.

413. Knowledge of the meteorological conditions that favor the development of loblolly pine seed is needed for predicting seed crop size. In Georgia, seed production was favored by a low mean temperature and a low evaporation rate during the month of May when pollination occurred; and by higher than average rainfall in July when flower primordia were differentiated. Hence seed production in loblolly pine orchards could be increased during drought years by irrigation in May and July. SE(806).

414. Forest landowners are concerned about slow diameter and volume growth in dense even-aged stands resulting from direct seeding. In an 11-year loblolly pine stand in central Louisiana, strip thinnings at 3 years of age were as effective as selective thinnings in increasing diameter growth of the residual trees. Power equipment can be used to do precommercial strip thinning quickly and at low cost. SO(807).

415. Seeds of white ash require exposure to a cold, moist environment for 60 days before they will germinate. This cold period was reduced to 30 days by presoaking the seeds for 12 hours in a solution containing 1 ppm of gibberellic acid. This reduction can speed up seed germination tests which are necessary to determine sowing rates in nursery beds. NC(809).

416. Seed yields of the southern pines are usually calculated in terms of pounds of seed per bushel of cones. The number of seeds per cone can be determined by slicing, but the number of cones per bushel varies greatly both within and between species. A study in Louisiana showed that the average cubic volume actually occupied by the cones in one bushel is relatively constant regardless of cone size. Using these averages—10,068 cubic centimeters per bushel for longleaf, slash, and shortleaf cones, and 18,162 cubic centimeters for loblolly cones--the number of cones per bushel can be estimated quite accurately by measuring the average volume of a few sample cones. SO(810).

417. The feasibility of planting needs to be evaluated as a means of regenerating northern red oak in young natural stands where it is now scarce. Seedlings planted in the Southern Appalachians of North Carolina survived well but their height growth has been unacceptable in spite of annual removal of competition. These results suggest that successful plantings will require much more intensive culture if planted oaks are to outgrow their competitors. SE(811).

418. Although container planting should provide better survival and subsequent growth than conventional methods, the amount of improvement must be determined before the method can be fully evaluated. Containerized Douglas-fir and noble fir planted in the Oregon Cascades averaged 2 1/2 times taller and had significantly higher survival than bare-rooted stock after one year. Widespread field trials underway will provide additional data over the next few years, but the method appears to have great potential. PNW(812).

419. Early attempts to plant and culture cottonwood in the Upper Mississippi Valley have not been very successful. Recent studies show that planting large seedlings to a depth of 24 to 30 inches will result in better survival and growth than use of cuttings or conventional planting of nursery-run seedlings. Site preparation and weed control are also essential. NC(813).

420. The relatively short growing season in northern latitudes causes difficulties in nursery production of seedlings that are large enought to plant when 2 years old. First-year survival of outplanted 2-year-old seedlings of white spruce and jack pine grown in the more favorable environment of an unheated plastic greenhouse was 60 to 100 percent greater than those grown in an outdoor nursery in Michigan. These results indicate the need for testing the effects of a plastic greenhouse enviornment on other northern species. NC(814).

421. Many failures of Engelmann spruce plantations in clearcut openings in the central Rocky Mountains have resulted from careless handling of nursery seedlings as well as from adverse environmental conditions after planting. Guidelines are now available that will provide the forest manager with the techniques needed to successfully plant spruce. Included are the kind of stock to plant, time of planting, transportation and storage, selection of planting spots, site preparation, planting methods, and plantation protection. RM(815,816).

422. The amount of animal depredation of direct-sown seed must be known so that adequate protection can be provided where needed. In a study of spot-seeded yellow-poplar in central Tennessee, it was found that screen-protected seedspots had three times as many seedlings as unprotected seedspots, but a sufficient number of the exposed seed escaped damage to provide fair stocking. Thus, unprotected seed can provide acceptable yellow-poplar stocking if a large number of seedspots are used. SO(818).

423. To improve chances for successful conifer regeneration in the Southwest, a provisional tree seed-zone and cone-crop rating scheme was developed. The forested areas of Arizona and New Mexico are divided into 10 physiographic-climatic regions, each with a number of 50-mile-wide seed collection zones. The system includes a 10-unit scheme for rating cone crop production. Collection zones will be used to define limits for segregating seed collected thus avoiding costly off-site planting errors; the cone-crop rating scheme will assist in planning regeneration cuts. RM(821).

424. A comprehensive summary of technical information on reforestation practices in California has been compiled which includes virtually all of the significant findings related to this subject during the past 60 years. It includes specific and detailed information on cone and seed handling practices, nursery practices, site preparation, planting and seeding. For the resource manager and timber grower in California this is a reference of inestimable value for planning and managing reforestation programs. RM(820).

425. To satisfy the demand for genetically improved seed, efforts are being made to stimulate cone production in seed orchards while trees are young. In a slash pine orchard in northern Florida, 168 kg of N per hectare was applied at 6

years and again at 9 years of age. This fertilization schedule doubled the cone crop at 8 years of age and tripled it at 11 years to an average of 66 cones per tree. SE(823).

426. It is frequently assumed that sufficient mycorrhizae will develop naturally to enhance the survival and growth of pines planted on soils containing the requisite fungi. However, in an experiment in Louisiana seedlings grown in inoculated soil had four times as many bifurcate roots (e.g. visible mycorrhizae) as did those grown in uninoculated soil, and 1/3 more of the inoculated seedlings survived when outplanted. Initial field survival was closely correlated with both the number of bifurcate roots at lifting and nitrogen and phosphorus levels in the plants. The results indicate that mycorrhizal development in the nursery can be increased by growing seedlings at uniform spacing in low-density beds that are fertilized at a moderate rate. SO(824).

427. Comparative performance data are needed to guide the selection of species for planting on various sites on the Cumberland Plateau. In a comparative test in Jefferson County, Alabama, loblolly pine outgrew Virginia pine at all topographic positions, and Virginia pine in turn outgrew yellow-poplar at all positions except in the bottoms. The study suggests that loblolly is the preferred species on most sites, with alternatives for Viginia pine on warm slopes and shallow soils, or yellow-poplar on lower slopes and bottoms. SO(825).

428. Douglas-fir regeneration is often unsatisfactory on hot, dry south-facing slopes in northwestern California. A recent study has shown that shade will improve germination and survival if direct seeding is used, but shading did not affect survival of planted seedlings. Thus, some means of providing partial shade—such as a nurse crop--should be used if these sites are to be direct seeded. PSW(827).

429. Kadam (*Anthocephalus chinesis*), an extermely fast-growing broadleaf species, appears very promising for industrial wood plantations in the tropics. Two simple methods of extracting the seeds have been developed in Puerto Rico, by macerating either dry or wet fruits. The seeds, separated by sieving, are 98 percent pure, and more than 90 percent germinate. ITF(828).

430. Black walnut seed is difficult to store. However, recent seed storage studies in Indiana show that walnut can be stored for at least 2 years in either deep outside pits or in cold rooms, if proper attention is given to seed moisture content. These results will help reduce annual fluctuations in walnut nursery production by reducing dependence on current seed crops. NC(834).

431. The search for nursery techniques that might improve subsequent growth and survival of black walnut seedlings has led to the suggestion that fibrous-rooted seedlings might perform better than normal tap-rooted seedlings. Root pruning was found effective in developing fibrous roots, but field plantings in Illinois and Indiana have not shown the expected advantages of fibrous roots. Instand, stem diameter appears to be a more important indicator of early height growth. NC(835).

3. Stand improvement

432. Response of trees to particular stocking levels often depends upon the condition of the trees or stand prior to thinning. Studies in Oregon show that 30-year-old ponderosa pine plantations containing trees with well-formed and full crowns can be reduced to as few as 140 trees per acre and still produce the same volume as stands containing four times as many trees. But small-crowned trees do not respond if thinned heavily, and stands containing dense understories may also fail to respond to heavy thinning because of competition. Thus, heavy thinnings in ponderosa pine should be limited to stands of large-crowned trees where understory vegetation is sparse. PNW(838).

433. Herbicidal sprays applied in drops large enough to assure effective placement on target species and prevent drift to nearby susceptible plants are usually less effective than when applied in fine mists. Research in Louisiana with radioactively labelled 2,4,5-T showed that the amount of chemical absorbed by the leaves of six hardwood brush species is proportional to the logarithm of the drop volume. This information should allow savings of 1 to 4 gallons of spray per acre with thickened sprays when compared to the old "rule of thumb" of 72 drops per square inch. SO(842).

434. Natural reproduction on moist sites at intermediate elevations in northern Idaho usually is a dense mixture of 4 to 8 coniferous species in which control of species composition has been difficult to attain. Recent observations on experimental plots installed 35 years ago, however, revealed that the best control of species composition was obtained by harvesting the overstory and cleaning out unwanted species before the young stand attained an age of 30 years. With this information, land managers can change the composition of the stand to favor development of species suitable for the planned use of such sites. INT(843).

435. The possibility that epicormic sprouting will degrade butt logs is an important deterrent to thinning in even-aged yellow-poplar stands in the southern Appalachians. In a North Carolina study, there was much individual tree variation in the production of epicormic sprouts, but the incidence was low and could not be correlated with thinning intensity, number of residual trees, age, or site index. Thus, thinnings can be made without incurring serious degrade due to epicormic sprouting. SE(844).

436. Research on tree development after thinning, fertilization, and other silvicultural treatments have shown that growth rate can be increased by release and often enhanced by the addition of nutrients. Using a new X-ray method for the rapid determination of wood density distribution, a study in California in a thinned 46-year-old ponderosa pine seed production stand indicated that unthinned trees decreased in growth rate and had a slight increase in wood density. In contrast, thinned and fertilized trees increased by 33 percent in growth rate and 3 percent in wood density. Although thinning with fertilization did not necessarily contribute to wood uniformity, these commonly applied cultural practices were beneficial by increasing growth. PSW(845).

437. How much should an individual tree be released to benefit from thinning, and for how many years will the effect last? In a 70-year-old white spruce stand in Maine, trees released on three or four sides grew twice as fast in diameter as the controls or those released on only one or two sides, and the effect was still evident 10 years after thinning. These results will help in prescribing silvicultural treatments for the many acres of pole-sized overstocked stands of even-aged spruce in the Northeast. NE(846).

438. Many different techniques have been developed in attempts to control unwanted drift of pesticide sprays from target areas. A recent summary of aerial spray additives for control of herbicide drift has just been published to assist foresters in choosing an appropriate technique. Materials covered include invert emulsions, particulating agents, thickening agents, and foam sprays, of which the latter two show special promise for forest use. PNW(847).

439. Injection with herbicides is a common method of removing undesirable hardwood trees from southern forests, but there is a continuing need for more effective and economical chemicals. Research in Louisiana showed that undiluted 2,4-D amine is effective and economical on many species, and that on those resistant to this chemical alone the addition of picloram or picloram and 2,4,5-T will give satisfactory results. Detailed recommendations have been developed for various species in terms of chemical mixture, dosage, and spacing of injections for best results. SO(849-851).

440. Selection of hardwood crop trees in young stands is often limited because many of these trees have sweep or crook in the lower stem. But a recent study of green ash saplings in Illinois has shown that considerable natural straightening of the stem will occur after release as a result of faster diameter growth on the inside of the crook. Thus sapling stems with a considerable amount of sweep and crook are acceptable as crop trees. NC(852).

441. Firetree, an agressive plant pest in Hawaii, can be controlled by herbicides but the falling limbs and stems of the dead trees reduce the form and quality of planted replacement species. Cutting the damaged trees and removing all but the dominant sprouts on each stump replaced the cull trees with valuable growing stock. Australian toon and tropical ash, because of their strong tendency to produce vigorous sprouts, gave the best response to treatment. Use of this method can maintain the volume and quality yield potential of plantations without costly replanting. PSW(854).

4. Animal damage

442. Although the protective coating generally recommended for Douglas-fir seed in the Pacific Northwest includes no Arasan, many foresters have begun including it in recent years. A reevaluation of the various components of the seed coating substantiates earlier findings that the addition of Arasan reduces germination without providing improved protection from deer mice. Since birds are not serious predators in the Pacific Northwest—as they are in the South where Arasan is used extensively--there appears to be no justification for adding Arasan to Douglas-fir seed coatings. PNW(855).

443. Feeding by various animals, expecially deer and hare, can seriously damage Douglas-fir regeneration. Chemical analysis and fermentation studies of foliage from clones and progenies with known and unknown susceptibility ratings to browsing revealed genotypic differences in tissue digestibility, in effects of essential oils on cellulose fermentation, and in chemical composition. Some of these resistance characteristics are transmittable through breeding. Breeding for resistance to browsing is of special importance now because of present objections to the use of synthetic chemicals and the high cost of mechanical protection. PNW(857).

444. Chlorogenic acid—a phenolic compound that may affect seed production and susceptibility to animal damage--has been isolated for the first time in the foliage of a north-temperate conifer, Douglas-fir. Concentrations of the compound varied among genotypes, suggesting the chlorogenic acid content may be a useful variable in tree breeding programs. PNW(858).

5. Growth requirements

445. Reasons for the distinct type lines which often occur between lodgepole and ponderosa pine stands in south-central Oregon are not clearly understood. Soil water levels close to saturation favored germination of lodgepole over ponderosa pine but had no differential effect on survival and growth of the two species during the early seedling stage. Exclusive occupancy of "wet sites" by lodgepole is thus not the result of differential species tolerance to mineral soils with high water contents as earlier suspected, unless such tolerance in ponderosa pine changes with age. PNW(862).

446. Early growth response to temperature differentials and soil fertility levels is not known for lodgepole and ponderosa pine in south-central Oregon. Fertilization at four levels increased growth of both species under nine different temperature regimes studied. Lodgepole pine growth was not as sensitive to night temperature changes of 8° to 16° C as was ponderosa pine growth. The number of daily degree hours necessary for maximum growth increased with increasing soil fertility level for night temperatures of 16° C. These findings demonstrate the importance of the fertility of the rooting medium when studying temperature effects on seedling growth of these two species. PNW(861).

447. It would be useful if selected cells which make up wood could be controlled or at least modified as wood is developed. Studies to determine if this is feasible are now under way by Israeli scientists supported by the United States PL-480 program. They are attempting to determine how selected cells originate and differentiate into wood. For example, they recently found in tree-of-heaven that the differentiation of the vascular rays is not controlled by the cambium but by associated tissues producing or consuming selected stimulating substances. Thus it may be possible to control the vascular ray development and perhaps other cells by externally controlling hormone production. WO(863,872).

448. Several species of the widely occurring fungus genus *Hebeloma* are thought to be ectomycorrhiza-formers. Recent joint studies between the Forest Service and the Claude Bernard University in France demonstrated that 16 different European species of the genus *Hebeloma* were able to form a mycorrhizal relationship with Virginia pine of the United States, while five species could not. These results suggest that marked physiological differences do exist between the different fungal species and that this type of data can be used to further the taxonomic characterization of this fungus genus. NE(865).

449. Mycorrhizae are symbiotic associations in which the smallest secondary roots of higher plants are invaded by specific fungi, and the roots retain their vital characteristics without pathological symptoms. Mycorrhizal fungi improve uptake of nutrients for host plants, protect roots against pathogens and move carbohydrates from one plant to another. Forest trees could not survive in their natural soil habitat without mycorrhizae. The little-known role of mycorrhizae in life cycles of the associated fungi and their hosts have now been summarized with data from the world's literature. NE(864).

450. Mature woody stems are hard and often dry or resinous and contain few living cells per unit volume. Enzyme studies of such material is difficult if not impossible. A new enzyme extraction method of woody plant material involving the homogenization of milled lyophilized tissue has been developed which is both reproducible and physiologically sound. The method has wide application in both biochemical and physiological investigations of tree tissues. NC(867).

451. Our understanding of the regulation of indoleacetic acid (IAA, a growth-regulating hormone) levels in plants is hindered by lack of knowledge on the location at which IAA oxidase activity occurs. An experiment in New Hampshire reveals that IAA oxidase and peroxidase activity occur

together, indicating that both are functions of a single enzyme. NE(869).

452. Habitat classifications and descriptions are needed in the mixed conifer zone of the Southwest to delineate where various tree species are apt to grow best. Dry-season moisture stresses in wild Arizona mixed conifer seedlings less than 15 cm tall were higher than those for larger seedlings. Night time recovery was complete. Stresses did not increase through the dry season and their day-to-day variability seemed a function of day-to-day weather differences. Species differences were small but significant. Although inconclusive for classifying regeneration habitats in Southwestern mixed conifers, these findings support the avoidance of small planting stock and the selection of those species with lowest stem moistures stress that are otherwise adapted to the planting site. RM(871).

453. Many landowners who grow black walnut for nut production are overlooking the timber values that could be obtained from these same trees. Research in Illinois shows that by planting more trees per acre, pruning side branches until a 9-foot log is produced and thinning at appropriate times, landowners can produce high-value walnut logs and still obtain good nut yields. NC(873).

454. One promising lead for genetic improvement of cottonwood is to select trees that retain their leaves longer than average. Annual growth on such trees is prolonged over a longer portion of the frost-free season. These late season increments are of high quality because they consist of wood fibers with very few or no conducting vessels. Thus leaf retention provides one index of selection for wood quality in cottonwood. NC(875).

455. Leaf surface area is a very good measure of photosynthetic productivity which determines wood yield, but leaf area measurements are difficult and time consuming. On two clones of hybrid poplars, leaf length proved to be a simple, nondestructive substitute for leaf area measurements. Hence the relative ability of various poplar clones to produce wood can be estimated by measuring their aggregate leaf lengths. NC(876).

456. Young developing leaves of many broadleaved species are both importers and exporters of the plant foods formed by photosynthesis, but the pathways for such bidirectional transport are not known. Experiments on young leaves using radioactive carbon dioxide revealed that plant food is photosynthesized in the mature tips and exported to other parts of the plant with a simultaneous import from other leaves to the immature basal region of the leaf. Transport from the mature tip directly to the base does not occur. This result is a significant contribution to our knowledge of how plant foods are translocated in trees. NC(877).

457. The process of water and nutrient movement in the xylem occurs within a preformed but complex structural system. Using radioisotopes to interpretate the translocation of nutrients, recent studies in Wisconsin with cottonwood seedlings have further demonstrated that mature leaves do not import assimilates from other leaves under normal conditions but developing leaves always will. But under special conditions where the mature leaf is above an isotope-fed leaf it will import assimilates and this movement represents a cross-transfer from phloem to xylem at a node. These results demonstrate the importance of knowing the position as well as the structural and the functional relationships of the vascular system if we are to understand nutrient movement in trees. NC(874).

458. The growth of slash pine appears to be limited by shortage of available phosphorus on a number of sites in the South. A study in Louisiana revealed that 1 year-old slash pine seedlings in a growth chamber accumulated more radioactive phosphorus in their needles at a 15-20° C day-night temperature and 2500 foot-candles than at higher temperatures and reduced light. This information will be helpful in selecting the proper season for fertilizer application. SO(878).

459. The presence of mycorrhizae on tree root systems is generally considered beneficial. Investigations by Indian scientists have recently shown that mycorrhizal inocula from hardwood litter were able to form mycorrhizae in conifers and vice versa. But beneficial effects in terms of faster growth occurred only when seedlings received the mycorrhizae from their own type of litter. These results further support the proposal that benefits accrue from mycorrhizae to higher plants only with specific fungi. NE(882).

460. Overwinter depletion of food reserves in tree seedlings may trigger mortality or limit rapid development when growth resumes. Studies in Colorado revealed that lipid, protein, and hemicellulose concentrations in Engelmann spruce seedlings varied little during the period December to June; in contrast carbohydrate concentrations were lower in June, but the differences were no greater than the variability during the winter. These trends disprove the hypothesis that food reserve depletion is a major cause of overwinter mortality of spruce in Colorado plantations, and support the search for other principal causes. RM(883).

461. In view of the potential use of ionizing radiation for producing mutations, we need to know when trees are most sensitive. Two-year-old jack pine seedlings were most sensitive to gamma rays at a dosage of 900R when the exposure was made in July at the time of shoot primordia differentiation. Hence radiation applied during this period could result in the maximum number of mutations from which promising ones might be selected. NC(884).

462. *Pinus merkusii*, a tropical Asian pine which is considered to be a potential plantation species for tropical and subtropical areas, apparently is not suitable for Puerto Rico. Two years after outplanting seedlings of both the insular and continental ecotypes of this species on two sites on the island, the plants were healthy but had made less than one-fourth as much height growth as seedlings of *Pinus caribaea* var. *hondurensis*. ITF(886).

6. Growth and yield

463. Preliminary tables and yield equations indicate that growth of thinned even-aged yellow-poplar stands in the mountains of Virginia, North Carolina, and Geogia may vary from 80 to 280 cubic feet per acre per year depending on site index and residual basal area. The yield tables permit managers to evaluate yields of alternative thinning schedules on sites of varying producitivity in stands of varying ages maintained at different stand densities. SE(889).

464. Wood processors and users of Black Hills ponderosa pine have questioned whether fast-grown wood is apt to have the same technical properties as a slow-grown wood. Stemwood analyses from trees in typical thinned and unthinned second growth stands revealed no differences in average specific gravity, percent late wood, or percent extractives between thinned and unthinned trees; wood at all heights from thinned trees had significantly larger rings and shorter fibers. Height had a significant influence on all variables examined, except ring width in unthinned trees and percent extractives in both the thinned and unthinned trees. Results sug-

gest that manipulation of stocking through thinning to hasten growth of Black Hills ponderosa pine is unlikely to cause major changes in wood properties. RM(891).

465. Intensive culture in northwest Douglas-fir forests is still in its early stages because of the remaining volumes of old-growth timber. An analysis of European data shows that potential yields of Douglas-fir from intensively managed plantations in Europe considerably exceed even gross yields (net yield plus all mortality) of natural stands of the same site index in the U.S. These data aid in predicting responses from future intensive management in the Douglas-fir region. PNW(892).

466. In an experiment in Arkansas, restraint from normal wind sway of 19- to 22-year-old loblolly pines resulted in lessened bole taper, ring width, wood density, compression wood incidence, latewood tracheid diameter, and number of earlywood and latewood tracheids within immobilized portions of the boles. These results help to explain the influence of wind on the growth of trees, and indicate that shortened rotations obtained through heavy thinning may lead to increased reaction wood. SO(893).

467. Although normal yield tables have been in use in this country since about 1900, we still lack tables that allow prediction of stand development as a function of time, site, and cultural treatment. A recently published article points out the need for yield tables for intensively managed stands, and summarizes the current state of the art in yield table construction. The article provides a stepping off point for further yield table work. PNW(894).

468. Ever-increasing demands for more wood as well as nontimber commodities require more intensified timber management practices for the upland oak forests of the East. Growth and yield predictions have been developed for thinned upland oak stands by age, basal area, and site class, all in increments of 10 units from age 20 to 110 years, from 20 to 130 square feet of basal area, and from site index 55 to 85. Application of the recommended thinning practices can increase volume yields from one-fourth to one-third over those obtained from unmanaged stands. NE(897).

469. Slash pine has been extensively planted west of the Mississippi River, where it is not native, but information on its growth and yield in this region is limited. Early results from two long-term thinning experiments in southwest Louisiana indicate that in terms of total volume growth there is little to be gained by thinning on good sites up to 27 years of age. Landowners who are interested in maximum return on short rotations, as for pulpwood, therefore probably do not need to incur the costs associated with thinning. SO(898,902).

470. Much of the potential timber yield in stands of western white pine has been lost because of overstocking. The best response to various degrees of commercial thinning in an 80-year-old stand was obtained by removing about 35 percent of the volume to release the crowns of the best dominant and codominant trees. Results show that this type of thinning can provide an early monetary return from the stand and still maintain volume growth near its maximum rate. INT(899).

471. The ability of unsightly coal mine spoils in Kansas to support rapid tree growth are largely unknown. Remeasurements of plantations established 22 years ago show that some species have made good growth—averaging as much as 35 feet in height and 8.5 inches in diameter in loblolly pine. Other promising species include shortleaf pine, Virginia pine, burr oak, and sycamore. These results show that many spoils in Kansas are sutiable sites for timber production. NC(900).

472. Data on optimum stocking levels for timber production are needed to provide guides for stand density control in ponderosa and Jeffrey pine pole stands. Results of a recent study on site index 65 to 80 land in northeastern California show that thinning to about 80 square feet of basal area per acre will bring such stands to merchantable size as rapidly as possible with negligible loss of total cubic-foot volume production. PSW(904).

473. There is relatively little data on the growth and development of managed stands over long periods of time. Results of repeated thinnings over a 21-year period have recently been summarized to provide this type of information for Douglas-fir in Washington. The thinnings have resulted in improved growth of residual trees with little change in total usable yield, and show that an earlier harvest is the primary benefit of such thinnings. PNW(905).

474. Before short-rotation coppice management for sycamore can be applied on a large scale, the effects of spacing and rotation age need to be determined. In a Georgia study, it was found that the percentage of leaves in the whole tree harvest increased with wider spacing, but decreased with age, while the proportions of branches increased with both wider spacing and increased age. Of the combinations tested, plantings made at 1 by 4 feet and allowed to grow for 4 years produced the largest annual yield (8.9 tons per acre fresh weight annually) with the smallest proportion of that yield in leaves and branches. SE(907).

475. Some foresters have been reluctant to use disking for reducing competition in young stands of pine because of the destruction of roots near the soil surface. In a 12-year-old slash pine plantation on moderately deep sandy soil in northern Florida, disking 2 to 4 times annually reduced the root surface area in the upper 15 cm of soil, doubled the root surface at depths of 30 to 45 cm but did not affect either total root surface area or bolewood production. These comparisons were made with trees in a nondisked part of the area that was cover cropped during the first 5 years but had no competing vegetation afterward. These results indicate that disking can be used to reduce competition in slash pine plantations on similar soils without risk of reducing the root surface area. SE(908).

7. Forest measurements

476. Site indices are often needed on sites where the trees are too young or otherwise unsuitable for direct measurement of the site index. For red pine stands in Minnesota, reliable prediction of height at 50 years of age were made from measurements on young trees of the 5-year height growth in the internodes beginning at 8 feet above the ground. Phosphorus content of the soil also affects site index. Of four methods frequently used for determining phosphorus in agricultural soils, the one involving extraction with dilute (0.01N) hydrochloric acid of the upper 25 cm of mineral soil provided the best estimate of site index for red pine. NC(909,910).

477. Yield tables are available for estimating the volume of pulpwood and saw logs in slash pine plantations, but estimated yields for additional products are needed. Taper curves were constructed for slash pine from which the volume of veneer bolts in a plantation can be estimated. This extension of multiple-product yields is valuable for both buyer and seller when timber sales are made. SE(912).

478. Site quality of newly abandoned fields planted with slash pine under the Soil Conservation Reserve program (soil bank) is better than that on old-field plantations. A new yield table was made that provides a more accurate yield estimate

for 10- to 15-year-old trees on soil bank sites than is provided by yield tables for old-field plantations. Site index curves, however, show that the initial superiority of newly abandoned fields is lost about 20 to 25 years after planting. Hence yield tables for old-field stands are reliable on these sites after 20 years. SE(911).

479. More complete tree utilization, made necessary by critical timber supplies, is creating frequent need for estimates of merchantable stem volume calculated to various merchantable top diameters. Research has shown that transformations can be made to the Behre equation of tree form, making it possible to estimate stem volume to any merchantable limit as a percentage of basal diameter or as a fixed top diameter. The transformations also make it possible to derive taper equations from data gathered on volume to variable top diameters. PSW(913).

480. Land managers in Montana, Idaho, and Utah need growth and yield data on Rocky Mountain lodgepole pine so that they can predict its response to various degrees of thinning. To satisfy this need, an equation was developed for predicting future diameter growth of this variety. Curves plotted from this equation proved to be valid in both managed and unmanaged stands for a time span of one decade. INT(109).

481. An understanding of the theory, development, and application of stocking equations is very useful to land management specialists in making decisions when timber is a part of the resource mix. A concise guide presenting this perspective has been published using newly synthesized equations for determining stocking from point sampling inventories in cutover and virgin ponderosa pine stands in Arizona to illustrate methodology. Stocking equations can be used to help evaluate land treatment potentials, to determine treatment feasibility on a single management unit, and as a basis for setting operating priorities on a number of management units. RM(915).

482. Foresters are increasingly faced with a changeover to metric units—a task that can be extremely complicated in the case of forest inventory systems. Recent modification of STX--a large computerized forest inventory system--allows it to process either U.S. or metric input and to convert from one to the other before final output. This revised system permits immediate use of metric units while retaining the existing capability of handling U.S. units of measure. SE(916).

483. Ring counts of decayed wood are difficult because the wood crumbles away, yet decayed sections of stemwood need to be counted to determine tree age. Accurate ring counts and radial growth measurements on decayed transverse wood sections are possible when rotted wood is frozen. This technique was developed and successfully used during stem analysis studies of old-growth mixed conifers in the Oregon-Washington Cascade Range and has universal utility. PNW(918,919).

484. Presale cruises of standing trees to estimate net volume and value can be quite inaccurate in stands of defective timber. Study in western Oregon of cruising by falling, bucking, and scaling sample trees shows that the magnitude of errors resulting from standing tree measurements is high and that fall, buck, and scale cruising is particularly appropriate for sales containing high-value, defective trees. PSW(920).

485. Volume, taper, and related tables are essential tools that forest managers use in estimating standing timber. Improved tables have been developed for ponderosa pine in Arizona and New Mexico. Volumes are in total cubic feet and cubic feet to a variable top. Tree heights are in feet and numbers of logs. Standards for basic mensurational values meet the needs of both the national Forest Survey and regional forest managers. RM(921).

486. Foresters frequently need quick accurate estimates of stand volumes. Improved tables and point-sampling factors are now available for rapid reliable determination of volumes for Engelmann spruce stands in Colorado and southern Wyoming. Volumes are in total cubic feet, cubic feet to a 4.0-inch top, and board feet (Scribner and International 1/4-inch Rules) to 6-inch and 8-inch tops. Tree heights are in feet and numbers of logs. RM(923).

487. Tree crown width is a commonly used statistic in forest measurement work, and growth and yield studies often require the measurement of thousands of tree crowns at a considerable expenditure of time. Field conditions in the Rocky Mountains prompted development of a small, lightweight mirror instrument of simple design to assist in rapidly and accurately measuring tree crowns. The instrument, which has a large field of view, can be handled easily and leveled horizontally by a user standing in any direction with respect to the tree. It is also rugged enought to survive jostling in a pack or vehicle without damage, since it has no moving parts or delicate mechanism. RM(924).

8. Management planning

488. Organizations faced with the decision of how to best obtain computer services have a bewildering array of alternatives from which to choose. Some helpful guidance for this decision is now available in a recently published article that summarized information on the alternative systems available and discussed the merits of in-house versus outside processing, various methods of access, and the limitations and possible improvement in teleprocessing. SE(928).

489. A comprehensive procedure for computation of yield tables for dwarf mistletoe-infested, even-aged ponderosa pine in Arizona and New Mexico has been developed to help forest managers predict yields and evaluate and make necessary comparisons of alternative management decisions for these stands. RM(929).

9. Maple syrup production

490. Sugarbush operators are interested in the relationship between sap-sugar yields and characteristics of sugar maple trees. In general, individual tree characteristics that express live crown ratio were highly related to sap volume yield and total sugar production. There was little evidence of a relationship between tree characteristics and sap-sugar concentration. NE(934).

491. The lack of an easy method of estimating relative sap volume produced between trees has long stymied efforts to make phenotypic selections of sugar maples for superior sap volume production. From a study in Vermont, a high correlation between sap-flow rate and total sap volume produced during the sugaring season was found among trees that were both single- and multi-tapped. Measurements of sap-flow rates over a short period of time in conjunction with other parameters provide a reliable measure for making plus-tree selections for super sap volumes. NE(935).

492. In northern Vermont, vacuum pumping for sap collection during the fall is not advisable even though profitable sap collection by gravity from fall tapping of sugar maple trees has been reported in certain localities. The volume yields were only 2/3 as much as in the spring. It is recommended that if favorable conditions are observed in a locale, vacuum pumping in the fall should be tried on a small scale before attempting a large operation. NE(939).

493. With the renewed interest in plastic tubing for collecting maple sap for processing into maple syrup, a how-to-do-it guide for constructing and installing the plastic tubing system has been prepared. Both aerial and ground lines are described, with details on hanging lateral lines, laying ground lines, tapping trees, and connecting spouts, droplines, and conduit lines. NE(940).

494. Bigleaf maple in California, Oregon, and Washington produces a sweet sap similar to that of eastern species of maple, but little is known about its potential for the production of syrup. In experiments in Oregon, the volumes and sugar percentages of sap from bigleaf maple were somewhat less than from sugar maple sap in the East. Color and flavor of the syrup produced were about equal to the lower grades of sugar maple syrup. Production of syrup from bigleaf maple is feasible and with the application of improved extraction, collection, and processing methods, commercial production may be possible. PNW(938).

495. The size of tubing needed for maple sap collection in a sugarbush is an important decision because of cost and its effect on sap yield. From studies in Vermont the most important factors affecting maple sap flow in plastic tubing are tubing diameter and slope percent. Estimates are given to help determine the most efficient tubing installation for a specific sugarbush situation. NE(941).

496. In general, fall tapping without vacuum of sugar maple trees for sap collection is not recommended. The sugar content of the fall collected sap is about 2/3 of the sugar content found in the spring and sap volume yields are very low, only 15 percent of the spring volume. However, it is suggested that fall sugar readings can be used to determine a tree's relative sweetness and thus serve as a guide for tree selection for sugarbush thinning. NE(942).

10. Naval stores

497. Very little is known about the factors affecting resin duct formation in pines and the mechanism of resin secretion. Studies on Aleppo pine in Israel showed that resin ducts are formed about 3 1/2 months after cambial activity is resumed and their number is determined by the temperature at the time the cambium becomes active. A resin droplet being forced through a cell wall was photographed in a resin duct fragment kept alive in an isotomic solution of mannitol. This research has advanced basic knowledge of the biology of pines. WO(946,947).

B. Breeding Improved Trees

1. Distribution and classification of forest trees

498. Data on the distribution of various trees species is usually presented in rather general terms. More specific information has recently been published for 86 forest and woodland tree taxa native to California. Maps are accompanied by descriptive notes that relate the distribution of each species to its ecological and taxonomic position. This publication provides basic information needed by foresters, ecologists, and others who work with the tree and woodland communities of California. PSW(948).

499. The true firs, one of the largest groups of coniferous forest trees in the world, are widely distributed over the Northern Hemisphere especially in the high altitudes and boreal regions. Because of their great morphological variability, the taxonomy of true firs has long been in a state of confusion. To clarify the botanical relationships among the world firs, a comprehensive investigation was conducted by members of the staff of the National Taiwan University in cooperation with the U.S. Forest Service. A fully illustrated monograph was prepared which provides detailed descriptions, keys, lists, indexes of common names, notes on habitat, and a general bibliography for 39 species, 23 varieties, and 8 hybrids, with special chapters devoted to the fossil history of the group and the geographical and altitudinal distribution. Tree scientists in the United States as well as those the world over will find this book of immediate use in the identification of the true firs, in genetic studies, and in selecting the proper species to be planted. WO(950).

500. There are six species of white pines native to western North America and Central America. These include the commercially important species of sugar pine, western white pine, Mexican white pine, and southwestern white pine. The two remaining white pine species, whitebark pine and limber pine, are not of commercial importance but have a high esthetic value. The distribution, variation, and site requirements for these six species are now summarized and evaluated for those interested in these species. INT(951).

2. Inherent variation

501. Black walnut shows great genetic diversity across its wide natural range and it is sometimes difficult to decide which seed source one should plant. Recent studies in Illinois have indicated that southern seed sources flush earlier in the spring and drop their leaves later in the fall than northern sources. In addition, southern sources also grew faster than northern sources. These early results suggest that where spring frosts are not a serious problem southern walnut seed sources should be planted. NC(953).

502. Late spring frosts will damage the tender new growth of black walnut trees and the earliest trees to break dormancy are vulnerable for the longest period. In an effort to avoid or to lessen the problems associated with frosts, the pattern of leaf flushing for walnut from various midwest geographic locations was determined. Southern seed sources were found to flush earlier than northern sources and trees that flushed early one year also flushed early the next year. In selecting the proper walnut seed source early flushing trees should be avoided in an area with a history of late spring frosts. NC(954).

503. The long generation interval in forest trees is often a hinderance to early selection and breeding programs. Thus, early flowering is a highly desirable trait but early flowering should not reduce vegetative growth. In a recent study 6-year-old Virginia pines produced sufficient numbers of both male and female strobili to allow early selections and rapid genetic gain, and the reproductive growth did not reduce height growth. SE(955).

504. Because trees with rapid juvenile height growth pass through the vulnerable seedling stage more quickly, nursery and outplanting practices which increase seedling growth are being developed but these methods are expensive. Genetic improvement of juvenile growth has been suggested as a possible alternative. A recent study with 54 Douglas-fir families planted at two sites in Washington showed that potential for genetic improvement in rapid early seedling height growth is good with an average genetic gain in seedling height growth of approximately 12 percent. PNW(956).

505. Polyembryony, more than one embryo per seed, is a rare occurrence particularly in angiosperms. Recently this unusual phenomenon has been found in sugar maple and two sets of twin seedlings were grown successfully for more than

3 years. This is the only known case of sugar maple twins actually developing into normal seedlings. Later studies will determine their genetic make up. NE(957).

506. The use of oleoresin analyses in identifying pine species and hybrids is now a well established method in many phases of genetics and tree improvement, especially those dealing with the development of pest-resistance for certain tree species. There are, however, limits to the usefulness of this technique. For example, recent oleoresin analyses throughout the range of loblolly and shortleaf pines demonstrate insignificant qualitative differences between comparable samples and no differences between samples in or out of southern pine bark beetle epidemic areas. There was, however, a decreasing percentage of the dominant terpene, a-pinene, in samples from east to west and from south to north throughout their natural range. SO(959).

507. The ability of a coniferous seedling to develop into a vigorously growing tree depends largely upon the size and rate of activity of its shoot apical meristem. By means of both anatomical and mathematical techniques the apical dome volume was found to increase 10-fold while meristematic volume growth rate declined from 20% to 12.5% per day in 20-days-old to 140-days-old spruce seedlings. Further analysis of this basic type of growth information will yield a better understanding of the problems and processes associated with the differences between slowly growing trees and genetically superior trees. NE(963, 964).

508. Wavy grain pattern is a desirable feature in selected high-quality hardwoods. It may be possible in the future to encourage the wavy grain development with a better understanding on how such patterns originated. Recent studies in Poland supported by the PL-480 program have now established how cell divisions in the cambium are preferentially oriented to the right or left and how they form a wave pattern. This is an important first step in establishing how the wavy patterns are initially developed. WO(965).

509. In order to obtain better yield estimates of genetically superior seed from young Jack pine seed orchards, cone production, cone characteristics, and seed yield per cone from different seed sources planted in Northern Wisconsin were studied. For seven of the seed sources nearly all cone characteristics and seed yields were similar, with two of the more western seed sources having the smallest seed. There was, however, considerable variation among trees within a given seed source. These results suggest that as much consideration and importance must be placed on variation between trees of the same source as has been previously placed on trees of different seed sources. NC(966).

510. A common problem in forest tree improvement research is the extrapolation of data from young progeny or provenance test plantations to predict tree performance at final rotation age. In a loblolly pine progeny test plantation measured at ages 3, 4, and 15 years, genetic correlations showed that little change in genetic control of height takes place after the 5th year, and that resistance to fusiform rust at age 15 differs little from that at ages 3 or 5. These high genetic correlations provide good evidence that families can be selected at early ages for breeding purposes. SE(968).

511. It is commonly assumed that improved trees from forest genetics programs can be evaluated only after many years. But a test of eastern cottonwood clones in Mississippi has shown that correlations between measurements made in the first three years and in the sixth year were high, suggesting that culling of clones after two growing seasons is a feasible way to reduce the time and cost of cottonwood improvement programs. The study also provides estimates of heritabilities and indicates that large increases in volume production can be obtained by selection for either height or diameter. SO(969).

512. Differences in the growth of forest trees of different seed sources are caused by unequal sensitivity to external factors, such as temperature and light. Recent studies in Poland have shown that trees of different origin also differ in their sensitivity to X-rays. Seedlings of high-elevation Scots pine of Polish origin grown from X-ray-irradiated seed were more radioresistant and recovered better from radiation effects than seedlings from low land. Low dose X-irradiation of seed was found also to stimulate dry matter production of and increased size as well as number of needles on Scots pine seedlings. WO(970,979).

513. A recent genetics study of several 53-year-old plantations of selected Douglas-fir in western Washington and northwestern Oregon indicate that genetic control of apical growth is strong in certain environments and that this control changes during a tree's lifetime. Genetic control is the strongest in the juvenile and early reproductive years but may decline under some conditions of environmental pressures. These results, support the feasibility of genetic selection for early rapid growth in Douglas-fir. PNW(971).

514. White spruce, one of the most widely distributed conifers in North America, has a considerable growth potential and is now used widely to produce a high-grade pulp for the production of quality paper. To assist the forest geneticists and tree breeders in planning future white spruce improvement programs, a review of its genetics now has been prepared. In this review white spruce evolution, migration history, distribution, phenology, heritability, and breeding behavior are summarized. NC(972).

515. Selected trees must be progeny tested but the tests are expensive and require many years to complete if decisions are delayed until the progenies are at or near rotation age. Data from 15-year-old progenies of 27 selected slash pine trees not only revealed differences among the progenies at 15 years but also showed that third-year data on height, survival, and rust infection could be used to accurately predict volume per acre at age 8 or at age 15. Data on these three traits could be used to choose among families much earlier than at rotation age. SE(974).

516. Basic information on variation of wood properties and other traits of yellow-poplar is needed. Analysis of wood samples from 500 yellow-poplar trees in the Southern Appalachians revealed that specific gravity in the species decreased with increasing site index, increasing elevation, and more northerly latitudes. Only part of the variation in specific gravity could be attributed to enviromental factors. This reliable description of the population is useful in designing future tree improvement programs for yellow-poplar in the Southern Appalachians. SE(975).

517. Spanish cedar (*Cedrela* spp.) produces a valuable tropical wood but is difficult to regenerate. The genus has a wide distribution in the Western Hemisphere which appears to be associated with considerable natural variation. A provenance trial in Puerto Rico and St. Croix has already revealed difference in survival, growth, and susceptibility to shoot borer attack in the nursery, with best and poorest overall performance by sources from Guanacaste, Costa Rica, and Misiones, Argentina, respectively. The results of field trials with these seedlings will help to identify superior provenances for use in Puerto Rico and similar areas in the American tropics. ITF(976).

518. Monoterpene compounds in the oleoresin of conifers are useful in defining genetic differentiation of populations of many tree species. None of the 9 measured monoterpenes differed significantly among geographic seed sources of red spruce in 14 populations growing in Quebec, Canada. Although the monoterpenes were not suitable for determining hybridization between red and black spruce, they may still be of value in studying hybridization of other spruce species. NE(977).

519. Trees of the same species but from different geographic areas often differ in both growth characteristics and selected wood properties. This general observation has been further supported by a recent Forest Service funded study at the University of Texas where sweetgum seedlings of different origins were grown under a series of controlled temperature and light conditions. Greater fiber tracheid length and lower specific gravity was correlated with a decrease in latitude of seed origin, while cell length varied more with environmental conditions of short day-length and cooler temperatures than seed origin. When cell length and wood density are important in sweetgum utilization both the origin and the environmental conditions under which the trees are growing should be considered. WO(978).

3. Insect-disease resistance

520. Western white pine is among the most valuable conifers native to western United States. However, because the blister rust disease limits its growth and survival, western white pine management is uneconomical over many parts of its native range. More than 20 years have been devoted to studying the inheritance of white pine blister rust disease resistance and now this information as well as features of the paleobotany, flowering, seed yield, and inbreeding of this species, have been summarized. This summary is of value to those who are interested in improving and selecting disease resistant western white pine for future planting. INT(982).

521. Blister rust disease is a major limiting factor in the successful growing of white pines in many parts of the world. Among the more than 20 species of white pines, there are possible differences in their resistance to blister rust. A re-examination of the world's literature on blister rust resistance and on field observations provided information for a tentative ranking among 14 of the different white pine species as to their disease resistance. This new ranking along with consolidated and updated information on botanical ranges as well as crossability of the white pines can be used in planning future white pine blister rust resistance breeding. INT(981).

522. Brown-spot needle blight is a major deterrent to natural regeneration of longleaf pine in the South. Recent research in Alabama shows that about 10 percent of wild seedlings remain nearly free of the disease, and most of these have superior juvenile growth. Natural regeneration following a good longleaf pine seed year often provides seedlings in such numbers that the disease-resistant fraction alone will provide a full stand. However, the land manager must recognize and exploit their existence, especially by modifying prescribed burning practices so that the superior seedlings are not injured. SO(983).

523. Methods are needed for producing varieties of tree species resistant to phytotoxic gases in the atmosphere. A selection system based on visible needle injury has been developed and tested for evaluating resistance of foliage of Scotch pine to injury by ozone and sulfur dioxide. The system is useful in evaluating breeding programs or in mass production of resistant clones or varieties, and has been shown to be successful in selecting resistant Scots pine. NE(984).

524. Improved methods are needed for the rapid identification of resistance in southern pines to the fusiform rust fungus. In recent investigations in Mississippi two artificial inoculation tests made it possible to identify resistance trees in 9 months, compared to at least 4 years required in field tests. The proportion of plants infected was a better measure of field performance than numbers of stem infections or total numbers of infections. This new artificial inoculation method has wide application in the fusiform rust resistance program now being developed in the South. SO(985).

525. The most serious pest on white pine in the Northeast is the white-pine weevil. Leader damage by the pine weevil on 27 seed sources tested in southern Maine suggests that chances of locating individual sources with acceptable levels of resistance are remote. Alternative methods of producing weevil resistant material must be explored. NE(986,987).

526. Tree rusts are the most serious of all forest diseases, and a knowledge of host-fungus resistance physiology is a prerequisite for efficient genetic gains. A recent review of the world's literature on this subject suggests that the principal resistance mechanisms against the major rust, white pine blister rust, are hypersensitivity of foliage and stem cells, and production of periderm barriers in the stem. The main rust of the southern pines, fusiform rust, appears to be more passive since resistance to it involves preformed fungus toxins. SO(988).

527. Among the most destructive pests of forest trees are members of the rust genera *Cronartium* and *Peridermium*. A recent review for forest tree breeders of the world's literature indicates that tree susceptibility to the rusts ranges from high to near immunity, with interspecies hybrids exhibiting a similar range. Heritability of resistance can be high and relatively stable to varying conditions of rust hazard. Although little is known of mechanisms of resistance, this review indicates a physiological basis, depending on the rate of response of the tree's cortical tissues to invasion by the rust. PSW(989).

528. Slow early height growth of longleaf pine is often partly due to the stunting effects of the brown-spot disease. In a test with wind-pollinated seedlings from 540 parents, infection in the seedlings of the best 10 percent of the parents average 48 percent compared to a population average of 63 percent. Inherent fast height growth was not a major resistance mechanism. In addition, the percentage of resistant, fast-growing seedlings identified varied greatly with seed source and was greatest in seedlings from southwest Alabama seed. Seeds and cuttings from the selected parents are now being used to establish a resistance breeding seed orchard. SO(990).

4. Tree breeding methodology

529. Black walnut seed orchards can provide nurserymen with easily accessible genetically improved seed of known origin. Guidelines and some useful hints have now been developed to assist those interested in establishing a walnut seed orchard. In developing a seed orchard consideration must be given to defining the seed collection area, as well as to the actual growing, planting, and maintaining of the superior seedlings. Planning for the next generation orchard is also explained. NC(991).

530. Self-pollination can be a major cause of seed loss in pine seed orcahrds. Using available data on seed development a theoretical model was developed to describe the relationship among multiple fertilizations, embryonic lethal alleles, and unsound seed following self-pollination. Tables from this model can now be used to predict the occurrence of unsound seed in

self-pollinated pines or to estimate the number of lethal alleles carried by the self-pollinated parent. SE(992).

531. In an attempt to induce genetic variation in river birch, its pollen was first irradiated by various levels of gamma radiation and then used in a series of controlled crosses. Pollen irradiation had no effect on the number of female flowers that matured nor on the number of seeds per flower. But viability of the filled seeds was reduced to 60 percent and young seedlings having an abnormal number or shape of cotyledons increased with an exposure of 8000 roetgens or greater. This increase in aberrant seedlings shows that the technique may be an effective means of inducing genetic variation, in spite of the reduced reproductive capacity resulting from use of the irradiated pollen. NC(993).

532. A 1970 evaluation to determine the scope of the artificial regeneration efforts in California by the Forest Tree Improvement Committee, Northern California Section, Society of American Foresters, indicated that approximately 34,000 acres were planted and 20,000 acres direct seeded. Fifty-four percent of the acres regenerated was with Douglas-fir and only a very small percentage of these acres were planted with genetically improved seeds. The appraisal indicates that the current tree improvement effort in the State of California falls far short of that required to improve this State's timber resources, and recommends an expanded program of silvicultural and genetic research to meet the future timber needs of California. PSW(994).

533. The identification of selected enzymes from various tissues of forest trees can serve as excellent genetic markers in tree breeding studies. Equipment and procedures employing starch gel electrophoresis for the identification of individual enzymes are fully described and illustrated for use by other scientists intersted in using this new technique. PSW(995).

534. The failure of the graft union to heal normally is a serious limitation to the development of the grafted Douglas-fir seed orchards. In a recent investigation in the Pacific Northwest where the graft compatibility of genetically related and unrelated rootstock scion combinations were compared, scion clones were 75 percent compatible when grafted on half-related rootstocks. In contrast scion clones were only 56 percent compatible when grafted on unrelated rootstocks. Grafting failures can be reduced by controlling the genetic origin of the rootstocks in future seed orchards. PNW(996).

535. Progeny testing to evaluate the genetic superiority of selected trees has the disadvantage of requiring a long period of time. In slash pine, good correlation between parents and their 3-year-old offspring was obtained for growth rate, specific gravity of wood, turpentine content of wood, and ethanol-benzine extractives. Hence, the progeny test period for these traits in slash pine may be reduced from 20 or more years to 3 years. SE(997).

536. Willow and cottonwood trees are most often propagated by rooting cuttings. Recent studies in India have demonstrated that rooting of willow stem segments is enhanced by 0.1 mg/l. IAA and 1.0% glucose while black cottonwood cuttings rooted best with 0.1 mg/l. IBA and 0.5% sucrose. Hormone effects on rooting of these species are determined partly by the nutritional status of the cutting, and the addition of a carbohydrate source such as glucose or sucrose provides a correct physiological balance. WO(999,1011).

537. To date only a few dawn redwood trees outside of China have produced both male and female cones. This had led to some misinformation concerning the origin and distribution of male cones on this rare and interesting tree. Recent observations in California have now clearly shown that the male cones are produced in lead axils on highly specialized flower branches and are not in florescences as previously reported. PSW(1000).

538. Grafting branches from selected superior walnut trees is a useful way of reproducing this species but often after a few years the grafted tree dies. In order to determine the cause or causes for this failure as well as the internal anatomical changes associated with the graft union, 22 combinations of walnut species and hybrids were grafted in Illinois. During the first 5 years, the bark grew faster on most of the black walnut stock plants than on the grafted branch of the Persian x black walnut hybrid. This is an early indication of incompatibility which must be avoided. NC(1001).

539. To better understand the fruiting process in forest trees and to aid in developing methods for reducing flower abortion, a Polish scientist has conducted a series of detailed investigations on the qualitative and quantitative levels of hormones in the reproductive organs of poplar. Immature flowers were generally high in auxins and low in inhibitors. The maximum amounts of inhibitors are found in the bursting buds, not in dormant ones as previously thought. Female flowers in the catkin stage contained more auxins that the male. WO(1002).

540. Tissue culture, the practice of growing a given plant tissue under controlled conditions, could provide a quicker and a better method of vegetatively reproducing desirable forest trees. An Indian scientist has now been able to establish tissue callus from five American pine species and two Indian species on known nutritive media. For chilgoza pine native to India he has obtained limited differentiation of root and presumptive shoot apical meristems. Although whole trees were not developed, these results do suggest that successful propagation of coniferous trees by tissue culture techniques will be possible in the near future. WO(1003).

541. Little is known about the financial returns that can be expected from hardwood tree improvement. Results of a preliminary economic analysis suggest that tree improvement with artificial regeneration is likely to be profitable for those hardwood species such as black cherry and paper birch that grow rapidly, produce large quantities of seed, and have higher than average market values. A secondary finding was that tree imporvement would pay only when combined with a program of intensive culture. These results emphasize the need for care in selection of hardwood species for tree improvement programs, and provide some guides for the selection of species on which tree improvement is likely to be profitable. NE(1006).

542. Rooted cuttings of yellow-poplar are most easily obtained from sprouts, but propagation of select trees by this method is of limited usefulness unless some means is devised to induce sprouting without cutting the entire tree. Studies in Georgia have shown that sprouting can be induced on most trees by partial girdling, or by severe pruning of various portions of the crown. These methods of forced sprouting and propagation offer a shortcut means of obtaining clonal lines for research or seed orchard uses. SE(1004,1005).

543. To further our understanding of the rooting phenomena of hardwoods the endogenous hormones of developing buds, shoots, and adventitious roots of willows were isolated and identified by a team of Polish scientists. A gibberellinlike hormone and growth inhibitor similar to abscisic acid were found in both the shoot and bud of willow cuttings. It appears that the root influences both the quantity and quality of the gibberellin in the shoot and vice versa, but the controlling factor of the inhibitor was not demonstrated.

547-679 O - 74 - 4

Later studies will deal with the actual interactions of the inhibitors and gibberellins on adventitious rooting. WO(1007,1008).

544. In the study of the dynamics associated with changes in forest tree populations, there is a recurring need to define these changes in terms of mathematical equations in order to estimate future changes. Applying existing Leslie-type matrix analyses, projections of forest tree population behavior were constructed which account for variation in both survival and reproduction due to possible sampling error as well as to variations in the environment. This model equation must still be tested as additional data becomes available. SE(1009).

545. Detailed investigations in India of the comparative rooting ability of a number of different tree species indicated that, in general, ease of rooting is associated with certain groups of trees. Furthermore, their results strongly support the widely held view that seasonal variation in the rooting ability of cuttings is related to the state of the dormancy which, in turn, governs starch and auxin levels in cuttings. Hormone induced rooting could be accelerated and increased by the addition of an energy source, such as sucrose. WO(1010).

546. One of the most important factors in determining the profitability of seedling seed orchards is amount of time required for the orchard to produce seed. Both fertilization and cultivation were found to enhance early male and female flower production in a 4- to 7-year-old loblolly seed orchard in southern Mississippi. Although increasing the level of fertilization had little effect on increasing the number of trees flowering, it did increase the number of cones on those trees which were bearing. SO(1013).

547. Before improved plant material is released for commercial use it is necessary to describe the methods and criteria used to select and reject clones, varieties, or parentages. Beginning with 13,000 hybrid poplar seedlings in 1924, 40 clones have now been selected in the Northeast on the basis of growth rate and pest resistance. These clones have been distributed to ten state nurseries and are now available for commercial plantings. NE(1014).

548. Forest tree improvement programs are now an established part of the total management of our public and industrial forests. In order to assist the natural resource manager with the terms and language employed by tree breeders and to provide a common and acceptable language for the tree breeders in different parts of the country, a revised glossary of 175 terms in genetics has been prepared. This new edition updates the 1959 edition by redefining most terms, adding some, and deleting others. SO(1015).

549. Among oak species are a number of valuable trees which are becoming more important in forest management with the increasing demand for hardwood fiber. Improved methods for the genetic improvement of oaks such as long-term pollen storage at 18°C and better methods of pollination using special bags were developed and tested in Yugoslavia under the PL-480 program. These investigations also described the basic details of flower development which will directly aid the U.S. tree breeder in applying these new methods to American species. WO(1020).

550. Walnut is a valuable forest tree in the United States, producing both a useful nut and highly valued wood for the furniture industry. PL-480 supported research in Yugoslavia has shown that superior Persian walnuts that grew faster and had fewer forks came from the higher altitudes or more northerly latitudes of Yugoslavia. By crossing the superior Persian and American black walnut additional genetic improvement can be achieved. WO(1021).

551. Seedlings of western white pine have a definite period of winter dormancy. For this species to be grown normally in a greenhouse or growth chamber for experimental purposes, a cold-conditioning method was developed. For 2-year-old seedlings to approach maximal growth without natural winter conditioning, they must receive a minimum of 14 weeks of artificial chilling at 40°F. INT(1019).

552. Polyploid races of forest trees have potential value as an intermediate stage in forest tree breeding research since such trees provide a mechanisms for controlled gene exchange. In a series of studies in Finland, germinating seeds of 10 birch species were treated with 0.06% concentration of the chemical colchicine. Colchicine induced polyploids in six of the birch species, but the polyploid seedlings were characterized by a high mortality and an initial slow growth. At age 3 to 5 trees had short trunks and bushlike branching with leaves greater than normal size, larger stomata, and larger mesh size of the venation network on the lower surface. Later studies will determine how these trees flower, and whether they will produce seeds following controlled pollination. WO(1022).

553. When different pine species are crossed, cone abortion is often high and seed-set low. Two Yugoslavian scientists have attempted to overcome these breeding barriers in the cross of Austrian and Scots pine by using pollen previously irradiated with gamma rays. Irradiated pollen permitted normal seed development for 11.5 months following pollination, while cones in the same cross treated with normal pollen aborted shortly after pollination. Although they did not get seed-set, their method will make it possible to overcome selected breeding barriers in some American pine species by preventing early cone abortion. WO(1023).

C. Improving Uses and Protection of Wood

1. Utilization potential and processing of wood

554. A reference handbook for those interested in, involved in, or contemplating utilization of the major woodland tree species of Arizona provides information on stand and stock characteristics and physical and chemical properties of the species. The occurrence, physical characteristics, and utilization potential of pinyon, juniper, and Gambel oak are reported here. RM(1220).

555. *Pinus caribaea* grown in Puerto Rican plantations were sampled to determine the amount, orientation, and effect of compression on some physical and mechanical properties of the wood. Compression wood did not affect specific gravity, modulus of rupture, or work to maximum load in beam tests. Modulus of elasticity decreased as did all shrinkages except longitudinal. Boards from thse trees were kiln dried without seasoning degrade and were easy to machine. The results indicate that plantation grown *Pinus caribaea*, although high in compression wood, can be utilized successfully. ITF(1221).

556. Newly developed standard eastern white pine logs and tree grades were tested for reliability in predicting the value of standard yard lumber products from a sample of white pine trees and logs. The tree grading system predicted a lumber value of the sample trees within 3 percent of the actual, and the log grading system predicted a lumber value of sample logs within 2 percent of the actual. The grades provide a working tool for both buyers and sellers of white pine logs. NE(1222).

557. Weight scaling, rather than log scaling, is the most common method of purchasing southern pine tree-length logs. Season of the year affects the weight loss due to drying dur-

ing storage. Woodyards in South Georgia should limit storage periods for pine tree-length logs to approximately one week in summer, but could extend storage up to 6 weeks during the winter months. SE(1225).

558. Heartwood and sapwood of black walnut (*Juglans nigra* L.) were prefrozen and then thawed prior to drying to test the effect of prefreezing on wood hygroscopicity, dimensional properties, and extractives. The tests indicated that prefreezing induced shrinkage reductions result from increased bulking of the fibers by extractives during drying and altered hygroscopicity of the wood. What is meaningful is that prefreezing favorably altered the wood-water system and therefore can provide the possibility of accelerating drying without increasing defects. NC(1227).

559. A computer program, written in FORTRAN, predicts the maximum yield of cuttings for softwood cut-up and edge- and end-gluing operations. The program calculates cutting recovery (given cutting width and length constraints and defect locations on the board) and specifies the maximum number of rips. This program will improve wood conversion efficiency through improving yields, reducing wastes, and reducing human errors. RM(1231).

560. Trees cut in thinning of Douglas-fir stands were processed into lumber by a profiled cant chipper, a square cant chipper with resaw, and a bandmill. Recovery totals by processing method are given in both lumber and chips. This information will be useful to mill operators and timber managers in predicting product yields and evaluating different production methods. PNW(1232).

561. Residual volumes of wood, bark, and needles were obtained in a lodgepole pine stand after clearcutting to a 6-inch-diameter top. A total of 4,333.1 cubic feet of residue per acre, excluding needles, remained after logging. In terms of utilization potential, 82 percent of the residue was 3 inches or larger. FPL(1233).

562. An adhesives evaluation system has been devised providing data for specific design situations that can be related to anticipated levels of stress and deflection, as well as the critical nature of each adhesive application. The system includes accelerated aging to forecast durability properties for 50 years and longer. This approach provides information needed by adhesive formulators, raw material manufacturers, and architects and design engineers to secure dependable adhesive bonding in building construction. FPL(1235).

563. A study of surface characteristics and lumber recovery information indicates that problems encountered in applying a log grading system to sugar pine trees can be alleviated by modifications in the specifications. The modifications do not appear to decrease the performance of the grading system used by foresters and cruisers in various operations requiring grading. PNW(1243).

564. Thirty-six resins formulated with varying ratios of phenol, sodium hydroxide, formaldehyde, and resin solids were used in tests of glue bond quality of southern pine. Of the three formulation variables, the NaOH/phenol ratio gave the most surprising results. The ratio of 0.4 is considerably lower than any used in industry, but resins thus catalyzed gave consistently better bonds than those with higher proportions of caustic. SO(1245).

565. Recently, interest has developed in applying the scanning electron microscope to study the stretching of cellulose fibers. A single and inexpensive fiber-stretching device was designed and fabricated using the existing standard stereostage and goniometer 360 degrees stage control. Thus, cellulose fibers can be stretched and photographed while being viewed with a scanning electron microscope. FPL(1253).

566. A two-volume handbook characterizes the 10 southern pine species as an industrial raw material and describes the processes by which it is converted to use. This handbook is the most comprehensive reference to the utilization of the southern pines available anywhere. SO(1254).

567. A public U.S. patent was issued for a process for steam straightening and kiln drying lumber, the lumber being held under total restraint with respect to warping during the process. The lumber is steam straightened and kiln dried to a moisture content of 9 percent. SO(1257).

568. About 98 percent of the Sitka spruce lumber sawn in a southeastern Alaska lumber recovery study was in cants 3 to 8 inches in thickness. Thirteen percent of the lumber volume was graded No. 2 and 3 clear by the Pacific Lumber Inspection Bureau rules and 9 percent high grade cants (Piano grade) under export rules. PNW(1259).

569. New log grades for cruising coast Douglas-fir have been developed and tested. They are for application to standing timber and provide an accurate estimate of lumber and veneer yields and timber values. PNW(1260).

570. Rate-related dimensional stability characteristics of wood-base composition board panels were found to be dependent on interparticle capillarity. Total moisture absorption was controlled by characteristics inherent to the varying types of composition board and was found to be a logarithm of time. Linear expansion and thickness swelling were found to be directly correlated with moisture absorption and, in some test, time. Methods which improved bonding or reduced the number of available absorption sites were proved to be better stabilizing methods. This research was awarded first place in the 1970 Wood Award Competition co-sponsored by the Forest Products Research Society and Wood and Wood Products Magazine. FPL(1262).

571. Gas chromatography has been found an efficient and relatively accurate method for quantitatively and qualitatively determining the major components of creosote. Results of this procedure compare favorably with the more tedious flask distillation procedures, and separation of individual components can be achieved in as short a time as half an hour. FPL(1263).

572. A prediction equation has been developed for estimating the initial moisture content of Douglas-fir based on green volume and green weight. The same two variables and elapsed drying time can also be used for efficiently estimating the percent of moisture content during drying. The same estimates can be made for ponderosa pine when hardness of green wood is considered as an additional variable. Commercial application of this predictive method is directed toward kiln-drying operations. INT(1264).

573. Properties that affect the manufacture and use of veneer and products made from veneer are described for 156 U.S. tree species. Each species is rated for use in the general categories of construction plywood, decorative face veneer, inner plies of decorative panels, or container veneer and plywood. This report serves as ready reference for all U.S. species likely to be used for veneer. FPL(1267).

574. Wood moisture content affects tool forces and thus the energy consumed in machining. Moisture also interacts with other factors in influence the process of chip formation and, hence, affects the shape and size of the separated particle. When boring holes in southern pine of varying moisture content, torque and thrust increased with increasing moisture content to a maximum of 5 to 10 percent, then decreased to a constant value at fiber-saturation point. SO(1270).

575. The Western Wood Density Survey is a long-term program to assess the wood quality via wood density for the major western timber species. Wood density data are presented for 15 commercially important western timber species, with emphasis on 6 species not previously evaluated. FPL(1271).

576. Maple logs were sawed by two grade-sawing methods and a live-sawing method into lumber. Live sawing gave the greatest yield of rough clear cuttings; however, when warp is considered, grade-sawing becomes superior. Where a remanufacturing plant has any control over the sawmilling operation, loss from warp can be reduced if the sawyer is instructed to avoid live-sawing as much as possible. FPL(1272).

577. Rootwood tracheids of southern pine average one-third longer and are one-third larger in diameter and have walls 18 percent thinner and lumens almost two-thirds larger than stemwood tracheids at stump height. Since the stump and roots left after logging contain about 25 percent as much wood as is found in the harvested stem, knowledge of root tracheid measurements is important in assessing the utilization potential. SO(1273).

578. Trees throughout the commercial range of spruce pine (*Pinus glabra* Walt.) were sampled to determine effects of tree age and growth rate on morphological, mechanical, physical, and chemical properties. The wood properties studied were those deemed of importance in the manufacture of fiber or solid products. SO(1275).

579. The feasibility of roll-laminating fiber overlays on low-quality ponderosa pine common to the Southwest U.S. was studied at the pilot plant stage. Lumber repair techniques involving abrasive planing, wood plugging, chemical repair, and selective cutup were also studied. Results indicated that a high proportion of defects cannot be successfully overlaid without repair and that repair techniques are restricted by the need for an effective automated defect repair system. RM(1279).

580. The relationships of wood specific gravity to height in tree were investigated for 28 commercially important timber species. In 17 species, specific gravity decreased with increase in height; in 5 species, specific gravity increased with increase in height; in 3 species, specific gravity initially decreased with increase in height, then increased; and in 3 species, no significant change was observed. Information bears on strength properties and pulp yields of upper logs of sticks. FPL(1282).

581. One-half-inch-thick slices of clear, flat-grained red oak and southern pine were cut with the knife and flitch inclined, relative to the cutting motion, at angles of 45, 67.5, and 90 degrees. The depth of fractures into the wood slices was least when using inclination-angle differences of 45 degrees; however, thickness uniformity was poorer, the surfaces rougher, and there was considerable twist. Best overall quality occurred when there was no angular difference between the knife edge and the grain direction of the wood. FPL(1283).

582. Slash pine has thicker bark, a higher percent of bark by weight, a lower bark moisture content, and a higher bark specific gravity than loblolly pine. Location differences within a species were large for bark moisture content and bark specific gravity. This research is directed toward the development of increasing the utilization of the bark residue resource. SE(1284).

583. A newly revised guide to hardwood log grading was prepared. The guide is to serve as a teaching aid and field reference. Basic principles and details on practical applications are given. NE(1286).

584. An apparatus to continuously and simultaneously measure the swelling and liquid uptake by cylinders of wood is described. The device measures swelling during liquid penetration into wood, and can also be used to study the effect of various chemical treatments on either swelling or shrinkage. Measurements of this kind are useful for determining the best chemical treatment conditions. NC(1288).

585. Production of a laminated structural wood product by an innovative system which markedly improves yield at reasonable cost is now technically feasible. The complete log-to-product system consists of rotary cutting veneer up to one-half inch thick, press-drying the thick veneer in less than 15 minutes, applying glue to the hot sheets, and laminating into a thick structural material. The entire processing time is 30 minutes. An economic analysis shows that the selling price for vertically laminated joists would be equivalent to the majority of sawn structural lumber used for housing. FPL(1291).

586. Nomographs have been prepared allowing accurate estimates of dimension yields from various grades of hard maple, black walnut, and red alder lumber. Upon determination of the yields, cost comparisons can be made of the various grades and grade mixes to obtain the most economical choice for the specific cutting order. Until computerized processing and decision making become commonplace, and for small manufacturers who cannot afford an elaborate system, these nomographs present a useful alternative. FPL(1293).

587. Four paint combinations applied to simulated vertical siding of four wood species have been exposed outdoors in Hawaii since 1964. A three-coat, self-primed latex application is holding up best, with oil base primed latex, three-coat oil base, and two-coat oil base combinations next best in that order. Australian toon is holding paint best, followed by redwood, robusta eucalyptus, and Douglas-fir. For good service, durability, and appearance, three coats of self-primed latex on Australian toon or redwood are recommended. PSW(1296).

588. In pilot tests, introducing an enzyme solution into round sections of Rocky Mountain Douglas-fir markedly improved the creosote treatment of this hard-to-treat species. The enzyme pretreatment increased the depth of creosote penetration to full sapwood uptake, increased the total creosote retention, and produced an essentially "dry" surface. FPL(1300).

589. In the wood density survey of the Southern forest resource, information was collected on the specific gravity of the four major southern pines over their natural growth range. In terms of geographic variability, only slash pine and longleaf pine showed a general trend of specific gravity increasing from north to south. This information contributes to the characterization of this major forest resource. FPL (1302).

590. Northern red oak (*Quercus rubra* L.) lumber from trees with heartwood infected by certain anaerobic bacteria proved more susceptible to honeycomb and ring failure during kiln drying than similar lumber from non-infected heartwood. The implication of this work is that it would be possible to speed up the drying of red oak and decrease drying degrade if boards containing bacterial infections could be sorted out and dried separately. FPL(1304).

591. Little information is available concerning the chemical composition of growth rings with ring shake present as compared with normal hardwood growth rings. Working with scarlet oak and black walnut, it was shown that concentrations of holocellulose, alpha cellulose, lignin, and wood methoxyl differed between "normal" and "shake" growth rings. Abnormal distribution of the various components may weaken the wood

structure and thus be one cause of separation. Knowledge of this type may help in identifying causes of ring shake. NE(1309).

592. Loblolly pine wood was coated with either phenolic varnish, pigmented alkyd paint, or pigmented latex paint and artificially weathered. Observations made under a scanning electron microscope showed each coating to have a different failure mode, with the latex being more durable than the other two. SO(1312).

593. Wood density distribution, or uniformity, is an important characteristic in the determination of wood quality. Studies with ponderosa pine in California have now shown that X-ray techniques can be effectively used to measure wood density distribution across large increment cores. When wood density distribution is considered along with the mean wood density, a useful index to the type of conversion processes and products most suitable for that wood can be constructed. PSW(1024).

594. Gelatinous fibers, usually found in tension wood, cause difficulties in sawing and machining. Paper made from pulpwood containing such fibers has low strength. A high proportion of water-conducting vessels to fibers in pulpwood also reduces the strength of paper made from it. Very high percentages of gelatinous fibers were found in straight-boled, dominant cottonwood trees that were growing rapidly under intensive management. In the same trees, the proportion of vessels in the wood increased with height. These results indicate that the incidence of gelatinous fibers and of xylem vessels should be a primary consideration in evaluating wood quality of trees exhibiting rapid growth. NC(1026,1027).

2. Wood chemistry and fiber products

595. In preparing pulpwood for pulping, the bark is ordinarily removed and not used in the pulp-making process. However, bark does contain fiber. Pulping the bark along with the wood increases the fiber yield from a rough cord of wood by 4 percent. This also has the benefits of eliminating the de-barking operation, eliminating the loss of white-wood during de-barking, and permits the use of small tree branches, tops, and other wood wastes that are difficult to de-bark. To get the additional 4 percent fiber requires 0.2 percent more pulping chemical and 0.2 percent more chlorine bleach. FPL(1313,1326).

596. In the recycling of mixed municipal refuse, reduction of particle size and removal of pieces of plastic are necessary but difficult steps. Size reduction by hammermilling has been practiced on a large scale. Analyses of hammermilled waste show that, except for producing a large percentage of overfine material, hammermilled paper fiber undergoes no important degradation. Two methods have been developed for effectively separating thermoplastic films from wastepaper. Most of the plastics of solid waste are thermoplastic and soften on heating. The first method utilizes the principle of the thermoplastic films adhering to a heated surface. The second consists of suspending the waste mixtures in hot gas to contract or ball up the films followed by air classification or screening. Both methods are practicable and the second can be used immediately with equipment presently available on the market. FPL(1314,1332,1333,1337,1338).

597. New studies on the potential of pulps and wood wastes for ruminant feeds have shown that at high lignin contents, pulps from hardwoods are more digestible by ruminants than softwoods. Below 7 percent lignin, pulps have equal digestibility. Digestibility is inversely proportional to lignin content and is independent of the method (pulping process) of lignin removal. The same relationships were found when lignin was removed by fungi. The rate at which nine white-rot fungi removed lignin and carbohydrates from wood was measured. All fungi removed carbohydrate along with the lignin. A procedure also has been developed for determining the digestibility of cellulose and wood with cellulytic enzymes. It is useful for screening various physical and chemical treatments for improving the digestibility of wood materials. Using this enzyme test, fine grinding was found to be the most effective wood treatment. Dilute alkali and liquid ammonia increased the digestibility of aspen to 50 percent and 36 percent respectively. FPL(1315,1327,1335).

598. Paper testing methods are frequently difficult to reproduce and interpret in meaningful ways. Some advances in this field have been made. Interfiber bonding is an important factor in mechanical and optical properties of paper. Most methods for measuring this property rely on indirect measurements of exposed surface areas. A new method is described whereby surface areas are measured by small X-ray scattering techniques. In ring and flat crush tests on paper the comparability of results using different machines was also examined. Significant differences were found between the flexible and rigid types of testing machines. Suggested modifications of test procedures should include restricting the tests to one type of machine or standardizing machine parameters. FPL(1317,1328).

599. The speed at which corrugated fiberboard can be produced is in part dependent upon the runnability of the corrugated medium itself. The runnability of the corrugated medium is influenced by the web tension during corrugation. Installation of an automatically controlled tension device, electronically coupled to a magnetically operated disk brake system, resulted in accurate measurement and also control of the web tension of the corrugating medium on an experimental fiberboard machine. FPL(1322).

600. In chemical pulping processes, as much as 50 percent of the wood raw material is solubilized and lost to fiber production. Up to half of this is carbohydrate that is removed unnecessarily under the lignin-removal conditions which are usually alkaline. Additional basic information on the complex alkaline degradation reactions of cellulose and other wood carbohydrates has been developed. Oxygen, alkalinity, and higher temperatures increase the rate of cellulose degradation. Neighboring hydroxyl groups promote the breaking of chemical links in cellulose chains. Reactions leading to the formation of butyric acid derivatives and the migration of acyl groups in carbohydrate in wood or pulp have been described. The method is based on dehydrating carbohydrates to furans using sulfuric acid and measuring the resulting furans spectrophotometrically. FPL(1324,1343-1346).

601. The morphology and characteristics of individual wood fibers markedly affects the properties of papers made from those fibers. Paper strengths increase with fiber cell wall thickness to a critical point and then there is an adverse effect. The ratio of fiber length to cell wall thickness appears to be a good indicator for the paper making potential of a pulp. Earlywood papers show consistently higher tensile strengths than latewood papers. Cell wall densities of the two kinds of fibers are the same. Earlywood fibers have more uniform fiber-to-fiber conformability than latewood fibers. The amount of bonding developed in latewood is insufficient to fully utilize greater fiber strength. FPL(1325,1348,1349,1351).

602. Fast-growing, short-rotation crops of poplar and sycamore trees can produce as much as 4 tons per acre per year of biomass useful for pulpwood. This is more than four times the average dry wood weight produced in conventional pulpwood growth. Satisfactory pulps were made from whole

rapid-grown poplar trees and the presence of bark presented no special problems in either pulping or bleaching. Eleven-year-old trees gave the highest yields and stronger papers than younger trees. With sycamore, 4-year-old seedlings gave maximum production. Good quality pulps were made from all of the short-rotation sycamore trees studied. The pulps were found suitable for making acceptable bleached kraft papers and dry-formed hardboards. FPL(1329-1331).

603. The chemical extractives of elm and Eastern white pine were examined further. A symptom of Dutch elm disease is the formation of a brown discoloration in the last one or two annual growth rings. Phenolic substances identical to those extracted from heartwood but not found in sapwood were found in the "brown ring" area. These substances may be related to natural disease resistance factors. Information on the phenolic constituents of various elms can be used to distinquish between anatomically closely related species. The major chemical constituents of extracts of Eastern white pine cortex tissue and a needle polyprenol were characterized. The diterpenes, strobol, strobal, and manoyl oxide were present. FPL(1342,1352).

604. Wax-treated corrugated containers account for 6 to 7 percent of the corrugated containers presently used. This is expected to increase to 10 percent or one million tons in the near future. virtually all of this good-quality, wax-treated material is burned or buried after use rather than recycled since the specialized, solvent extraction equipment necessary for wax removal is not available at most recycling mills. A practical, more generally applicable method for wax removal has been developed that uses conventional mill equipment. Wood fiber reclaimed from such material shows promise as a significant source of fiber for use in corrugating medium either as 100 percent reclaimed fiber or blended with virgin fiber stock to improve properties. FPL(1334).

3. Wood engineering

605. Condensation because of excessive moisture is an aggravating problem for many building owners. A summary report suggests several design alternatives, including vapor barriers, insulation, ventilation, and other construction details to alleviate this problem. FPL(1353).

606. New clear wood mechanical properties were established for five western softwoods. Properties for white and sugar pine, western redcedar, and Port-Orford-cedar were not greatly different than those obtained in 1915. For a fifth, however, subalpine fir, new estimates are significantly higher; 22 percent in modulus of elasticity and 10-14 percent in other properties. Results will ultimately lead to more efficient and reliable structural design with these species. FPL(1355,1356).

607. Paper sacks are an important container for shipping many materials. When sacks fail, valuable contents are lost or damaged. If too much paper is used in constructing sacks, valuable paper resources are wasted. A theoretical design formula has been developed to predict failure of sacks subjected to certain drop heights. This is important to designers, manufacturers, and users. FPL(1365).

608. About 80 percent of the observed variation in tensile strength of dimension lumber can be explained by nondestructive measurements. These include a variety of stiffness measurements, several strength ratio estimators, slope or grain, and, to a small degree, specific gravity. FPL(1371).

609. Lumberlike products can be successfully built from laminated veneers. Strength properties can be maintained if butt joints are properly engineered and fabricated. Tensile strengths can exceed those of solid lumber of equal log grades. FPL(1381).

610. Observations and measurements of moisture contents in wood framing members used in natatoriums reveals that when design standards for occupancy are met, there is no hazard to wood framing or roof. Guidelines are provided to assure proper insulation and ventilation to avoid problems. This information is important to designers, developers, and builders of swimming pools and other wood structures where like humidity and moisture conditions exist. FPL(1383).

611. Hardboard can be used in many engineered structures. Recent studies provide the means for calculating the design characteristics of hardboard used as the webbing for I-beam construction. FPL(1384).

612. Experimental evidence shows that creep deformation of wood is accelerated by both increasing and decreasing moisture exposures, a factor that may be of considerable importance in design of wood structures. The research proposes a kinetic model for creep response of wood which attributes accelerated creep with a changing moisture environment to the diffusion of moisture within loaded specimens. Material basic to development of predictive models for creep of wood in uncontrolled environments is presented. FPL(1385).

613. A limited survey was made of users of the FPL low-cost house designs. Of the 225 homes constructed based on FPL plans, 93 percent reported costs less than $12,000. Ninety-two percent of the homeowners had annual incomes less than $7,000. Conclusions reached indicate that structurally sound houses can be built to meet the needs of most people at prices they can afford. FPL(1387).

614. Timber bridges offer an important alternative to steel or concrete for many sites. Recent research results provide a simplified method for designing wood bridge decks and connector systems using charts, rather than computers. RPL(1390).

615. Nail popping is a frequent, unsightly, and costly defect in resilient tile flooring used in houses. Techniques have been developed which significantly reduce such problems through use of short screws or nails used to fasten underlayment to subflooring. This practice eliminates nail popping resulting from penetration of longer nails into floor joists subject to large dimensional changes, and should be of particular interest to home builders, flooring contractors, and the like. SE(1391-1393).

4. Biological degradation

616. The function of protozoans occurring in the hindgut of the Formosan termite (*Coptotermes formosanus*) is not well understood. Tests were conducted to determine the ability of these termites to break down cellulose and to synthesize lipids with and without the normal gut fauna. Results suggested that the protozoan, *Pseudotrichonympha grassii*, is essential to normal cellulose breakdown, lipid synthesis, and survival of worker Formosan termites. Controlling this protozoan could control the termite. SO(436).

617. Recent studies have provided more information on the importance of protozoan symbionts in the feeding, soldier differentiation, and survival of the Formosan termite workers. Workers with protozoa fed more and survived better than did termites lacking all or some of their protozoan associates. SO(438).

618. Temperature affects the feeding and survival of native and Formosan subterranean termites. Workers lacking normal intestinal protozoa ate less and survived for shorter periods

than did colonies having these organisms. Formosan termites and their associated protozoans were able to survive higher temperature than *Reticulitermes flavipes*. These findings contribute to understanding the distribution, survival, and potential destructiveness of the respective termite species. SO(437,439).

619. The synthesis and subsequent breakdown of lipid materials form an important aspect of termite metabolism. The fatty acids in the body fat of *Reticulitermes flavipes* were determined after feeding on sound wood and wood rotted by the fungus *Lenzites trabea* in the field and laboratory. The major components were palmitic, palmitoleic, stearic, oleic, and linoleic acids. Differences in composition occurred between termite colonies and among termites fed on sound versus rotted wood. This information helps clarify the biochemical relationship between termites and their associated microorganisms. SO(433-435).

620. The psocid *Liposcelis bostrychophilus* is usually associated with anobiid-infested wood, but its role has not been clearly established. Laboratory tests revealed that as psocid numbers increase consumption of beetle eggs also increases. Under natural conditions, psocids apparently contribute in a limited way to the natural regulationg of anobiid infestations SO(442).

621. New, environmentally safe ways for extending the life of wood in use are being sought at the Southern Forest Experiment Station. Trials on malt agar showed that a *Streptomyces* sp. could strongly inhibit growth of decay fungi, but in pine sapwood the antagonist tended to hasten decay rates. This illustrates the need to completely understand systems when formulating biological control technology. SO(608).

622. Several previously undiscovered wood decay fungi have been described and identified by scientists at the Center for Forest Mycology. Identification of the numerous and diverse fungi involved in decay will allow more efficient research on clarifying the complex systems involved in decay processes. FPL(607,611,612).

623. Lignin was found to be selectively removed from wood by certain white-rot fungi. As a result, these woods were more digestible by rumen fluid and a mixture of polysaccharidases. This finding could lead to new uses for unused wood. FPL(610).

5. Prevention and control of wood-destroying organisms

624. Research on preventing decay in wooden structures has been summarized for technical use. Lumber cut from second-growth timber contains mostly sapwood, which is susceptible to decay. Such lumber can be protected by building designs which keep untreated wood dry and by pressure-treating wood with preservatives whenever contact with soil or other sources of moisture cannot be avoided. Increasing the service life of wood is an important step in conserving our valuable natural resources. SO(614,615).

625. Standard ground-board and stake tests to evaluate various soil treatments for protecting wooden structures from subterranean termites have been underway at Gulfport, Mississippi, since 1944. Many chemicals have failed to provide acceptable, long-term protection. Chlordane, aldrin, dieldrin, and heptachlor applied at various concentrations and rates continue to be fully effective after 18-22 years. SO(446).

626. Alternatives for the chlorinated hydrocarbons aldrin, chlordane, dieldrin, and heptachlor are needed for subterranean termite control. Carbamate and organophosphate insecticides have provided full protection for more than 4 years in southern Mississippi tests. Continuing standard slab and ground-board tests will determine the full potential of these materials and provide the necessary efficacy data for registration. SO(443).

627. Destruction of polymeric insulating materials, used on communication lines and in electronic equipment, by termites is of considerable concern in south temperate and tropical environments. Cooperative tests with the U.S. Naval Research Laboratory in Panama and Mississippi showed that polymeric formulations plasticized with tricresyl phosphate fortified with ortho isomer were least damaged by termites after 75 months of exposure, either with or without lindane, dieldrin, and aldrin. Formulations plasticized with tri-p-cresyl phosphate were nearly as good. This information will be utilized in fabricating termite-resistant insulating materials. SO(444).

628. The use of toxic inhibitors to protect cables sheathed or insulated with flexible polyvinyl chloride plastics against termite damage is hazardous in many situations. A cooperative study was undertaken with the Naval Research Laboratory to determine the effectiveness of nontoxic, termite-proof polymers in Panama, Mississippi, and Louisiana. Best results were obtained with ethylene propylene rubber and chlorosulfonated polyethylene formulations, which were highly resistant to termites, innocuous to man, and promised long service life without loss of effectiveness. SO(447).

629. Long-term stockpiling of pulp chips outdoors is facilitated by a newly patented technique which controls chip deterioration. Pulp chips are treated with an aqueous mixture of sodium sulfide and sodium carbonate, both of which are readily available and inexpensive at pulpmill sites. The treatment reduces natural heat evolving processes and wood rot, and reduces losses in stored chips by two-thirds. FPL(618).

630. Increased decay resistance of ammonia- and sodium hydroxide-treated wood was attributed to increased pH and to abnormally high levels of ammonical nitrogen rather than destruction of thiamine. These treatments were effective only against brown-rot fungi under low decay-promoting conditions. Refinements in wood preservation will become more feasible when mechanisms such as these are understood. FPL(616,617).

631. A number of wood-destroying beetles, commonly referred to as powder-post beetles, cause damage to seasoned wood in buildings. A new bulletin answers questions about the appearance, habits, damage, and control of these pests. SO(448).

6. Miscellaneous

632. Lobeline, an alkaloid used in antismoking preparations, usually is extracted from wild plants of *Lobelia inflata*. But harvesting of wild plants is not very profitable because much labor is required to find suitable plants in the widely scattered population. Cultivated lobelia plants in central Kentucky were larger and had a higher lobeline content than wild plants in the same area. Yield estimates indicate that cultivation of lobelia probably will be economically profitable at the current selling price of dried plant material. Analyses showed that *L. cardinalis* and *L. puberula* contain at least as much lobeline as *L. inflata*. Small groups of plants of each of the three species are scattered over the southeastern United States. A potential grower, therefore, can select the species best suited to his local conditions. SE(1034,1035).

D. Marketing Under-Used Species and Residues

633. The conversion of logs into lumber often results in the creation of fine wood residues equivalent to about one-third of total log weight. Profitable markets for fine wood residues are not available in all areas because of the variable character of these materials and economic constraints of handling and transportation. However, there are many potential uses for such residues and managers should carefully evaluate these opportunities within the context of their individual plant circumstances. SE(1140).

634. Prices paid by mills in the northwestern United States for "fiberwood" (chips, sawdust, shavings, and logs) varied widely during 1971 and 1972. Chip prices ranged from $7.75, to $21.25 per bone dry ton (BDT) during 1970-1971. Prices vary by location and species group. Sawdust prices have remained stable within a range of $3 to $7.50 per BDT over the past 10 years. Shavings prices averaged from $5 to $7.25 per BDT in 1970-1971, very close to the price range prevailing over the past decade. PNW(1142).

635. A detailed evaluation of a new hardwood sawmill system which includes a debarker, automatic carriage and circular headrig, chipper, and a sawyer-controlled vertical edger, showed that low-grade hardwood logs (Grade 3 and below) can be processed profitably. This new sawmill system should find wide application in the Appalachian region where about 70 percent of the sawtimber consists of Grade 3 and poorer quality material. NE(1143).

636. Until recently, bark has been the most difficult wood processing residue to convert into useable, marketable products. However, bark producers now are effectively developing markets for bark as mulches and soil conditioners. This article describes these markets and discusses the ways in which bark is used effectively for mulches and soil amendments. NE(1145).

E. Supply, Demand, and Price Analysis

637. A computer model was developed for projecting long-run demand for housing by type of dwelling unit and region. The model also provides a framework for projecting future housing demand under specified assumptions relating to population and economic growth. Tabular data showing these respective projections are presented. Since residential housing is the market for more than a third of all lumber and plywood consumed annually in the United States, trends in housing construction are of major significance in assessing future demand for timber products from the nation's forest resources. NC(1169).

638. More than 42 percent of the $2.1 billion dollars of Canadian forest products shipped into the United States in 1969 entered through northcentral region Custom Districts. Three commodities—newsprint, softwood lumber, and woodpulp--accounted for 89 percent of the value of all forest products imported from Canada and 87 percent of all Canadian forest products that come imto the northcentral region. NC(1152).

639. Construction of apartments in the United States in 1969 involved an average use per apartment unit of 3,886 board feet of lumber, 1,879 square feet of plywood (3/8-inch basis), 49 square feet of hardboard (1/8-inch basis), 52 square feet of particleboard (3/4-inch basis), and 45 square feet of insulation board (1/2-inch basis). These wood-use data were derived from a sample of apartment projects for which architectural plans and specifications were completed during 1969. Comparisons with data from a similar 1962 study indicate that while plywood use per housing unit has increased slightly, lumber use has declined by about 14 percent. WO(1176).

640. An analysis of the demand and price situation for forest products provides up-to-date information on timber product markets as of 1971. Information is presented on price and consumption trends for lumber, plywood, pulp and paper, and miscellaneous products, including naval stores. WO(1158).

641. In 1972, nearly 300 secondary wood-processing firms were in operation in West Virginia. A directory classifies these forms into categories and shows their location, raw material requirements, and type of product manufactured. This information aids area lumber and other primary processors in finding markets for their products, and also assists consumers of secondary products in locating sources of supply. NE(1156).

642. An analysis of the production and shipments of mobile homes in the contiguous 48 states during the period 1966-69 disclosed a shifting pattern of geographical concentration of production. On the basis of a model which relates mobile home use to state population characteristics, projections of mobile home shipments were made for the 1971-75 period. Since mobile homes now are a significant component of housing demand, these data will be helpful to wood materials suppliers in planning production and marketing strategies. SE(1162,1164).

643. A detailed time and motion study on sample logging operations in West Virginia indicated that in the log skidding phase, cost savings can be greater from improved machine operator practices than from upgrading skidding equipment. Since skidding represents one of the major costs in logging, logging managers should consider skidder operator training or production bonuses as promising opportunities to reduce harvesting costs. NE(1170).

644. An adaptation of a method developed in Sweden was used in an analysis in the Lake States relating timber inventory volumes to transport distances to specified delivery points. The procedure developed shows promise as a useful aid in analyzing area timber resources information and making decisions on wood procurement strategies. NC(1148).

645. A quarterly marketing report on the Northwest forest industries presents current timber industry information on production; prices; employment; trade; volume and average prices of stumpage sold by public agencies; and other related items. PNW(1153,1154,1159,1160).

646. The tariff on imported maple syrup products was abolished in 1972, and domestic maple syrup producers have anticipated market changes due to its removal. However, an analysis of syrup price and tariff levels since World War II concludes that no substantial increase of shipments of maple syrup products from Canada into the United States is likely as a result of the tariff removal. NE(1173,1174).

647. The major end-product for hardwood plywood (usually birch) stock panels in 1970 was kitchen cabinets. Over one-third of the wholesalers contacted in a study of the hardwood stock panel market reported that stock panel sales have been declining because of market inroads by plastic-overlaid particleboard panels. NC(1167).

F. Marketing—General

648. Wood craftsmen in Kentucky increased their production, received higher returns from sale of their crafted wood

products, and improved their skills by belonging to a handicraft cooperative. Some of the problems identified in this study of a handicraft cooperative were the underutilization of fixed assets, the need for more business knowledge and improved training facilities for craftsmen. NC(1187).

649. A bibliography of current published information on Christmas tree production and marketing provides a means of quickly and easily locating up-to-date information on this industry. This information can assist Christmas tree producers, wholesalers, and retailers in making better decisions on production and marketing practices. NE(1189).

650. An analysis of the commercial potential of a new kiln-drying method for hardwood lumber involving presurfacing green lumber and kiln-drying on an accelerated schedule, showed several advantages. this new drying method, developed by the Forest Products Laboratory, reduces drying time by nearly one-half permits a higher kiln-capacity use, produces slightly lower lumber degrade loss, and reduces fuel and power costs. NE(1185).

651. Model building codes covering high-density, commercial sections of urban areas usually contain provisions for "fire districts," "fire limits," or "fire zones." Within such areas, construction materials and methods are closely regulated so as to minimize the danger of fire. Results from regional studies of fire zones review the purpose, use, and implications for construction methods and materials use. SE(1178-1181).

FOREST ENVIRONMENT RESEARCH

WATERSHED MANAGEMENT RESEARCH

Controlling soil erosion

1. Clayton, J.L., and J.F. Arnold
   1972. Practical grain size, fracturing density, and weathering classification of intrusive rocks of the Idaho Batholith. USDA For. Serv. Gen. Tech. Rep. INT-2, 17p. (INT)

2. Curtis, W.R.
   1971. Vegetating strip-mine spoils for runoff and erosion control. Proc. Revegetation and Economic Use of Surface Mined Land and Mine Refuse Symp., Dec. 2-4, 1971, Pipestem State Park, W. Va., p. 40-41, W. Va. Univ., Morgantown. (NE)

3. DeBano, L.F., and R.M. Rice
   1971. Fire in vegetation management: Its effect on soil. Am. Soc. Civ. Engin. Proc. of Symp. on Interdisciplinary Aspects of Watershed Management, Bozeman, Mont., 1970: 327-346. (PSW)

4. Dyrness, C.T.
   1972. Soil surface conditions following balloon logging. USDA For. Serv. Res. Note PNW-182, 7p. (PNW)

5. Klock, G.O.
   1973. Mission Ridge - a case history of soil disturbance and revegetation of a winter sports area development. USDA For. Serv., National Winter Sports Symp., Feb. 26 - Mar. 2, 1973, Denver, Colo., 17p. (PNW)

6. Megahan, W.F.
   1972. Volume weight of reservoir sediment in forested areas. J. Hydraul. Div., ASCE, Vol. 98, No. HYS, Proc. Pap. 9129, p. 1335-1342. (INT)

7. Megahan, W.F.
   1972. Subsurface flow interception by a logging road in mountains of Central Idaho. Am. Water Resour. Assoc. Symp. on Watersheds in Transition Proc., p. 350-356, Colo. State Univ. (INT)

8. Megaham, W.F., and W.J. Kidd
   1972. Effect of logging roads on sediment production rates in the Idaho Batholith. USDA For. Serv. Res. Pap. INT-123, 14p. (INT)

9. Mersereau, R.C., and C.T. Dyrness
   1972. Accelerated mass wasting after logging and slash burning in western Oregon. J. Soil and Water Conserv. 27(3): 112-114. (PNW)

10. Rice, R.M., J.S. Rothacher, and W.F. Megahan
    1972. Erosional consequences of timber harvesting: An appraisal. Am. Water Resour. Assoc. Symp. on Watersheds in Transition Proc., p. 321-329, Colo. State Univ. (PSW)

11. Swanston, D.N.
    1972. Practical analysis of landslide potential in glaciated valleys of sourtheastern Alaska and similar sub-arctic or alpine regions. Arctic and Mountain Environments Symp., Mich. State Univ., April 22-23, 1972. (PNW)

12. Ursic, S.J., and P.D. Duffy
    1972. Hydrologic performance of eroded lands stabilized with pine. Miss. Water Resour. Conf. Proc., p. 203-216. (SO)

13. Vogel, W.G.
    1971. Needs in revegetation research on surface-mined lands. In Proc. Revegetation and Economic Use of Surface-Mined Land and Mine Refuse Symp., Dec. 2-4, Pipestem State Park, W. Va., p.17-18. (NE)

14. Vogel, W.G.
    1973. The effect of herbaceous vegetation on survival and growth of trees planted on coal-mine spoils. In Proc. Research and Applied Technology Symp. on Mined-Land Reclamation, Mar. 7-8, 1973, Pittsburgh, Pa., p.197-207, Bituminous Coal Research Inc., Monroeville, Pa. (NE)

15. Yamamoto, T., and H.W. Anderson
    1973. Splash erosion related to soil erodibility indices and other forest soil properties in Hawaii. Water Resour. Res. 9(2): 336-345. (PSW)

Chemicals in the forest environment

16. Ballard, T.M.
    1971. Role of humic carrier substances in DDT movement through forest soil. Soil Sci. Soc. Am. Proc. 35:146-147. (PNW)

17. Bollen, W.B., and C.M. Tu
    1972. Effects of an organotin on microbial activities in soil. Tin and Its Uses, Q. Bull. Tin Res. Inst., No. 94, p. 13-35. (PNW)

18. Curtis, W.R.
    1972. Chemical changes in streamflow following surface mining in eastern Kentucky. Mellon Institute Coal Mine Drainage Research Symp. 4:19-31. Bituminous Coal Research, Inc. (NE)

19. DeByle, N.V., and P.E. Packer
    1972. Plant nutrient and soil losses in overland flow from burned forest clearcuts. Am. Water Resour. Assoc. Symp. on Watersheds in Transition Proc., p. 296-307. Colo. State Univ. (INT)

20. Dickinson, J.O.
    1972. Toxicity of the arsenical herbicide monosodium acid methanearsonate in cattle. Am. J. Vet. Res. 33(9):1889-1892. (PNW)

21. Juntunen, E.T., and L.A. Norris
1972. Field application of herbicides—avoiding danger to fish. Agric. Exp. Sta., Oregon State Univ., Corvallis, OR, Special Rep. 354, 26 p. (PNW)

22. Li, C.Y., K.C. Lu, J.M. Trappe, and W.B. Bollen
1972. *Poria weirri*—inhibiting and other phenolic compounds in roots of red alder and Douglas-fir. Microbios 5:65-68. (PNW)

23. Massey, H.F.
1972. pH and soluble Cu, Ni, and Zn in eastern Kentucky coal mine spoil materials. Soil Sci. 114:217-221. (NE)

24. Pierce, R.S., C.W. Martin, C.C. Reeves, G.E. Likens, and F.H. Bormann
1972. Nutrient loss from clearcutting in New Hampshire. Am. Water Resour. Assoc. Symp. Watersheds in Transition Proc., p. 285-295. Colo. State Univ. (NE)

25. Stutzbach, S.J., A.L. Leaf, and R.E. Leonard
1972. Variation in the forest floor under a red pine plantation. Soil Sci. 114: 24-28. (NE)

26. Tarrant, R.F., D.G. Moore, W.B. Bollen, and B.R. Loper
1972. DDT residues in forest floor and soil after aerial spraying, Oregon—1965-1968. Pestic. Monit. J. 6:65-72. (PNW)

Improving water yields

27. Boelter, D.H.
1972. Preliminary results of water level control on small plots in a peat bog. Fourth Int. Peat Congr. Proc., Otaniemi, Finland, June 25-30, 1972; Vol. 3: 347-354. (NC)

28. Federer, C.A., R.S. Pierce, and J.W. Hornbeck
1972. Snow management seems unlikely in the Northeast. Am. Water Resour. Assoc. Symp. on Watersheds in Transition Proc., p. 212-219. Colo. State Univ. (NE)

29. Gary, H.L.
1972. Rime contributes to water balance in high-elevation aspen forests. J. For. 70: 93-97. (RM)

30. Halverson, H.G.
1972. Seasonal snow surface energy balance in a forest opening. US Atomic Energy Comm., Div. of Tech. Inf., TID-26242, 73 p. (PSW)

31. Harris, A.R.
1972. Relation of wind exposure and forest cutting to changes in snow accumulation. Proc. Int. Symp. on Role of Snow and Ice in Hydrol., Banff, Alberta, Can., Sept. 1972, 11 p. (NC)

32. Hibbert, A.R., and P.A. Ingebo
1971. Chaparral treatment effects on streamflow. 15 Annu. Ariz. Watershed Symp. Proc., Sept. 22, p.25-34. (RM)

33. Leaf, C.F., and G.E. Brink
1972. Simulating effects of harvest cutting on snowmelt in Colorado subalpine forest. Am. Water Resour. Assoc. Symp. on Watersheds in Transition Proc., p. 191-196. Colo. State Univ. (RM)

34. Martin, C.W., and J.W. Hornbeck
1972. Lysimeter snowmelt and streamflow on forested and cleared sites. 1972 Eastern Snow Conf. Proc., p. 111-118. (NE)

35. Martinelli, M., Jr.
1972. Snow fences for influencing snow accumulation. In Proc. Int. Symp. on Role of Snow and Ice in Hydrol., Banff, Alterta, Can., Sept. 1972, Sess. WMO-5, 6p. (Preprint). (RM)

36. Owston, P.W., J.L. Smith, and H.G. Halverson
1972. Seasonal water movement in tree stems. For. Sci. 18(4):266-272. (PSW)

37. Rich, L.R.
1972. Managing a ponderosa pine forest to increase water yields. Water Resour. Res. 8(2): 422-428. (RM)

38. Sartz, R.S.
1972. Soil water depletion by a hardwood forest in southwestern Wisconsin. Soil Sci. Soc. Amer. Proc. 36: 961-964. (NC)

39. Sartz, R.S., and A.R. Harris
1972. Growth and hydrologic influence of European larch and red pine 10 years after planting. USDA For. Serv. Res. Note NC-144, 4 p. (NC)

40. Satterlund, D.R., and H.F. Haupt
1972. Vegetation management to control snow accumulation and melt in the northern Rocky Mountains. Am. Water Resour. Assoc. Symp. on Watersheds in Transition Proc., 200-205. Colo. State Univ. (RM)

41. Schmidt, R.A., Jr.
1972. Sublimation of wind-transported snow—a model. USDA For. Serv. Res. Pap. RM-90, 24p. (RM)

42. Smith, J.L., and H.G. Halverson
1970. Suppression of evaporation losses from snowpacks. Am. Soc. Civ. Engin. Proc. of Symp. on Interdisciplinary Aspects of Watershed Management, Bozeman, Mont., 1970:5-25. (PSW)

43. Smith, J.L., H.G. Halverson, and R.A. Jones
1972. Central Sierra profiling snow gage: a guide to fabrication and operation. US Atomic Energy Comm., Div. of Tech. Inf., TID-25986, 53 p. (PSW)

44. Smith, J.L., H.G. Halverson, and R.A. Jones
1972. Development of a radioisotope profiling snow gage. US Atomic Energy Comm., Div. of Tech. Inf. TID-25987, 86 p. (PSW)

45. Sommerfield, R.A., and F. Wolfe, Jr.
1972. A centrifugal tensile tester for snow. USDA For. Serv. Res. Note RM-227, 4 p. (RM)

46. Tabler, R.D., and R.A. Schmidt, Jr.
1972. Weather conditions that determine snow transport distances at a site in Wyoming. In Proc. Int. Symp. on Role of Snow and Ice in Hydrol., Banff, Alberta, Can., Sept. 1972, 11 p. (RM)

47. Ursic, S.J., and P.D. Duffy
1972. Hydrologic performance of eroded lands stabilized with pine. Miss. Water Resour. Conf. Proc., p. 203-216. (SO)

48. Verry, E.S.
1972. Effect of an aspen clearcutting on water yield and quality in northern Minnesota. Am. Water Resour. Assoc. Symp. on Watersheds in Transition Proc., p. 276-284, Colo. State Univ. (RM)

Managing, rehabilitating, and improving watersheds

49. Aldon, E.F.
1972. Critical soil moisture levels for field planting fourwing saltbush. J. Range Manage. 25: 311-312. (RM)

50. Aldon, E.F., and G. Garcia
1972. Vegetation changes as a result of soil ripping on the Rio Puerco in New Mexico. J. Range Manage. 25: 381-383. (RM)

51. Anderson, H.W.
1972. Water yield as an index of lee and windward topographic effects on precipitation. Proc. Symp. on Distribution of Precipitation in Mountainous Areas, Geilo, Norway, July 31-Aug. 5, 1972, Vol. 2, p. 346-358, World Meteorol. Organ., Geneva, No. 326, 1972. (PSW)

52. Anonymous
1971. A blizzard wizard's role in snow control. The Highwayman 21(4): 2-5. (RM)

53. Anonymous
1972. Fencing parries winter's thrusts. The Highwayman 22(1): 12-14. (RM)

54. Anonymous
1972. Test proves snow fences at road cuts reduce plowing. Rural and Urban Roads 10(8): 56-57. (RM)

55. Anonymous
1972. Studying the problem of drifting snow. Public Works 103(8): 70-72. (RM)

56. Bethlahmy, N.
1972. Hydrograph analysis: A computerized separation technique. USDA For. Serv. Res. Pap. INT-122, 19 p. (INT)

57. Boelter, D.H.
1972. Water table drawdown around an open ditch in organic soils. J. Hydrol. 15: 329-340. (NC)

58. Brown, J.M.
1972. The effect of overstory removal upon surface wind in a black spruce bog. USDA For. Serv. Res. Note NC-137, 2 p. (NC)

59. Brown, J.M.
1972. Effect of clearcutting a black spruce bog on net radiation. For. Sci. 18: 273-277. (NC)

60. Campbell, C.J., and C.P. Pase
1972. Pressure bomb measures changes in moisture stress of birchleaf mountain mahogany after partial crown removal. USDA For. Serv. Res. Note RM-221, 4 p. (RM)

61. Campbell, R.E.
1972. Prediction of air temperature at a remote site from official weather station records. USDA For. Serv. Res. Note RM-223, 4 p. (RM)

62. Copeland, O.L., and P.E. Packer
1972. Land use aspects of the energy crisis and western mining. J. For. 70(11): 671-675. (INT)

63. Curtis, W.R.
1972. Strip-mining increases flood potential of mountain watersheds. Am. Water Resour. Assoc. Symp. on Watersheds in Transistion Proc., p. 357-360, Colo. State Univ. (NE)

64. Davidson, W.H., and G. Davis
1972. Sprouting of thinned hybrid poplars on bituminous strip-mine spoils in Pennsylvania. USDA For. Serv. Res. Note NE-147, 6 p. (NE)

65. Dickerson, B.P.
1972. Changes in the forest floor under upland oak stands and managed loblolly pine plantations. J. For. 70: 560-562. (SO)

66. Doty, R.D.
1972. Soil water distribution on a contour trenched area. USDA For. Serv. Res. Note INT-163, 6 p. (INT)

67. Douglass, J.E.
1972. Annotated bibliography of publications on watershed management by the Southeastern Forest Experiment Station, 1928-1970. USDA For. Serv. Res. Pap. SE-93, 47 p. (SE)

68. Douglass, J.E., and W.T. Swank
1972. Streamflow modifications through management of eastern forests. USDA For. Serv. Res. Pap. SE-94, 15 p. (SE)

69. Duffy, P.D., and T.R. Dell
1972. System for measuring and computing ephemeral runoff from small watersheds. USDA For. Serv. Res. Note SO-138, 4 p. (SO)

70. Dyrness, C.T.
1973. Early stages of plant succession following logging and burning in the western Cascades of Oregon. Ecol. 54(1): 57-59. (PNW)

71. Farmer, E.E., and J.E. Fletcher
1972. Some intrastorm characteristics of high intensity rainfall bursts. Proc. Symp. on Distribution of Precipitation in Mountainous Areas, Geilo, Norway, July 31-Aug. 5, 1972, Vol. 2, p. 525-531, World Meteorol. Organ., Geneva. (INT)

72. Federer, C.A., R.S. Pierce, and J.W. Hornbeck
1972. Snow management seems unlikely in the Northeast. Am. Water Resour. Assoc. Symp. on Watersheds in Transition Proc., p. 212-219 Colo. State Univ. (NE)

73. Franklin, J.F., F.C. Hall, C.T. Dyrness, and C. Maser
1972. Federal Research Natural Areas in Oregon and Washington: a guidebook for scientists and educators. USDA For. Serv., Pacific Northwest Forest and Range Exp. Stn., 498 p. (PNW)

74. Gary, H.L.
1971. Seasonal diurnal changes in moisture contents and water deficits of Engelmann spruce needles. Bot. Gaz. 132: 327-332. (RM)

75. Gee, G.W., and C.A. Federer
1972. Stomatal resistance during senescence of hardwood leaves. Water Resour. Res. 8: 1456-1460. (NE)

76. Haeffner, A.D., and A.H. Barnes
1972. Photogrammetric determinations of snow cover extent from uncontrolled aerial photographs. Am. Soc. Photogramm. (Columbus, Ohio, Oct. 1972) ASP Tech. Sess. Proc., p. 319-340. (RM)

77. Harris, A.R.
1972. Infiltration rate as affected by soil freezing under three cover types. Soil Sci. Soc. Am. Proc. 36: 489-492. (NC)

78. Heede, B.H.
1972. Influences of a forest on the hydraulic geometry of two mountain streams. Water Resour. Bull. 8(3): 523-529. (RM)

79. Heede, B.H.
1972. Flow and channel characteristics of two high mountain streams. USDA For. Serv. Res. Pap. RM-96, 12 p. (RM)

80. Helvey, J.D.
1972. First year effects of wildfire on water yield and stream temperature in north central Washington. Am. Water Resour. Assoc. Symp. on Watersheds in Transition Proc., p. 308-312. (PNW)

81. Helvey, J.D., J.D. Hewlett, and J.E. Douglass
1972. Predicting soil moisture in the southern Appalachians. Soil Sci. Soc. Am. Proc. 36(6): 954-959. (SE)

82. Herrington, L.P., G.E. Bertolin, and R.E. Leonard
1972. Microclimate of a suburban park. Am. Meteorol. Soc. Conf. on Urban Environment Proc., p. 43-44. (NE)

83. Herrington, L.P., R.E. Leonard, J.E. Hamilton, and G.M. Heisler
1972. The response of moving radiometers. Boundry-Layer Meteorol. 2(1972): 395-405. (NE)

84. Johnston, R.S., and R.D. Doty
1972. Description and hydrologic analysis of two small watersheds in Utah's Wasatch mountains. USDA For. Serv. Res. Pap. INT-127, 53 p. (INT)

85. Klock, G.O., and W.B. Fowler
1972. An inexpensive water sample. USDA For. Serv. Res. Note PNW-188, 6 p. (PNW)

86. Knipe. O.D.
1973. Western wheatgress germination as related to temperature, light, and moisture. J. Range Manage. 26: 68-69. (RM)

87. Knipe, O.D., and H.W. Springfield
1972. Germinable alkali sacaton seed content of soils in the Rio Puerco basin, west central New Mexico. Ecol. 53: 965-968. (RM)

88. Leaf, C.F., and G.E. Brink
1972. Annual streamflow summaries from four subalpine watersheds in Colorado. USDA For. Serv, General Tech. Rep. RM-1, 24 p. (RM)

89. Leaf, C.F., and G.E. Brink
1973. Computer simulation of snowmelt within a Colorado subalpine watershed. USDA For. Serv. Res. Pap. RM-99, 22 p. (RM)

90. Leaf, C.G., and J. Kovner
1972. Sampling requirements for areal water equivalent estimates in forested subalpine watersheds. Water Resour. Res. 8(3): 713-716. (RM)

91. Leonard, R.E.
1972. Making our lives more pleasant—plants as climate changers. USDA Yearbook 1972: 5-9. (NE)

92. Leonard, R.E., and C.A. Federer
1972. Soil microclimate. In Proc. First Soil Microcummunities Conf., D.L. Dindal (Ed.), CONF-711018 USAEC, Natl. Tech. Inf. Serv., US Dep. Commerce, Springfield, VA. (NE)

93. Lull, H.W., and K.G. Reinhart
1972. Forests and floods in the eastern United States. USDA For. Serv., U.S. Dep. Commerce, Springfield, VA. (NE)

94. Martinelli, M., Jr.
1972. Take the plunge. Ski Area Manage. 11(1): 27-28. (RM)

95. Megahan, W.F.
1972. Logging, erosion, sedimentation—are they dirty words? J. For. 70: 403-407. (INT)

96. Miller, R.L., and F.R. Larson
1973. A cost analysis of clearing a ponderosa pine watershed. USDA For. Serv. Res. Note RM-231, 7 p. (RM)

97. O'Connell, P.F., and H.E. Brown
1972. Use of production functions to evaluate multiple use treatments on forested watersheds. Water Resour. Res. 8(5): 1188-1198. (RM)

98. Orr, H.K., and T. VanderHeide
1973. Water yield characteristics of three small watersheds in the Black Hills of South Dakota. USDA For. Serv. Stn. Pap. RM-100, 8 p. (RM)

99. Pase, C.P.
1972. Litter production by oak-mountain mahogany chaparral in central Arizona. USDA For. Serv. Res. Note RM-214, 7 p. (RM)

100. Patric, J.H., and G.R. Trimble
1972. Transition in research on small forested watersheds in West Virginia. Am. Water Resour. Assoc. Symp. on Watersheds in Transition Proc., Series 14, p. 272-275. (NE)

101. Plass, W.T.
1971. Highwalls—An environmental nightmare. In Proc. Revegetation and Economic Use of Surface-Mined Land and Mine Refuse Symp., Dec. 2-4, 1971, Pipestem State Park, W. Va., p. 9-13. (NE)

102. Plass, W.T.
1972. Fertilization treatments increase black locust growth on extremely acid surface-mine spoils. Tree Planters' Notes 23(4): 10-12. (NE)

103. Rice, R.M.
1972. Using canonical correlations for hydrological predictions. Bulletin of the IASH, XVII 3 10/1972: 315-321. (PSW)

104. Rice, R.M., J.S. Rothacher, and W.F. Megahan
1972. Erosional consequences of timber harvesting: an appraisal. Am. Water Resour. Assoc. on Watersheds in Transition Proc., p. 312-329. (PSW)

105. Richardson, B.Z., and E.R. Burroughs, Jr.
1972. Effects of air gaps and saturated voids on accuracy of neutron moisture measurements. USDA For. Serv. Res. Pap. INT-120. (INT)

106. Rogers, J.J.
1972. A methodology for testing system theoretic models. In Proc. Int. Symp. on Uncertainties in Hydrologic and Water Resource Systems, Vol. 3, Dec. 11-14, 1972, Tucson, AZ, p. 1029-1038. (RM)

107. Rothacher, J.
1970. Managing forest land for water quality. In Proc. Joint FAO/USSR Int. Symp. on Forest Influences and Watershed Management, Moscow, Aug. 17-Sept. 6, 1970, Publ. by FAO, Rome, p. 232-244. (PSW)

108. Schier, G.A.
1973. Apical dominance in multishoot cultures from aspen roots. For. Sci. 18(2): 147-149. (INT)

109. Schier, G.A.
1973. Effect of abscisic acid on sucker development and callus formation on excised roots of *Populus tremuloides*. Physiol. Plant. 28: 143-145. (INT)

110. Schier, G.A.
1973. Origin and development of aspen root suckers. Can. J. For. Res. 3(1):45-53. (INT)

111. Schier, G.A.
1973. Effects of gibberellic acid and an inhibitor of gibberellin action on suckering from aspen root cuttings. Can. J. For. Res. 3(1): 39-44. (INT)

112. Solomon, M.J., and E.A. Hansen
1972. Canoeist suggestions for stream management in the Manistee National Forest of Michigan. USDA For. Serv. Res. Pap. NC-77, 10 p. (NC)

113. Swank, W.T., N.B. Goebel, and J.D. Helvey
1972. Interception loss in loblolly pine stands of the South Carolina Piedmont. J. Soil and Water Conserv. 27: 160-164. (SE)

114. Swanson, R.H.
1972. Water transpired by trees is indicated by heat pulse velocity. Agric. Meteorol. 10: 277-281. (RM)

115. Tabler, R.D., and D.L. Veal
1971. Effect of snow fence height on wind speed. Int. Assoc. Sci. Hydrol. Bull. XVI (4): 49-56. (RM)

116. Tarrant, R.F.
1972. Managing young forests in the Douglas-fir region. Proc. Symp., June 15-18, 1970, Alan B. Berg, (Ed.), Oreg. State Univ., Sch. For. (PNW)

117. Tarrant, R.F., and J.M. Trappe
1971. The role of *Alnus* in improving the forest environment. Plant and Soil, Special Volume 1971, 335-348. (PNW)

118. Thompson, J.R., and A.D. Ozment
1972. The Rocky Mountain millivolt integrator for use with solar radiation sensors. USDA For. Serv. Res. Note RM-225, 8 p. (RM)

119. Tiedemann, A.R.
1972. Soil properties and nutrient availability in tarweed communities of central Washington. J. Range Manage. 25: 438-443. (PNW)

120. Tiedemann, A.R., and J.O. Klemmedson
1973. Effect of mesquite on physical and chemical properties of the soil. J. Range Manage. 26: 27-29. (PNW)

121. Tiedemann, A.R., and J.O. Klemmedson
1973. Nutrient availability in desert grassland soils under mesquite (*Prosopis juliflora*) trees and adjacent open areas. Soil Sci. Soc. Am. Proc. 37: 107-111. (PNW)

122. Tiedemann, A.R., G.O. Klock, H.W. Berndt, and F.C. Hall
1972. Meeks Table Research Natural Area. In Federal Research Natural Areas in Oregon and Washington: a guidebook for scientists and educators, prepared by Jerry F. Franklin, Frederick C. Hall, C.T. Dyrness, and Chris Maser, USDA For. Serv. PNW Forest and Range Exp. Stn., 498 p. (PNW)

123. Walkotten, W.J.
1972. A recording soil moisture tensiometer. USDA For. Serv. Res. Note PNW-180, 12 p. (PNW)

124. Williams, G.P., Jr.
1973. Changed spoil dump shape increases stability on contour strip mines. In Proc. Research and Applied Technology Symp. on Mined-Land Reclamation, Mar. 7-8, 1973, Pittsburgh, PA., p. 243-249, Bituminous Coal Research Inc., Monroeville, PA. (NE)

125. Yamamoto, T., and H.K. Orr
1972. Morphometry of three small watersheds, Black Hills, South Dakota, and some hydrologic implications. USDA For. Serv., Rocky Mt. Forest and Range Exp. Stn., Res. Pap. RM-93, 15 p. (RM)

Preventing watershed damage

126. Anderson, H.W.
1972. Major floods, poor land use delay return of sedimentation to normal rates. USDA For. Serv. Res. Note PSW-268, 4 p. (PSW)

127. Brown, J.M.
1972. Effect of clearcutting a black spruce bog on net radiation. For. Sci. 18: 273-277. (NC)

128. Brown, J.M.
1972. The effect of overstory removal upon surface wind in a black spruce bog. USDA For. Serv. Res. Note NC-137, 2 p. (NC)

129. Clayton, J.L.
1972. Salt spray and mineral cycling in two California coastal ecosystems. Ecol. 53: 74-81. (INT)

130. Lull, H.W., and K.G. Reinhart
1972. Forests and floods in the eastern United States. USDA For. Serv. Res. Pap. NE-226, 94 p. (NE)

131. Martinelli, M., Jr.
1972. Simulated sonic boom as an avalanche trigger. USDA For. Serv. Res. Note RM-224, 7 p. (RM)

132. Martinelli, M., Jr.
1973. Avalanche. 1973 update for McGraw-Hill Yearbook of Science and Technology (a requested article). (RM)

133. Navon, D.I., and H.W. Anderson
1972. Modern techniques in planning the management of wildland resources. Proc. Joint FAO/USSR Int. Symp. on Forest Influences and Watershed Management, Moscow, Aug. 17 - Sept. 6, 1970, p. 359-368. Publ. by FAO, Rome. (PSW)

134. Orr, H.K.
1973. The Black Hills (South Dakota) floods of June 1972: impacts and implications. USDA For. Serv. Tech. Rep. RM-2, 12 p. (RM)

135. Packer, P.E.
1972. Site preparation in relation to environmental quality. In Maintaining Productivity of Forest Soils, Annu. Meet. West. For. and Conserv. Assoc. Proc. 1971:23-28. (INT)

136. Perla, R.E.
1972. Generalization of Haefeli's creep angle analysis. J. Glaciol. 11(63): 447-450. (RM)

137. Rice, R.M., and J.S. Krammes
1971. Mass-wasting processes in watershed management. Am. Soc. Civ. Engin. Proc. of Symp. on Interdisciplinary Aspects of watershed Management, Bozemen, Mont., 1970: 231-259. (PSW)

138. Smith, F.W.
1972. Elastic stresses in layered snow packs. J. Glaciol. 11(63): 407-414. (RM)

139. Williams, K.
1972. Avalanches in our western mountains: what are we doing about them? Weatherwise 25(5): 220-227. (RM)

Water quality

140. Aubertin, G.M., and D.W. Smith
1972. Streamflow quality after urea fertilization of a forested watershed. Agron. Abstracts, Amer. Soc. Agron. Meet., Miami Beach, Fla., 10/29-11/2, p. 175. (NE)

141. Branson, B.A., and D.L. Batch
1972. Effects of strip mining on small-stream fishes in east-central Kentucky. Proc. Biol. Soc. of Washington, Vol. 84, No. 59, 507-518. (NE)

142. Fredriksen, R.L.
1972. Nutrient budget of a Douglas-fir forest on an experimental watershed in western Oregon. In Proc. Research on Coniferous Forest Ecosystems--A Symposium, Bellingham, Wash., Mar. 23-24, 1972. Publ. by USDA For. Serv., Pacific Northwest Forest and Range Exp. Stn. (PNW)

143. Massey, H.F., and R.I. Barnhisel
1972. Copper, nickel, and zinc released from acid coal mine spoil materials of eastern Kentucky. Soil Sci. 113: 207-212. (NE)

144. Moore, D.G.
1972. Fertilization and water quality. In Maintaining Productiivity of Forest Soils, Annu. Meet. West. For. and Conserv. Assoc. Proc. 1971: p. 28-31. (PNW)

145. Norris L.A., and D.G. Moore
1972. The entry, fate and toxicity of forest chemicals in streams. IN Forestry and Water Quality, p. 45-67. Oreg. State Univ., Sch. For. (PNW)

146. Reinhart, K.G.
1972. Effects of clearcutting upon soil/water relations. IN A perspective on Clearcutting in a Changing World, p. 67-74. Applied Forestry Res. Inst., N.Y. State Coll. For., Misc. Rep. 4. (NE)

147. Tarrant, R.F.
1970. Man-caused fluctuations in quality of water from forested watersheds. Proc. Joint FAO/USSR Int. Symp. on Forest Influences and Watershed Management, Moscow, Aug. 17 - Sept. 6, 1970, p. 209-218. Publ. by FAO, Rome. (PNW)

WILDLIFE HABITAT AND RANGE RESEARCH

Evaluating and improving animal habitat resources

148. Anderson, A.E., D.E. Medin, and D.C. Bowden
1972. Carotene and vitamin A in the liver and blood serum of a Rocky Mountain mule deer, *Odocoileus hemionus hemionus*, population. Comp. Biochem. Physiol. 41B: 745-758. (INT)

149. Anderson, A.E., D.E. Medin, and D.C. Bowden
1972. Indices of carcass fat in a Colorado mule deer population. J. Wildl. Manage. 36: 579-594. (INT)

150. Anderson, A.E., D.E. Medin, and D.C. Bowden
1972. Mule deer fecal group counts related to site factors on winter range. J. Range Manage. 25: 66-68. (INT)

151. Anderson, A.E., and D.E. Medin, Bowden, D.C.
1972. Mule deer numbers and shrub yield-utilization on winter range. J. Manage. 36: 571-578. (INT)

152. Baker, M.F., and N.C. Frischknecht
1973. Small mammals increase on recently cleared and seeded juniper rangeland. J. Range Manage. 26(2): 101-103. (INT)

153. Books, D.J.
1972. Little Sioux Burn: year 2. Nat. 23(3,4): 2-7. (NC)

154. Buchanan, H, W.A. Laycock, and D.A. Price
1972. Botanical and nutritive content of the summer diet of sheep on a tall-forb range in southwestern Montana. J. Animal Sci. 35(2): 423-430. (INT)

155. Driscoll, R.S.
1972. Pattern recognition of native plant communities—Manitou Colorado Test Site. In 4th Annu. Earth Resour. Program Rev. NASA-Manned Spacecraft Center - 05937; Vol. 5, Sect. 123: 1-28. (RM)

156. Driscoll, R.S., P.O. Currie, and M.J. Morris
1972. Estimates of herbaceous standing crop by microdensitometry. Photogramm. Eng. 38: 590. (Abstracts of Technical Papers.). (RM)

157. Driscoll, R.S., and R.E. Francis
1972. Multistage, multiband, and dequential imagery to identify and quantify non-forest vegetation resources. 5th Annual and Final Progress Report for the NASA Earth Resources Survey Program; Office of Space Science and Applications. 44p. (Available in Microfiche, Scientific and Technical Aerospace Reports.). (RM)

158. Driscoll, R.S., and R.E. Francis
1973. 70-mm aerial photographs for range vegetation analysis. In Abstracts, 26th Annual Meet. Soc. for Range Management, Boise, Idaho, Feb. 4-9, 1973, p.13. (RM)

159. Driscoll, R.S., and M.M. Spencer
1972. Multispectral data scanner for plant community classification. In Summaries, 8th Int. Symp. on Remote Sensing of Environment, Univ. of Mich., Ann Arbor, Oct. 2-6, 1972, p. 127-128. (RM)

160. Evans, K.E., and R.R. Kerbs
1972. Some birds of Jackson County, South Dakota. S. D. Bird Notes Vol. XXIV, No. 4, 3 p. (RM)

161. Francis, R.E., R.S. Driscoll, and J.N. Reppert
1972. Loop-frequency as related to plant cover, herbage production, and plant density. USDA For. Serv. Res. Pap. RM-94, 8 p. (RM)

162. Frischknecht, N.C., and M.F. Baker
1972. Voles can improve sagebrush rangelands. J. Range Manage. 25(6): 466-468. (INT)

163. Frischknecht, N.C., and L.E. Harris
1973. Sheep can control sagebrush on seeded rangelands, if—. Utah Sci. 34(1). (INT)

164. Gartner, F.R., and K.E. Severson
1972. Fee hunting in western South Dakota. J. Range Manage. 25(3): 234-237. (RM)

165. Grelen, H.E., L.B. Whitaker, and R.E. Lohrey
1972. Herbage response to precommercial thinning in direct-seeded slash pine. J. Range Manage. 25(6): 435-437. (SO)

166. Holmgren, R.C., and S.F. Brewster, Jr.
1972. Distribution of organic matter reserve in a desert shrub community. USDA For. Serv. Res. Pap. INT-130, 15 p. (INT)

167. Jameson, D.A.
1971. Optimum stand selection for juniper control on southwestern woodland ranges. J. Range Manage. 24(2): 94-99. (RM)

168. Lennartz, M.R., and J.W. McMinn
1973. Growth of two varieties of slash pine on prepared sites in South Florida: 10-year results. USDA For. Serv. Res. Pap. SE-103, 10 p. (SE)

169. Lewis, C.E.
1972. Chopping and webbing control saw-palmetto in south Florida. USDA For. Serv. Res. Note SE-177, 6 p. (SE)

170. Mueggler, W.F.
1972. Plant development and yield on mountain grasslands in southwestern Montana. USDA For. Serv. Res. Pap. INT-124, 20 p. (INT)

171. Patton, D.R., V.E. Scott, and E.L. Boeker
1972. Construction of an 8-mm time-lapse camera for biological research. USDA For. Serv. Res. Pap. RM-88, 8 p. (RM)

172. Pearson, H.A., J.R. Davis, and G.H. Schubert
1972. Effects of wildlife on timber and forage production in Arizona. J. Range Manage. 25(4): 250-253. (RM)

173. Pond, F.W.
1971. Chaparral: 47 years later. USDA For. Serv. Res. Pap. RM-69, 11 p. (RM)

174. Pond, F.W., and J.W. Bohning
1971. The Arizona chaparral. Ariz. Cattelog 27(10): 16, 18, 20, 22-28, 27(11): 13-16, 18-24. (RM)

175. Ratliff, R.D., and S.E. Westfall
1973. A simple stereophotographic technique for analyzing small plots. J. Range Manage. 26(2): 147-148. (PSW)

176. Reppert, J.N., and R.E. Francis
1973. Interpretation of trend in range condition from 3-step data. USDA For. Serv. Res. Pap. RM-103, 15 p. (RM)

177. Reynolds, H. G.
1972. Wildlife habitat improvement in relation to watershed management in the Southwest. Ariz. Watershed Symp. Proc. 16: 10-17. (RM)

178. Ringger, D., and F. Stearns
1972. Influence of forest openings on climate. Univ. Wis. Field Stn. Bull. 5(2): 8-12. (NC)

179. Segelquist, C.A., and R.E. Pennington
1972. Browse resources of the Ouachita National Forest in Arkansas. USDA For. Serv. Res. Note SO-140, 4 p. (SO)

180. Segelquist, C.A., H.L. Short, and R.G. Leonard
1972. Quality of some winter deer forages in the Arkansas Ozarks. J. Wildlife Manage. 36: 174-177. (SO)

181. Thilenius, J.F.
1972. Classification of deer habitat in the ponderosa pine forest of the Black Hills, South Dakota. USDA For. Serv. Res. Pap. RM-91, 28 p. (RM)

182. Thilenius, J. F.
1972. Environmental factors affecting the species composition and productivity of alpine tundra on Medicine Bow Peak and high subalpine meadows in Libby Flats. In Interim Progress Report, The Medicine Bow Project, p. 37-57, Univ. of Wyo., Rocky Mtn. For. and Range Expt. Sta., Wyo. Water Resour. Res. Inst., Laramie. (RM)

183. Thomas, J.W., R.O. Brush, and R.M. DeGraaf
1973. The National Wildlife Federation says: invite wildlife to your backyard. Nat. Wildf. 11(3): 5-16. (NE)

184. Tingey, W.M., C.D. Jorgansen, and N.C. Frischknecht
1972. Thrips of the sagebrush grass range community in west-central Utah. J. Range Manage. 25(4): 304-308. (INT)

Characteristics and values of forage and browse plants

185. Carpenter, L.H., O.C. Wallmo, and M.J. Morris
1973. Effect of woody stems on estimating herbage with a capacitance meter. J. Range Manage. 26(2): 151-152. (RM)

186. Conrad, P.W., and W.T. McDonough
1972. Growth and reproduction of red elderberry on subalpine rangeland in Utah. Northwest Sci. 46:140-148. (INT)

187. Dealy, J.E.
1972. Idaho fescue preference. West Livestock J. 50(26): 40, 42, and 44. (PNW)

188. Dietz, D.R.
1972. Nutritive value of shrubs. In Wildland Shrubs - Their Biology and Utilization. USDA For. Serv. Gen. Tech. Rep. INT-1, p.289-302. (RM)

189. Feddema, C.
1972. *Sclerocarpus uniserialis* (Compositae) in Texas and Mexico. Phytologia 23(2): 201-209. (RM)

190. Ferguson, R.B.
1972. Bitterbrush topping: Shrub response and cost factors. USDA For. Serv. Res. Pap. INT-125, 11 p. (INT)

191. Florez, A, W.T. McDonough, and L.D. Balls
1972. Western coneflower—a noxious species? J. Range Manage. 25: 403-404. (INT)

547-679 O - 74 - 5

192. Garrison, G.A.
1972. Carbohydrate reserves and response to use. In Wildland Shrubs--Their Biology and Utilization. USDA For. Serv. Gen. Tech. Rep. INT-1, p. 271-278. (PNW)

193. Geist, J.M.
1972. NS relations in northeastern Oregon upland soils—key to increased productivity. 1972 Annual Meetins Soil Sci. Amer. Oct. 29-Nov. 2, Miami Beach, Fla., p. 106. (Abstr.). (PNW)

194. Haferkamp, M.R., and P.O. Currier
1973. Effect of fertilizer on root strength of Sherman big bluegrass (*Poa ampla* Merr.). Agron. J. 65: 511-512. (RM)

195. Halls, L.K., and R. Alcaniz
1972. Growth patterns of deer-browse plants in southern forests. USDA For. Serv. Res. Pap. SO-75, 14 p. (SO)

196. Harlow, R.F., and L.K. Halls
1972. Response of yellow-poplar and dogwood seedlings to clipping. J. Wildlife Manage. 36: 1076-1080. (SO)

197. Hermann, F.J.
1972. A new variety of *Carex bicknellii* from Arkansas. SIDA 5(1): 49. (RM)

198. Hermann, F.J.
1972. Book review (Moss flora of the Pacific Northwest. By Elva Lawton. 362 p. and 195 pl. Hattori Botanical Laboratory, Miyazaki, Japan. 1971.). Bull. Torrey Bot. Club 11(2): 101-102. (RM)

199. Hermann, F.J.
1972. Sphagnum magellanicum new to the West Indies. The Bryologist 75(3): 359-360. (RM)

200. Holmgren, R.C., and S.S. Hutchings
1972. Salt desert shrub response to grazing use. In Wildland Shrubs--Their Biology and Utilization. USDA For. Serv. Gen. Tech. Rep. INT-1, p. 153-164. (INT)

201. Krueger, W.C.
1972. Evaluating animal forage preference. J. Range Manage. 25: 471-475. (INT)

202. McKell, C.M., J.P. Blaisdell, and J.R. Goodin
1972. Wildland shrubs—their biology and utilization, an international symposium, Utah State University, Logan, Utah, July 1971. USDA For. Serv. Gen. Tech. Rep. INT-1, 494 p. (INT)

203. Medin, D.E., and R.B. Ferguson
1972. Shrub establishment on game ranges in the Northwestern United States. In Wildland Shrubs--Their Biology and Utilization . USDA For. Serv. Gen. Tech. Rep. INT-1, p. 359-368. (INT)

204. Plummer, A.P.
1972. Selection. In Wildland Shrubs--Their Biology and Utilization. USDA For. Serv. Gen. Tech. Rep. INT-1, p. 121-126. (INT)

205. Ratliff, R.D., S.E. Westfall, and R.W. Roberts
1972. More California-poppy in stubble field than in old field. USDA For. Serv. Res. Note PSW-271, 4 p. (PSW)

206. Segelquist, C.A., M. Rogers, and F.D. Ward
1972. Quantity and quality of Japanese honeysuckle on Arkansas Ozark food plots. In 25th Annu. Southeast. Assoc. Game and Fish Comm. Conf. Proc. 1971: 47-53. (SO)

207. Short, H.L., R.M. Blair, and L. Burkart
1972. Factors affecting nutritive values. In Wildland Shrubs--Their Biology and Utilization. USDA For. Serv. Gen. Tech. Rep. INT-1, p. 311-318. (SO)

208. Skovlin, J.M.
1972. The influence of fire on important range grasses of East Africa. 1971 Annual Tall Timbers Fire Ecology Conf. Proc., April 22-23, Tallahassee, Fla., p. 201-217. (PNW)

209. Turner, G.T.
1972. A new approach to estimating herbage moisture content. J. Range Manage. 25(3): 229-231. (RM)

210. Ward, D.E., and S.C. Martin
1972. Tanglehead-a dual purpose grass. Arizona Cattlelog 28(8): 18-20. (RM)

211. Wolters, G.L.
1972. Production and persistence of common carpetgrass in relation to site and harvest frequency. J. Range Manage. 25(5): 360-364. (SO)

212. Wolters, G.L.
1972. Responses of southern bluestems to pine straw mulch, leachage, and ash. J. Range Manage. 25: 20-23. (OP)

Livestock and grazing management

213. Anderson, A.E., D.E. Medin, and D.C. Bowden
1972. Blood serum electrolytes in a Colorado mule deer population. J. Wildlife Dis. 8: 183-190. (INT)

214. Anderson, A.E., D.E. Medin, and D.C. Bowden
1972. Total serum protein in a population of mule deer. J. Mammal. 53: 384-487. (INT)

215. Boeker, E.L., V.E. Scott, H.G. Reynolds, and B.A. Donaldson
1972. Seasonal food habits of mule deer in southwestern New Mexico. J. Wildl. Manage. 36(1): 36-63. (RM)

216. Brander, R.B., and D.J. Books
1973. Return of the fisher. Nat. Hist. 82(1): 52-57. (NC)

217. Duffy, I.T.
1972. Landowner and stockmen gain from forest forage. Forest Farmer 32(1): 15-16. (SO)

218. Duncan, D.A., and J.R. Merton
1973. Yearlong tops seasonal grazing in extended rangeland study. Western Livestock Journal, Mountain, Plains, and Southwest Edition 51(21):32, 48; Pacific Slope Edition 51(21):32, 53. (PSW)

219. Harlow, R.F., and W.A. Guthrie
1972. Grouse crops—a habitat barometer. Virginia Wildlife, p. 10-11, Comm. of Game and Inland Fish. Richmond, Va. (SE)

220. Healy, W.M.
1972. Wild turkey brood habitat evaluation using human-imprinted poults. Available from Information Services, Northeastern Forest Experiment Station, Forest Service, USDA, Upper Darby, Pa. 19082. (NE)

221. Kundaeli, J.N., and H.G. Reynolds
1972. Desert cottontail use of natural and modified pinyon-juniper woodland. J. Range Manage. 25(2): 116-118. (RM)

222. Laycock, W.A., H. Buchanan, and W.C. Krueger
1972. Three methods for determining diet, utilization, and trampling damage on sheep ranges. J. Range Manage. 25(5): 352-356. (INT)

223. Martin, S.C.
1972. Semidesert ecosystems—Who will use them? How will we manage them? J. Range Manage. 25(4): 317-319. (RM)

224. Martin, S.C.
1972. Some effects of continuous grazing on forage production. Arizona Cattlelog 28(10): 17, 18, 23-25. (RM)

225. Martin, S.C., and D.E. Ward
1973. Salt and meal-salt help distribute cattle use on semidesert range. J. Range Manage. 26(2): 94-97. (RM)

226. Parker, H.D., Jr.
1972. Airborne infrared detection of deer. Ph. D. Thesis, Colo. State Univ., Ft. Collins, 186 p. (Available through University Microfilms, Ann Arbor, Mich.). (RM)

227. Parker, H.D., Jr., and R.S. Driscoll
1972. An experiment in deer detection by thermal scanning. J. Range Manage. 25(6): 480-481. (RM)

228. Parker, H.D., Jr., and J.C. Harlan
1972. Solar radiation affects radiant temperatures of a deer surface. USDA For. Serv. Res. Note RM-215, 4 p. (RM)

229. Pearson, H.A.
1972. Estimating cattle gains from consumption of digestible forage on ponderosa pine range. J. Range Manage. 25(1): 18-20. (RM)

230. Pearson, H.A., J.F. Mann, and D.A. Howard
1971. Timing use of cool- and warm-season grasses on pine ranges. J. Range Manage. 24(2): 162-163. (RM)

231. Pearson, H.A., Whitaker, L.B.
1972. Thrice-weekly supplementation adequate for cows on pine-bluestem range. J. Range Manage. 25(4): 315-316. (SO)

232. Pond, F.W., and H.A. Pearson
1971. Freeze branding cattle for individual identification. J. Range Manage. 24(6): 466-467. (RM)

233. Ratliff, R.D.
1972. Livestock grazing not detrimental to meadow wildflowers. USDA For. Serv. Res. Note PSW-270, 4 p. (PSW)

234. Ratliff, R.D., J.N. Reppert, and R.J. McConnen
1972. Rest-rotation grazing at Harvey valley—range health, cattle gains, cost. USDA Forest Serv. Res. Pap. PSW-77, 24 p. (PSW)

235. Reynolds, H.G.
1972. An albino Gambel quail from southern Arizona. J. Ariz. Acad. of Sci. 7(2): 46. (RM)

236. Reynolds, H.G., and F. Turkowski
1972. Reproductive variations in the round-tailed ground squirrel as related to winter rainfall. J. Mammal. 53(4): 893-898. (RM)

237. Schneeweis, J.C., K.E. K.E. Severson, and L.E. Petersen
1972. Food habits of deer in the Black Hills. Part 1. Northern Black Hills. S.D. State Univ. Agric. Exp. Sta. Bull. 606, pp. 5-18. (RM)

238. Severson, K.E., H.E. Messner, and D.R. Dietz
1972. Two-headed white-tailed deer fetus. Am. Midl. Nat. 88(2): 464-465. (RM)

239. Short, H.L.
1972. Ecological framework for deer management. J. For. 70(4): 200-203. (SO)

240. Turner, G.T.
1973. Effects of pocket gophers on the range. IN Pocket Gophers and Colorado Mountain Rangeland. Colo. State Univ. Exp. Sta. Bull. 554S, p. 51-61. (RM)

241. Watkins, R.K., and P.J. Urness
1972. Maxillary canine and supernumerary incisors in Arizona Coues white-tailed deer. Southwest. Nat. 17(2): 211-213. (RM)

242. Weeks, R.W., J.J. Cupal, and A.L. Ward
1972. Telemetry tracking of summer transplanted elk in southcentral Wyoming. Int. Telem. Conf. (Los Angeles, Calif. Oct. 1972) Proc. 1972: 238-244. (RM)

243. Weeks, R.W., A.L. Ward, and J. Cupal
1972. A telemetry system for studying elk behavior in the Rocky Mountains. 9th Annual Rocky Mtn. Bioeng. Symp. and 10th Int. ISA Biomed. Sci. Instrum. 9: 177-181. (RM)

Integrating other uses and wildlife

244. De Graaf, R.M., and J.S. Larson
1972. A technique for the observation of sex chromatin in hair roots. J. Mammal. 53(2): 368-371. (NE)

245. Marburger, R.G., R.M. Robinson, J.W. Thomas, M.J. Andregg, and K.A. Clark
1972. Antler malformation produced by leg injury in white-tailed deer. J. Wildlife Disease 8: 311-314.

246. Maxwell, R.K., L.L. Rogers, and R.B. Brander
1972. The energetics of wintering bears (*Ursus americanus*) in northeastern Minnesota. Ecol. Soc. Am. Bull. 53(2): 21(Abstr.). (NC)

247. Sanderson, H.R., and L.R. Berry
1972. A 2-acre enclosure for tree squirrel research. SE Assoc. Game and Fish Comm. 26th Ann. Conf. Proc., in press. (NE)

248. Sassaman, R.W., J.W. Barrett, and J.G. Smith
1972. Economics of thinning stagnated ponderosa pine sapling stands in the pine-grass areas of central Washington. USDA For. Serv. Res. Pap. PNW-144, 17 p. (PNW)

249. Severson, K.E., and F.R. Gartner
1972. Problems in commercial hunting systems: South Dakota and Texas compared. J. Range Manage. 25(5): 342-345. (RM)

250. Thill, R.E., K.E. Severson, and Y.A. Grechius
1972. Effects of aldrin on young pheasants under seminatural conditions. Bull. Environ. Contam. and Toxicol. 7(2/3): 188-192. (RM)

251. Thomas, J.W., and B.R. Payne
1972. The Pinchot Institute of Environmental Forestry—research underway at the Amherst, Mass. Work Unit. Tree Wardens, Arborists, and Utilities 1972 Conf. Proc., p. 27-31. (NE)

252. Wallmo, O.C., W.L. Regelin, and D.W. Reichert
1972. Forage use by mule deer relative to logging in Colorado. J. Wildlife Manage. 36(4): 1025-1033. (RM)

FOREST RECREATION RESEARCH

Environmental amenities—landscapes and open space

253. Kaplan, R.
1973. Predictors of environmental preference: Designers and "clients". In Environmental Research Design, Wolfgang F.E. Preiser (Ed.), Dowden, Hutchinson & Ross, Inc., Stroudsburg, Pa., Vol. 1: 265-274. (NC)

254. Kaplan, S.
1973. Cognitive maps, human needs and the designed environment. In Environmental Research Design, Wolfgang F.E. Preiser (Ed.), Dowden, Hutchinson & Ross, Inc., Stroudsburg, Pa., Vol. 1: 275-283. (NC)

255. Kaplan, S., R. Kaplan, and J.S. Wendt
1972. Rated preferences and complexity for natural and urban visual material. Perception and Psychophysics 12(4): 354-356. (NC)

256. Kojima, M., and J.A. Wagar
1972. Computer-generated drawings of ground form and vegetation. J. For. 70: 282-285. (PNW)

257. Litton, R.B., Jr.
1972. Aesthetic dimensions of the landscape. In Natural Environments--Studies in Theoretical and Applied Analysis, John V. Krutilla, (Ed.), p. 262-291, Publ. for Resources for the Future, Inc. Baltimore and London: Johns Hopkins Univ. Press. (PSW)

258. Thomas, J.W., and R.A. Dixon
1973. Cemetery ecology. Natur. Hist. 82(3): 60-67. (NE)

Environmental amenities—wilderness

259. Books, D.J.
1972. Little Sioux Burn: Year 2. Naturalist 23 (Special Issue 3 and 4): 2-7. (NC)

260. Elsner, G.H.
1972. Wilderness management—a computerized system for summarizing permit information. USDA For. Serv. Gen. Tech. Rep. PWS-2, 8 p. (PSW)

261. Gilbert, C.G., G.L. Peterson, and D.W. Lime
1972. Toward a model of travel behavior in the Boundary Waters Canoe Area. Environ. and Behav. 4(2): 131-157. (NC)

262. King, J.G., and A.C. Mace, Jr.
1972. Can we keep our canoe country waters clean? Volunteer (Minn. Dept. Natur. Resour.) 35(205): 57-61. (NC)

263. Lime, D.W.
1972. Large groups in the Boundary Waters Canoe Area—Their numbers, characteristics, and impact. USDA Forest Serv. Res. Note NC-142, 4 p. (NC)

264. Lucas, R.C.
1970. User concepts of wilderness and their implications for resource management. In Environmental Psychology: Man and His Physical Setting, Harold M. Proshansky, William H. Ittelson, Leanne G. Rivlin, (Eds.), Environmental Psychology Program, the City University of New York, New York, Chicago, San Francisco, Atlanta, Dallas, Montreal, Toronto, London, Sydney. p. 297-303. (NC)

265. Stankey, G.H.
1972. A strategy for the definition and management of wilderness quality. In Natural Environments--Studies in Theoretical and Applied Analysis, John V. Krutilla, (Ed.), Publ. for Resources for the Future, Inc., Baltimore and London: Johns Hopkins Univ. Press, p. 88-114. (INT)

266. Swain, A.M.
1972. A fire history of the Boundary Waters Canoe Area as recorded in Lake Sediment. Naturalist 23 (Special Issue 3 and 4): 24-31. (NC)

Managing recreational opportunities

267. Bultena, G.L., and J.C. Hendee
1972. Foresters' views of interest group positions on forest policy. J. Forest. 70(6): 337-342. (PNW)

268. Burdge, R., and J. Hendee
1972. The demand survey dilemma: Assessing the credibility of State outdoor recreation plans. Guideline 2(6): 65-68. (PNW)

269. Clark, R.N., R.L. Burgess, and J.C. Hendee
1972. The development of anti-litter behavior in a forest campground. J. Appl. Behav. Anal. 5(1): 1-5. (PNW)

270. Clark, R.N., J.C. Hendee, and R.L. Burgess
1972. The experimental control of littering. J. Environ. Educ. 4(2): 22-28. (PNW)

271. Clark, R.N., J.C. Hendee, and R.F. Washburne
1972. Litterbags: An evaluation of their use. USDA Forest Serv. Res. Note PNW-184, 5 p. (PNW)

272. Cordell, H.K., and G.A. James
1972. Visitors' preferences for certain physical characteristics of developed campsites. USDA Forest Serv. Res. Pap. SE-100, 21 p. (SE)

273. Dickerson, A.L.
1972. A report on the Youth Conservation Corps—Motivating teenagers. Nature Study (J. of the Am. Nature Study Soc.) 26(2): 3-5. (WO)

274. Hendee, J.C.
1972. Challenging the folklore of environmental education. J. Environ. Educ. 3(3): 19-23. (PNW)

275. Hutchinson, J.G., and D.W. Lime
1972. In search of campsites. Naturalist 23 (Special Issue 3 and 4): 18-23. (NC)

276. James, G.A., and A.K. Quinkert
1972. Estimating recreational use at developed observation sites. USDA Forest Serv. Res. Pap. SE-97, 6 p. (SE)

277. James, G.A., and H.T. Schreuder
1972. Estimating dispersed recreation use along trails and in general undeveloped areas with electric-eye counters: Some preliminary findings. USDA Forest Serv. Res. Note SE-181, 8 p. (SE)

278. James, G.A., and G.R. Sanford, Searcy, A.
1972. Origin of visitors to developed recreational sites on National Forests. J. Leisure Res. 4(2): 108-118. (SE)

279. Johnston, W.E., and G.H. Elsner
1972. Variability in use among ski areas: A statistical study of the California market region. J. Leisure Res. 4(1): 43-49. (PSW)

280. LaPage, W.F.
1973. Growth potential of the family camping market. USDA Forest Serv. Res. Pap. NE-252, 25 p. (NE)

281. LaPage, W.F.
1972. Understanding camping market growth. In 2nd Annu. Fam. Fed. Amer. Camping Congr. Resume "Camping Opportunities in the 70's," May 9-10, 1972, Palmer House, Chicago, IL 60605 (Fam. Camping Fed., Bradford Woods, Martinsville, IN 46151): 47-66. (NE)

282. LaPage, W.F., and D.P. Ragain
1972. Campground marketing—the impulse camper. USDA Forest Serv. Res. Note NE-150, 4 p. (NE)

283. LaPage, W.F., P.L. Cormier, and S.C. Maurice
1972. The commercial campground industry in New Hampshire—A report on a 1971 campground census. USDA Forest Serv. Res. Pap. NE-255, 41 p. (NE)

284. Lime, D.W.
1972. Behavioral research in outdoor recreation management: An example of how visitors select campgrounds. In Environment and the Social Sciences: Perspectives and Applications, J.F. Wohlwill and S.H. Carson (Eds.), Am. Psychol. Assoc., p. 198-206. (NC)

285. Lloyd, R.D., and V.L. Fischer
1972. Dispersed versus concentrated recreation as forest policy. Seventh World Forest. Congr. Proc. (Buenos Aires, Argentina) Oct. 4-18, 1972: 196(E)-1-16. (WO)

286. Magill, A.W., and A.T. Leiser
1972. Growing plants on view landscapes and recreation areas. Guideline 2(5): 57, 59-61. (PSW)

287. Moeller, G.H., and J.H. Engelken
1972. What fishermen look for in a fishing experience. J. Wildlife Manage. 36(4): 1253-1257. (NE)

288. Moeller, G.H., and D.A. Morrison
1971. Campground owners: Your business may be influenced by how you answer mail. Dep. Resour. Manage., State Univ. N.Y., Coll. For., Syracuse Univ., 7 p. (NE)

289. Potter, D.R., K.M. Sharpe, and J.C. Hendee
1973. Human behavior aspects of fish and wildlife conservation—an annotated bibliography. USDA For. Serv. Gen. Tech. Rep. PNW-4, 288 p. (PNW)

290. Potter, D.R., K.M. Sharpe, J.C. Hendee, and R.N. Clark
1972. Questionnaires for research: An annotated bibliography on design, construction, and use. USDA For. Serv. Res. Pap. PNW-140, 80 p. (PNW)

291. Shafer, E.L., Jr., P.H. Amidon, and C.W. Severinghaus
1972. A comparison of violators and nonviolators of New York's deer-hunting laws. J. Wildlife Manage. 36(3): 933-939. (NE)

292. Solomon, M.J., and E.A. Hansen
1972. Canoeist suggestions for stream management in the Manistee National Forest of Michigan. USDA Forest Serv. Res. Pap. NC-77, 10 p. (NC)

293. Tyre, G.L.
1972. The implications of cost-benefit analysis in recreation systems management. Ph.D. Dissertation, 137 p. (May be purchased from Univ. Microfilms, Univ. Mich., Ann Arbor, MI 48104). (SE)

294. Tyre, G.L., and G.R. Welch
1972. Program manual for estimating use and related statistics on developed recreation sites. USDA For. Serv. Gen. Tech. Rep. SE-1, 44 p. (SE)

295. USDA Forest Service, Northeastern Forest Experiment Station
1972. Summary of the forest recreation symposium. USDA Forest Serv. Res. Pap. NE-235, 21 p. (NE)

296. Wagar, J.A.
1972. The recording quizboard: A device for evaluating interpretive services. USDA For. Serv. Res. Pap. PNW-139, 12 p. (PNW)

FOREST INSECT AND DISEASE RESEARCH

FOREST INSECT RESEARCH

Chemicals in the forest environment

297. Look, M., and L.R. White
1972. Synthesis of 2-acyl-3-amino-1,2,4-triazoles and 2-acetyl-3-amino-1,2,4-triazoles-5-^{14}C (isotope 14). J. Agr. Food Chem. 20(4): 824. (PSW)

298. Pieper, G.R.
1972. Effect of target size and air movement on drop impingment efficiency and drop size distribution. J. Econ. Entomol. 65(3): 884-886. (PSW)

299. Roberts, R.B., R.L. Lyon, C.K. Duckles, and M. Look
1972. Influence of selected synergists on the action of five insecticides on larval western spruce budworm: absence of synergism and in vitro oxidation of Zectran. J. Econ. Entomol. 6(5): 1277-1282. (PSW)

Detection and evaluation

300. Batzer, H.O.
1973. Net effect of spruce budworm defoliation on mortality and growth of balsam fir. J. For. 71(1): 34-37. (NC)

301. Campbell, R.W., and H.T. Valentine
1972. Tree condition and mortality following defoliation by the gypsy moth. USDA Forest. Serv. Res. Pap. NE-236, 331 p. (NE)

302. Carolin, V.M., and W.K. Coulter
1972. Sampling populations of western spruce budworm and predicting defoliation on Douglas-fir in eastern Oregon. USDA Forest Serv. Res. Pap. PNW-149, 38 p. (PNW)

303. Ebel, B.H., E.P. Merkel, and R.J. Kowal
1972. Key to damage of southern forest trees by insects. For. Farmers Man. 31(7):31-36. (SE)

304. Heller, R.C.
1972. Remote sensing in forestry—Promises and problems. Proc. SAF-Natl. Meet., Hot Springs, Ark., Oct. 1-5, 1972. (PSW)

305. McKnight, M.E.
1971. Natural mortality of the western spruce budworm, *Choristoneura occidentalis*, in Colorado. USDA For. Serv. Res. Pap. RM-81, 12 p. (RM)

306. McManus, M.L.
1972. Aerobiology and airborne insects. IN Proc. Conf. on Ecological Systems Approaches to Aerobiology I. Identification of componsnet elements and their functional relationships. Manhattan, Kansas, Jan. 6-8, 1972. US/IBP Aerobiol. Program Hdbk. No. 2, Univ. Mich., Ann Arbor, Mich., p. 76-78. (NE)

307. McManus, M.L.
1973. A dispersal model for larvae of the gypsy moth *Porthetria diapar* L. IN Proc. Conf. on Ecological Systems Approaches to Aerobiology II. Development, demonstration, and evaluation of models. Natl. Ctr. for atmos. Res., Boulder, Colo., July 12-14, 1972. US/IBP Aerobiol. Program Hdbk. No. 2, Univ. Mich., Ann Arbor, Mich., p. 129-38. (NE)

308. Martignoni, M.E.
1972. A rapid method for the identification of nucleopolyhedron types. J. Invert. Pathol. 19(2): 281-283. (PNW)

309. Mata, S.A.,
1972. Accuracy of determining mountain pine beetle attacks in ponderosa pine utilizing pitch tubes, frass, and entrance holes. USDA For. Serv. Res. Note RM-222, 2 p. (RM)

310. Thornbury, J.R., and R.E. Denton
1972. Low kilovoltage radiography in the detection of parasites of the western spruce budworm, *Choristoneura occidentalis*. Annu. Entomol. Soc. Amer. 65(5): 1232-1235. (INT)

311. Weber, F.P.
1973. The use of airborne spectrometers and multispectral scanners for previsual detection of ponderosa pine trees under stress from insects and disease. IN Monitoring Forest Land from High Altitude and from Space. Final Rpt. (Available in microfiche from NASA, Wash., D.C., Sci. and Tech. Aerospace Rpts.). (PSW)

312. Weber, F.P., T.H. Waite, and R.C. Heller
Trend and spread of bark beetle infestations in the Black Hills. IN Monitoring Forest Land from High Altitude and from Space. Final Rpt. (Available in microfiche form NASA, Wash., D.C., Sci. and Tech. Aerospace Rpts.). (PSW)

313. Yates, H.O., III, and B.H. Ebel
1972. Light-trap collections and review of biologies of two species of pine-feeding Geometridae. J. Ga. Entomol. Soc. 7(4): 265-271. (SE)

Biology and understanding

314. Abrahamson, L.P., and F.I. McCracken
1971. Insect and disease pests of southern hardwoods. Proc. Symp. Southeast. Hardwoods 1971, p. 80-89. USDA For. Serv. State and Private Forestry, Atlanta, Ga. (SO)

315. Abrahamson, L.P., and L. Newsome
1972. Tree age influences trunk borer infestations in cottonwood plantations. For. Sci. 18(3): 231-232. (SO)

316. Amman, C.D.
1972. Prey consumption and development of *Thanasimus undatulus*, a predator of the mountain pine beetle. J. Environ. Entomol. 1(4): 528-530. (INT)

317. Amman, G.D.
1972. Some factors affecting oviposition behavior of the mountain pine beetle. J. Environ. Entomol. 1(6): 691-695. (INT)

318. Andrews, T.L., and R.P. Miskus
1972. Some effects of fatty acids and oils on western spruce budworm larvae and pupae. Pest. Biochem. Physiol. 2(3): 257-261. (PSW)

319. Asher, W.C.
1972. Receptor unit activity in pales weevil larvae. J. Elisha Mitchell Sci. Soc. 88: 195 Abstr. (SE)

320. Barras, S.J.
1972. Improved White's solution for surface sterilization of pupae of *Dendroctonus frontalis*. J. Econ. Entomol. 65:1504. (SO)

321. Barras, S.J., and T. Perry
1972. Fungal symbionts in the prothoracic mycangium of *Dendroctonus frontalis* (Coleopt.: Scolytidae). Z. Angew. Entomol. 71: 95-104. (SO)

322. Batzer, H.O.
1972. Aspen insects. IN Aspen Symp. Proc. USDA For. Serv. Gen. Tech. Rep. NC-1: 83-87. (NC)

323. Beckwith, R.C.
1972. Key to adult bark beetles commonly associated with white spruce stands in interior Alaska. USDA For. Serv. Res. Note PNW-189, 6p. (PNW)

324. Beckwith, R.C.
1972. Scolytid flight in white spruce stands in Alaska. Can. Entomol. 104: 1977-1983. (PNW)

325. Bennett, W.H., and H.E. Ostmark
1972. Insect pests of southern pines. USDA For. Serv., South. For. Exp. Sta., 38 p. (SO)

326. Bohart, G.E., and T.W. Koerber
1972. Insects and seed production. IN Seed Biology, T.T. Koslowski (Ed.), Vol. III, p. 1-53, Acad. Press, Inc., New York. (PSW)

327. Brewer, J.W., and R.E. Stevens
1972. Biology of the pinyon pitch nodule moth in Colorado. J. Colo.-Wyo. Acad. Sci. 7: 68-69 Abstr. (RM)

328. Carolin, V.M., Jr., and G.E. Daterman
1972. European pine shoot moth hazard. Am. Forests (Forum) 78(10): 5. (PNW)

329. Clark, E.W.
1972. The role of *Pityophthorus pulicarius* Zimmerman in tip dieback of young loblolly pine. J. Ga. Entomol. Soc. 7(2): 151-152. (SE)

330. Daterman, G.E.
1972. Effects of high temperature and vapor pressure deficit on European pine shoot moth, *Rhyacionia buoliana* (Lepidoptera: Olethreutidae), egg production and survival. Can. Entomol. 104: 1387-1396. (PNW)

331. Drooz, A.T., and V.H. Fedde
1972. Discriminate host selection by *Monodontomerus dentipes*. Environ. Entomol. 1: 522-523. (SE)

332. Drooz, A.T., and H.T. Schreuder
1972. Elm spanworm: Models of eclosion as related to temperature and relative humidity in the laboratory. Environ. Entomol. 1(5): 582-588. (SE)

333. Ebel, B.H., and H.O. Yates, III
1973. Rearing and biological observations of a southern pine cone insect, *Eucosma cocana* (Lepidoptera: Olethreutidae). Annu. Entomol. Soc. Am. 66(1): 88-92. (SE)

334. Fatzinger, C.W.
1972. Bioassay, morphology, and histology of the female sex pheromone gland of *Dioryctria abietella* (Lepidoptera: Pyralidae (Phycitinae)). Annu. Entomol. Soc. Am. 65(5): 1208-1214. (SE)

335. Fedde, G.F.
1971. The spruce aphid, *Elatobium abietinum*, observed in eastern United States. Entomol. News 82: 176. (SE)

336. Fedde, G.F.
1972. An anomalous occurrence of the spruce aphid, *Elatobium abietinum* (Walker) (Homoptera: Aphididae), in western North Carolina. Proc. Entomol. Soc. Wash. 74(4): 379-381. (SE)

337. Fellin, D.G., and P.C. Kennedy
1972. Abundance of arthropods inhabiting duff and soil after prescribed burning on forest clearcuts in northern Idaho. USDA For. Serv. Res. Note INT-162, 8 p. (INT)

338. Frye, R.H., and Schmid
1972. Polypropylene jars improve performance of bark beetle emergence cages. J. Econ. Entomol. 65: 931. (RM)

339. Furniss, M.M., and R.L. Furniss
1972. Scolytids (*Coleoptera*) on snowfields above timberline in Oregon and Washington. Can. Entomol. 104: 1471-1478. (INT)

340. Furniss, M.M., R.D. Hungerford, and E.F. Wicker
1972. Insects and Mites associated with western white pine blister rust cankers in Idaho. Can. Entomol. 104: 1713-1715. (INT)

341. Furniss, M.M., and R.G. Krebill
1972. Insects and diseases of shrubs on western big game ranges. In Wildland Shrubs -- Their Biology and Utilization USDA For. Serv. Gen. Tech. Rpt. INT-1, p. 218-227. (INT)

342. Galford, J.R.
1972. Some basic nutritional requirements of smaller European elm bark beetle larvae. J. Econ. Entomol. 65(3): 681-684. (NE)

343. Grimble, D.G., F.B. Knight, and J.C. Nord
1971. Associated insects reared from galls of *Saperda inornata* (Coleoptera: Cerambycidae) on trembling aspen in Michigan. Mich. Entomol. 4(2): 53-57. (SE)

344. Hay, C.J.
1972. Red oak borer (Coleoptera: Cerambycidae) emergence from oak in Ohio. Annu. Entomol. Soc. Am. 65(5): 1243-1244. (NE)

345. Hay, C.J.
1972. Woodpecker predation on red oak borer in black, scarlet, and northern red oak. Annu. Entomol. Soc. Am. 65(6): 1421-1423. (NE)

346. Herring, J.L.
1972. A new species of Neoborella from dwarf mistletoe in Colorado (Hemiptera: Miridae). Proc. Entomol. Soc. Wash. 74: 9-10. (RM)

347. Jennings, D.T.
1972. An overwintering aggregation of spiders (Araneae) on cottonwood in New Mexico. Entomol. News 83: 61-67. (RM)

348. Kennedy, B.H., and J.R. Galford
1972. Development of *Dendrosoter protuberans* (Hymenoptera: Braconidae) on larvae of the smaller European elm bark beetle being reared on an artificial medium. Annu. Entomol. Soc. Am. 65: 757-759. (NE)

349. Koplin, J.R.
1972. Measuring predator impact of woodpeckers on spruce beetles. J. Wildl. Manage. 36: 308-320. (RM)

350. Kowal, R.J., and B.H. Ebel
1972. Insects attacking forest trees in the South. For. Farmer Man. 31(7): 28-30. (SE)

351. Lanier, C.N., M.C. Birch, R.F. Schmitz, and M.M. Furniss
1972. Pheromones of *Ips pini* (Coleoptera: Scolytidae): Variation in response among three populations. Can. Entomol. 104: 1917-1923. (INT)

352. Lorio, P.L., Jr., V.K. Howe, and C.N. Martin
1972. Loblolly pine rooting varies with microrelief on wet sites. Ecol. 53(6): 1134-1140. (SO)

353. Lyon, R.L., C.E. Richmond, J.L. Robertson, and B.A. Lucas
1972. Rearing diapause and diapause-free western spruce budworm (*Choristoneura occidentalis* (Lepidoptera: Tortircidae)) on an artificial diet. Can. Entomol. 104: 417-426. (PSW)

354. McCambridge, W.F., and F.B. Knight
1972. Factors affecting spruce beetles during a small outbreak. Ecol. 53(5): 830-839. (RM)

355. McCambridge, W.F., and G.C. Trostle
1972. The mountain pine beetle. USDA For. Pest Leafl. 2, 6 p. (RM)

356. McKnight, M.E., and D.G. Aarhus
1973. Notes on weevils from trees and shrubs in North Dakota. USDA For. Serv. Res. Note RM-230, 4p. (RM)

357. McManus, M.L., and Smith H.R.
1972. Importance of the silk trails in the diel behavior of lake instars of the gypsy moth. Environ. Entomol. 1(6): 793-795. (NE)

358. Mason, R.R., and T.C. Tigner
1972. Forest-site relationships within an outbreak of lodgepole needle miner in central Oregon. USDA For. Serv. Res. Pap. PNW-146, 18 p. (PNW)

359. Moore, G.E.
1972. Pathogenicity of ten strains of bacteria to larvae of the southern pine beetle. J. Invert. Pathol. 20(1): 41-45. (SE)

360. Moore, G.E.
1973. Pathogenicity of tree entomogenous fungi to the southern pine beetle at various temperatures and humidities. Environ. Entomol. 2(1): 54-57. (SE)

361. Moore, G.E., and R.C. Thatcher
1973. How safe are your pines from bark beetles? For. Farmer 32(3): 12-13, 18. (SE)

362. Moser, J.C., and L.M. Roton
1972. Reproductive compatability between two widely separated populations of *Pyemotes scolyti*. Pan. Pac. Entomol. 48(2): 97-99. (SO)

363. Nelson, R.E., and C.J. Davis
1972. Black twig borer—A tree killer in Hawaii. USDA For. Serv. Res. Note PSW-274. (PSW)

364. Nord, J.C., and F.B. Knight
1971. The nomenclatural status of *Saperda inornata* Say (Coleoptera: Cerambycidae). Mich. Entomol. 4(2): 33-38. (SE)

365. Nord, J.C., and F.B. Knight
1971. The geographic variation of *Saperda inornata* Say (Coleoptera: Cerambycidae) in eastern North America. Mich. Entomol. 4(2): 39-52. (SE)

366. Nord, J.C., and F.B. Knight
1972. The distribution of *Saperda inornata* and *Oberea schaumii* (Coleoptera: Cerambycidae) within the crown of large trembling aspen, *Populus tremuloides*. Great Lakes Entomol. 5(1): 28-32. (SE)

367. Nord, J.C., and F.B. Knight
1972. The importance of *Saperda inornata* and *Oberea schaumii* (Coleoptera: Cerambycidae) galleries as infection courts of *Hypoxylon pruinatum* in trembling aspen, *Populus tremuloides*. Great Lakes Entomol. 5(3): 87-92. (SE)

368. Nord, J.C., and F.B. Knight
1972. The relationship of the abundance of *Saperda inornata* and *Oberea schaumii* (Coleoptera: Cerambycidae) in large trembling aspen, *Populus tremuloides*, to site quality. Great Lakes Entomol. 5(3): 93-97. (SE)

369. Nord, J.C., and M.L. McManus
1972. The Columbian timber beetle. USDA For. Pest Leafl. 132, 7 p. (SE)

370. Podgwaite, J.D., and R.W. Campbell
1972. The disease complex of the gypsy moth. II. Aerobic bacterial pathogens. J. Invert. Pathol. 20: 303-308. (NE)

371. Rexrode, C.O., T.W. Jones, and R.R. Jones
1972. Overwintering stages and spring emergence of *Pseudopityophthorus* spp. in Missouri and West Virginia. J. Econ. Entomol. 65: 1520. (NE)

372. Schmid, J.M.
1972. A problem in the Front Range: Pine beetles. Colo. Outdoors 21(6): 37-39. (RM)

373. Schmitz, R.F.
1972. Behavior of *Ips pini* during mating, oviposition, and larval development (Coleoptera: Scolytidae). Can. Entomol. 104: 1723-1728. (INT)

374. Smith, R.H.
1972. Xylem resin in the resistance of the Pinaceae to bark beetles. USDA For. Serv. Gen. Tech. Rpt. PSW-1, 7 p. (PSW)

375. Solomon, J.D.
1972. Biology and habits of the living beech borer in red oaks. J. Econ. Entomol. 65(5): 1307-1310. (SO)

376. Solomon, J.D., R.E. Doolittle, and M. Beroza
1972. Isolation and analysis of the carpenterworm sex pheromone. Annu. Entomol. Soc. Am. 65(5): 1058-1061. (SO)

377. Solomon, J.D., and W.W. Neel
1972. Emergence behavior and rhythms in the carpenterworm moth, *Prionoxystus robiniae*. Annu. Entomol. Soc. Am. 65(6): 1296-1299. (SO)

378. Stevens, R.E.
1971. Pine needle-sheath miner. USDA For. Pest Leafl. 65 (Rev.), 5 p. (RM)

379. Struble, G.R.
1973. Biology, ecology, and control of the lodgepole needle miner. USDA Tech. Bull. 1458, 38 p. (PSW)

380. Taylor, J.F., and E. Hodgson
1972. Phospholipid ethanolamine biosynthesis in larval fat body of the blowfly, *Phormia regina*. Insect Biochem. 2: 243-248. (SE)

381. Thomas, H.A., B.F. Swindel, W.L. Hafley, and R.J. Monroe
1972. Probabilistic phototactic behavior in a bark beetle: comment. Can. J. Zool. 50(5): 473-476. (SE)

382. Torgersen, T.R.
1972. A *Perithous* (Hymenoptera: Ichneumonidae) introduced from Europe. Great Lakes Entomol. 5: 99. (PNW)

383. Werner, R.A.
1972. Aggregation behaviour of the beetle *Ips grandicollis* in response to insect-produced attractants. J. Insect Physiol. 18: 1001-1013. (SE)

384. Werner, R.A.
1972. Response of the beetle, *Ips grandicollis*, to combinations of host- and insect-produced attractants. J. Insect Physiol. 18: 1403-1412. (SE)

385. White, J.D., L.T. Alexander, and E.W. Clark
1972. Fluctuations in the inorganic constituents of inner bark of loblolly pine with season and soil series. Can. J. Bot. 50(6): 1287-1293. (SE)

386. Wickman, B.E., and R.F. Scharpf
1972. Decay in white fir top-killed by Douglas-fir tussock moth. USDA For. Serv. Res. Pap. PNW-133, 9 p. (PNW)

387. Wilson, L.F.
1972. Life history and outbreaks of an oak leafroller, *Archips semiferanus* (Lepidoptera: Tortricidae), in Michigan. Great Lakes Entomol. 5(3): 71-77. (NC)

388. Wilson, L.F.
1972. Lions and tigers in the sand. Mich. Entomol. Soc. Entomol. Notes No. 1, 2 p. (NC)

389. Wilson, L.F.
1972. Insect galls. Mich. Entomol. Soc., Entomol. Notes No. 2, 2 p. (NC)

390. Wilson, L.F.
1972. Eastern pineshoot borer. USDA For. Pest Leafl. 134, 6 p. (NC)

391. Yates, H.O., III
1972. Bark beetles attacking Caribbean pine in northeastern Nicaragua. FAO Plant Prot. Bull. 20(2): 25-27. (SE)

392. Yates, H.O., III
1972. Identifying three pine bark beetles (Coleoptera: Scolytidae) likely to be found in northeastern Nicaragua. FAO Plant Prot. Bull. 20(5): 101-104. (SE)

Pest control techniques

393. Barger, J.H., R.A. Cuthbert, and D.G. Seegrist
1973. Statistical correlations between GLC assay and smaller European elm bark beetle bioassay. J. Econ. Entomol. 66(1): 79-81. (NE)

394. Bennett, W.H.
1971. Silvicultural techniques will help control bark beetles. In Soc. Amer. Foresters South. Region Tech. Conf. Proc., p. 289-295. (SO)

395. Breillatt, J.P., J.N. Brantley, H.M. Mazzone, M.E. Martignoni, J.E. Franklin, and N.G. Anderson
1972. Mass purification of nucleopolyhedrosis virus inclusion bodies in the K-series centrifuge. Appl. Microbiol. 23(5): 923-930. (NE)

396. Daterman, G.E., G.D. Daves, Jr., and M. Jacobson
1972. Inhibition of pheromone perception in European pine shoot moth by synthetic acetates. Environ. Entomol. 1: 382-383. (PNW)

397. DeBarr, G.L., E.P. Merkel, C.H. O'Gwynn, and M.H. Zoerb, Jr.
1972. Differences in insect infestation in slash pine seed orchards due to phorate treatment and clonal variation. For. Sci. 18(1): 56-64. (SE)

398. Denton, R.E.
1972. Establishment of *Agathis pumila*(Ratz.) for control of larch casebearer, and notes on native parasitism and predation in Idaho. USDA For. Serv. Res. Note INT-164, 6 p. (INT)

399. Fedde, G.F.
1972. Status of imported and native predators of the balsam woolly aphid on Mt. Mitchell, North Carolina. USDA For. Serv. Res. Note SE-175, 4 p. (SE)

400. Fellin, D.G.
1973. Weevils attracted to thinned lodgepole pine stands in Montana. USDA For. Serv. Res. Pap. INT-136, 20 p. (INT)

401. Fowler, R.F., I. Millers, and L.F. Wilson
1973. Aerial and ground applications of malathion control redheaded pine sawfly. J. Econ. Entomol. 66(11): 288. (NC)

402. Fowler, R.F., and L.F. Wilson
1971. Evaluation of three aldrin application methods for white grub, *Phyllophaga* spp., control (Colcoptera: Scarabaeidae). Mich. Entomol. 4: 89-91. (NC)

403. Furniss, M.M., L.N. Kline, R.F. Schmitz, and J.A. Rudinsky
1972. Tests of three pheromones to induce or disrupt aggregation of Douglas-fir beetles (Coleoptera: Scolytidae) on live trees. Annu. Entomol. Soc. Am. 65(5): 1227-1232. (INT)

404. Galford, J.R., and L.R. Schreiber
1972. Effect of feeding on twigs from benomyl-treated seedlings on the smaller European elm bark beetle. J. Econ. Entomol. 65: 1542. (NE)

405. Goolsby, R.P., J.L. Ruehle, and H.O. Yates
1972. Insects and diseases of seed orchards in the South. Ga. For. Res. Council Rep. 28: 25 p. (SE)

406. Hubbard, H.B., Jr., and F.B. Lewis
1973. A spray tower for the application of microbial aerial sprays. USDA For. Serv. Res. Pap. NE-249, 8 p. (NE)

407. Lyon, R.L., S.J. Brown, and J.L. Robertson
1972. Contact toxicity of sixteen insecticides applied to forest tent caterpillars reared on artificial diet. J. Econ. Entomol. 65(3): 928-930. (PSW)

408. McCambridge, W.F.
1972. Treatment height for mountain pine beetles in Front Range ponderosa pine. USDA For. Serv. Res. Note RM-218, 2 p. (RM)

409. Page, M., and R.L. Lyon
1973. Insecticides applied to western tussock moth reared on artificial diet: laboratory tests. J. Econ. Entomol. 66(1): 53-55. (PSW)

410. Rasmussen, L.A.
1972. Attraction of mountain pine beetles to small-diameter lodgepole pines baited with *trans*-verbenol and *Alpha*-pinene. J. Econ. Entomol. 65: 1396-1399. (INT)

411. Reardon, R.W., M.W. Statler, and W.H. McLane
1973. Rearing techniques and biology of five gypsy moth parasites. Environ. Entomol. 2: 124-27. (NE)

412. Richmond, C.E.
1972. Juvenile hormone analogs tested on larvae of western spruce budworm. J. Econ. Entomol. 65(4): 950-953. (PSW)

413. Robertson, J.L., R.L. Lyon, F.L. Shon, and N.L. Gillette
1972. Contact toxicity of 20 insecticides applied to the red-humped oakworm, *Symmerista canicosta*. J. Econ. Entomol. 65(4): 1560-1562. (PSW)

414. Robertson, J.L., R.L. Lyon, F.L. Shon, and M. Page
1973. Western blackheaded budworm: toxicity of five insecticides to two populations. J. Econ. Entomol. 66(1): 274-275. (PSW)

415. Rudinsky, J.A., M.M. Furniss, L.N. Kline, and R.F. Schmitz
1972. Attraction and repression of *Dendroctonus pseudotsugae* (Coleoptera: Scolytidae) by three synthetic pheromones in sticky traps in Oregon and Idaho. Can. Entomol. 104: 815-22. (INT)

416. Schmid, J.M.
1972. Reduced ethylene dibromide concentrations or fuel oil alone kills spruce beetles. J. Econ. Entomol. 65: 1520-1521. (RM)

417. Sonnet, P.E., and J.C. Moser
1972. Synthetic analogs of the trail pheromone of the leaf-cutting ant, *Atta texana*(Buckley). J. Agric. Food Chem. 20(6): 1191-1194. (SO)

418. Stevens, R.E.
1972. Use of silviculture in control of bark beetles (Scolytidae). VII. Natl. Cong. Mex. Entomol., Mexico City, Oct. 1970, Folia, Nos. 23-24, p. 87-88. (RM)

419. Tumlinson, J.H., J.C. Moser, R.M. Silverstein, R.G. Brownlee, and J.M. Ruth
1972. A volatile trail pheromone of the leaf-cutting ant, *Atta texana*. J. Insect Physiol. 18: 809-814. (SO)

420. Wickman, B.E., R.R. Mason, and C.G. Thompson
1973. Major outbreaks of the Douglas-fir tussock moth in Oregon and California. USDA For. Serv. Gen. Tech. Rpt. PNW-5, 18 p. (PNW)

421. Yendol, W.G., R.A. Hamlen, and F.B. Lewis
1973. Evaluation of *Bacillus thuringiensis* for gypsy moth suppression. J. Econ. Entomol. 66(1): 183-186. (NE)

Pest management strategies

422. Amman, G.D., and B.H. Baker
1972. Mountain pine beetle influence on lodgepole pine stand structure. J. For. 70(4): 204-209. (INT)

Remote sensing methods

423. Aldrich, R.C., and W.J. Greentree
1972. Forest and nonforest land classification using aircraft and space imagery. In Monitoring Forest Land from High Altitude and from Space. Final Rep. available in microfiche from NASA, Wash., D.C., Sci. and Tech. Aerospace Reps. (PSW)

424. Aldrich, R.C., J. von Mosch, and W.J. Greentree
1972. Projection-viewer for microscale aerial photography. USDA For. Serv. Res. Note PSW-277, 4 p. (PSW)

425. Dana, R.W.
1973. Digital sensitometry of color infrared film as an aid to pattern recognition studies. In Proc. 2nd Annu. Remote Sensing Earth Resources Conf., Univ. Tenn. Space Inst., Tullahoma, TN, March 26-28, 1973. (PSW)

426. Dana, R.W., and B. Marx
1972. Calibration of focal plane shutters. Final Rep. available in microfiche from NASA, Wash., DC, Sci. and Tech. Aerospace Reps. (PSW)

427. Dana, R.W., and N.X. Norick
1972. Mathematical modeling of film characteristic curves. Final Rep. available in microfiche from NASA, Wash., DC Sci. and Tech. Aerospace Reps. (PSW)

428. Heller, R.C., B. Spada, and A.M. Woll
1972. Remote sensing in resource evaluation, planning, protection, and management. In Proc. 7th World Forestry Cong., Buenos Aires, Argentina, October 4-18, 1972. (PSW)

429. Hildebrandt, G., and R.C. Heller
1973. The implication of remote sensors for forestry research and practice. In Proc. XV IUFRO Cong., Gainesville, FL, March 14-20, 1971. (PSW)

430. Norick, N.X., and M. Wilkes
1972. Classification of land use by automated procedures. In Monitoring Forest Land from High Altitude and from Space. Final rep. available in microfiche from NASA, Wash., DC, Sci. and Tech. Aerospace Reps. (PSW)

431. Weber, F.P., R.C. Aldrich, F.G. Sadowski, and F.J. Thomson
1972. Land use classification in the southeastern forest region by multispectral scanning and computerized mapping. In Monitoring Forest Land from High Altitude and from Space. Final rep. available in microfiche from NASA, Wash., DC Sci. and Tech. Aerospace Reps. (PSW)

Biological degradation

432. Carter, F.L., and R.V. Smythe
1972. Extractives of baldcypress, black walnut, and redwood and survival of the eastern subterranean termite, RETICULITERMES FLAVIPES. Annu. Entomol. Soc. Am. 65: 686-689. (SO)

433. Carter, F.L., L.A. Dinus, and R.V. Smythe
1972. Effect of wood decayed by *Lenzites trabea* on the fatty acid composition of the eastern subterranean termite, *Reticulitermes flavipes*. J. Insect Physiol. 18: 1387-1393. (SO)

434. Carter, F.L., L.A. Dinus, and R.V. Smythe
1972. Fatty acids of the eastern subterranean termite, *Reticulitermes flavipes* (Isoptera: Rhinotermitidae). Annu. Entomol. Soc. Am. 65(3): 655-658. (SO)

435. Carter, F.L., C.A. Stringer, and R.V. Smythe
1972. Survival of six colonies of *Reticulitermes flavipes* on unfavorable woods. Annu. Entomol. Soc. Am. 65(4): 984-985. (SO)

436. Mauldin, J.K., R.V. Smythe, and C.C. Baxter
1972. Cellulose catabolism and lipid synthesis by the subterranean termite, *Coptotermes formosanus*. Insect Biochem. 2: 209217. (SO)

437. Smythe, R.V.
1972. Feeding and survival at constant temperatures by normally and abnormally faunated *Reticulitermes virginicus* (Isoptera: Rhinotermitidiae). Annu. Entomol. Soc. Am. 65(3): 756-757. (SO)

438. Smythe, R.V., and J.K. Mauldin
1972. Soldier differentiation, survival, and wood consumption by normally and abnormally faunated workers of the Formosan termite, *Coptotermes formosanus*. Annu. Entomol. Soc. Am. 65(5): 1001-1004. (SO)

439. Smythe, R.V., and L.H. Williams
1972. Feeding and survival of two subterranean termite species at constant temperatures. Annu. Entomol. Soc. Am. 65(1): 226-229. (SO)

440. Solomon, J.D., L. Newsome, and W.N. Darwin
1972. Carpenterworm moths and cerambycid hardwood borers caught in light traps. J. Ga. Entomol. Soc. 7(1): 76-79. (SO)

441. Williams, L.H.
1972. Roller cages for study of wood-products insects in building crawl spaces. J. Econ. Entomol. 65(3): 613-615. (SO)

442. Williams, L.H.
1972. Anobiid beetle eggs consumed by a psocid (Psocoptera: Liposcelidae). Annu. Entomol. Soc. Am. 65(3): 533-536. (SO)

Prevention and control of wood-destroying organisms

443. Beal, R.H., and V.K. Smith
1972. Carbamate or phosphate insecticides for subterranean termite control? Pest Control 40(7): 20, 22, 43. (SO)

444. Bultman, J.D., C.R. Southwell, and R.H. Beal
1972. Termite resistance of polyvinyl chloride plastics in southern temperate and tropical environments: Final Report Phase I - Effect of Plasticizers and Insecticides. U.S. Nav. Res. Lab. Rep. 7417, 20 p. (SO)

445. Johnston, H.R., V.K. Smith, and R.H. Beal
1972. Subterranean termites, their prevention and control in buildings. USDA Home and Garden Bull. 64 (Rev.), 30 p. (SO)

446. Smith, V.K., R.H. Beal, and H.R. Johnston
1972. Twenty-seven years of termite control tests. Pest Control 40(6): 28, 42, 44. (SO)

447. Southwell, C.R., J.D. Bultman, and R.H. Beal
1972. Termite resistance of polymeric materials: Phase 2—Nontoxic polymers. U.S. Nav. Res. Lab. Rep. 7418, 22 p. (SO)

448. Williams, L.H., and H.R. Johnston
1972. Controlling wood-destroying beetles in buildings and furniture. USDA Leafl. 558, 8 p. (SO)

FOREST DISEASE RESEARCH

Air pollution

449. Bega, R.V., P.R. Miller, and J.R. McBride
1973. Oxidant air pollutant effects on a western coniferous forest ecosystem. IN Task B Rep., Air Pollut. Res. Cent., Univ. Calif., Riverside, p. A1-A36. (PSW)

450. Dochinger, L.S.
1972. Impact of air pollution on forest tree plantings. Soil Conserv. Soc. Am. Proc. 27: 134-138. (NE)

451. Dobhinger, L.S.
1972. Can trees cleanse the air of particulate pollutants? Int. Shade Tree Conf. Proc. 48: 45-48. (NE)

452. Dochinger, L.S., A.M. Townsend, D.W. Seegrist, and F.W. Bender
1972. Responses of hybrid poplar trees to sulfur dioxide fumigation. J. Air Pollut. Control Assoc. 22(5): 369-371. (NE)

453. Edinger, J.G., M.H. McCutchan, P.R. Miller, B.C. Ryan, M.J. Schroeder, and J.F. Behar
1972. Penetration and duration of oxidant air pollution in the south coast air basin of California. J. Air Pollut. Control Assoc. 22(11): 882-886. (PSW)

454. Evans, L.S., and P.R. Miller
1972. Comparative needle anatomy and relative ozone sensitivity of four pine species. Can. J. Bot. 50(5): 1067-1071. (PSW)

455. Miller, P.R.
1973. Oxidant damage to conifers at selected study sites, 1972. In Task C Rep., Air Pollut. Res. Cont., Univ. Calif., Riverside, III: 1-19. (PSW)

456. Miller, P.R., M.H. McCutchan, and H.P. Milligan
1972. Oxidant air pollution in the Central Valley, Sierra Nevada foothills, and Mineral King Valley of California. Atmos. Environ. 6: 623-633. (PSW)

457. Miller, P.R., M.H. McCutchan, and B.C. Ryan
1972. Influence of climate and topography on oxidant air pollution concentrations that damage conifer forests in southern California. In seventh Int. Symp. of Forest Fume Damage Experts, Forstliche Bundesversuchanst. alt., Wien, I: 585-607. (PSW)

458. Miller, P.R., and R.P. Milligan
1973. Oxidant air monitoring in the San Bernardino Mountains. In Task C Rep., Air Pollut. Res. Cent., Univ. Calif., Riverside, VI: 1-8. (PSW)

459. Miller, P.R., and R.M. Yoshiyama
1973. Self-ventilated chambers for identification of oxidant damage to vegetation at remote sites. Environ. Sci. and Techol. 7: 66-68. (PSW)

Detection and evaluation

460. Beaton, J.A., W.B. White, and F.H. Berry
1972. Radiography of trees and wood products. Mater. Eval. 30(10): 14a-17a. (NE)

461. Heller, R.C., and R.V. Bega
1973. Detection of forest diseases by remote sensing. J. For. 71: 18-21. (PSW)

462. Houston, D.R.
1972. Large-scale aerial color photography—a tool for studying forest tree disease (Abstract). Phytopathol. 62: 495. (NE)

463. Houston, D.R.
1972. The use of large-scale aerial color photography for assessing forest tree diseases. I. Basal canker of white pine: A case study. USDA For. Serv. Res. Pap. NE-230, 7 p. (NE)

464. Kessler, K.J., Jr., and R.W. Blank
1972. Endogone sporocarps associated with sugar maple. Mycol. 64(3): 634-638. (NC)

465. Knutson, D.M.
1972. Cylindrical rods: more efficient spore samplers. Plant Dis. Rep. 56(8): 719-720. (PNW)

466. McCain, A.H., and R.S. Smith, Jr.
1972. Quantitative assay of *Macrophomina phaseoli* from soil. Phytopathol. 62: 1098. (PSW)

467. Myers, C.A., F.G. Hawksworth, and P.C. Lightle
1972. Simulating yields of southwestern ponderosa pine stands, including effects of dwarf mistletoe. USDA For. Serv. Res. Pap. RM-87, 16 p. (RM)

468. Nicholls, T.H., and H.D. Brown
1972. How to identify *Lophodermium* and brown spot diseases on pine. USDA For. Serv., North Cent. Forest Exp. Stn., 5 p. (NC)

469. Nicholls, T.H., and D.D. Skilling
1972. *Lophodermium* needlecast disease of Scotch pine Christmas trees. Am. Christmas Tree J. 16(2): 11-13. (NC)

470. Skutt, H.R., A.L. Shigo, and R.A. Lessard
1972. Detection of discolored and decayed wood in living trees using a pulsed electric current. Can. J. For. Res. 2(1): 54-56. (NE)

471. Tattar, T.A., and A.L. Shigo
1972. Relationship between the degree of resistance to a pulsed electric current and wood in progressive stages of discoloration and decay in living trees. Phytopathol. 72: 792 (Abstr.). (NE)

472. Tattar, T.A., A.L. Shigo, and T. Chase
1972. Relationship between the degree of resistance to pulsed electric current and wood in progressive stages of discoloration and decay in living trees. Can. J. For., 2(3): 236-243. (NE)

Biology and understanding

473. Anderson, G.W.
1972. Diseases. In USDA For. Serv. Gen. Tech. Rep. NC-1, p. 74-82. (NC)

474. Anderson, G.W.
1972. Development of *Hypoxylon mammatum* cankers on artificially shaded aspen stems. For. Sci. 18: 316-318. (NC)

475. Bakshi, B.K.
1972. Gummosis in eucalypts. Indian For. 98: 647-648. (WO)

476. Bakshi, B.K., and S. Singh
1972. Susceptibility of exotic pines to *Cronartium himalayense*. Indian For. 98: 239-240. (WO)

477. Bakshi, B.K., M.A. Ram Reddy, Y.N. Puri, and S. Singh
1973. PL-480 final technical report, 1967-1972. Forest Path. Branch, Forest Res. Instit. and Colleges, Dehra Dun. 117 p. (WO)

478. Baur, P.S., C.H. Walkinshaw, R.S. Halliwell, and V.E. Scholes
1973. Morphology of *Nicotiana tabacum* cells grown in contact with lunar material. Can. J. Bot. 51(1): 151-156.

479. Berry, F.H., and J.A. Beaton
1972. Decay causes little loss in hickory. USDA For. Serv. Res. Note NE-152, 4 p. (NE)

480. Berry, F.H., and J.A. Beaton
1972. Decay in oak in the central hardwood region. USDA For. Serv. Res. Pap. NE-242, 11 p. (NE)

481. Berry, F.H., and W. Lautz
1972. Anthracnose of eastern hardwoods. USDA For. Serv., Forest Pest Leafl. 133, 6 p. (NE)

482. Bingham, R.T., and J. Cremmen
1971. A proposed international program for testing white pine blister rust resistance. Eur. J. For. Pathol. 1(2): 93-100. (INT)

483. Campbell, W.A., F.F. Hendrix, and W.M. Powell
1972. *Pythium* and nematode species implicated in root rot. Tree Plant. Notes 23(1): 5-7. (SE)

484. Dochinger, L.S., and F.W. Bender
1972. Additional hybrid poplar hosts of *Fusarium* canker in Ohio. Plant Dis. Rep. 56(6): 523-524. (NE)

485. Gilbertson, R.L., and H.H. Burdsall, Jr.
1972. *Phellinus torulosus* in North America. Mycol. 64(6): 1258-1269. (FPL)

486. Haissig, B.E., and A.L. Schipper, Jr.
1972. Easy extraction of enzymes from small amounts of woody tissue. Anal. Biochem. 48(1): 129-146. (NC)

487. Haissig, B.E., and A.L. Schipper, Jr.
1972. Quantifying enzyme activity in woody tissue. II. Enzyme protection. In Second North Am. Forest Biol. Workshop Proc., p. 37, (Abstr.). (NC)

488. Hall, R.H., P.S. Baur, and C.H. Walkinshaw
1972. Variability in oxygen consumption and cell morphology in slash pine tissue cultures. For. Sci. 18: 298-307. (SO)

489. Harvey, A.E.
1972. Influence of host dormancy and temperature on teliospore induction by *Cronartium ribicola*. For. Sci. 18: 321-323. (INT0)

490. Harvey, A.E.
1972. SOLAR, a management and education tool for research. In Proc. Workshop on Computer and Information Systems in Resources Management Decisions, R.N. Stone and K.D. Ware (Eds.), USDA For. Serv., p. 124-127. (INT)

491. Hendrix, J.W., and N.E. Martin
1972. The interaction of rust and wheat roots in mist culture. In Proc. Eur. and Mediterr. Cereal Rusts Conf., Prague, Czech., p. 137-140. (INT)

492. Hepting, G.H., and A.L. Shigo
1972. Difference in decay rate following fire between oaks in North Carolina and Maine. Plant Dis. Rep. 56(5): 406-407. (NE)

493. Hinds, T.E., and R.W. Davidson
1972. *Ceratocystis* species associated with aspen ambrosia beetle. Mycol. 64(2): 405-409. (RM)

494. Hodges, C.S., and L.C. May
1972. A root disease of pine, araucaria, and eucalyptus in Brazil caused by a new species of *Cylindrocladium*. Phytopathol. 62(8): 898-901. (SE)

495. Jensen, K.F., and L.S. Dochinger
1972. Gibberellic acid and height growth of white pine seedlings. For. Sci. 18: 196-197. (NE)

496. Jones, T.W., and W.R. Phelps
1972. Oak wilt. USDA For. Serv., Forest Pest Leafl. 29, 7 p. (NE)

497. Kais, A.G.
1972. Variation between southern and northern isolates of *Scirrhia acicola*. Phytopathol. 62(7): 768, (Abstr.). (SO)

498. Kamensky, S.A., and G.G. Erdmann
1973. *Diaporthe* outbreak in yellow birch seedlings. Plant Dis. Rep. 57(3): 230-233. (NC)

499. Kessler, K.J., Jr.
1972. Sapstreak disease of sugar maple. USDA For. Serv., Forest Pest Leafl. 128, 4p. (NC)

500. Kessler, K.J., Jr.
1972. Sapstreak disease of sugar maple found in Wisconsin for the first time. USDA For. Serv. Res. Note NC-140, 2 p. (NC)

501. Knutson, D.M.
1972. Growth of dwarf mistletoe-infected ponderosa pine seedlings. Phytopathol. 62: 769 (Abstr.). (PNW)

502. Knutson, D.M., and W.J. Toevs
1973. Dwarf mistletoe reduces root growth of ponderosa pine seedlings. For. Sci. 18(4): 323-324. (PNW)

503. Koenigs, J.W.
1972. Effects of hydrogen peroxide on cellulose and on its susceptibility to cellulase. Mater. und Organ. 7(2): 133-147. (SE)

504. Koenigs, J.W.
1972. *Poria weirii* as a possible commercial source of peroxidase. Appl. Microbiol. 23(4): 835-836. (SE)

505. Krebill, R.G.
1972. Germination of basidiospores of *Cronartium comandrae* on rocks and vegetation. Phytopathol. 62(3): 389-390. (INT)

506. Krebill, R.G.
1972. Mortality of aspen on the Gros Ventre elk winter range. USDA For. Serv. Res. Pap. INT-129, 16 p. (INT)

507. Krebill, R.G.
1972. *Pucciniastrum goeppertianum* in *Abies lasiocarpa* in the Rocky Mountain States. Am. Midl. Nat. 87(2): 570-576. (INT)

508. Kuhlman, E.G.
1972. Variation in zygospore formation among species of *Mortierella*. Mycol. 64(2): 325-341. (SE)

509. Laemmlen, F.F., and R.V. Bega
1972. Decline of ohia and koa forests in Hawaii. Phytopathol. 62: 770 (Abstr.). (PSW)

510. Laseter, J.L., G.W. Lawler, C.H. Walkinshaw, and J.D. Weete
1973. Fatty acids of *Pinus elliottii* tissues. Phytochem. 12: 817-821. (SO)

511. Laseter, J.L., J.D. Weete, and C.H. Walkinshaw
1973. Volatile terpenoids from aeciospores of *Cronartium fusiforme*. Phytochem. 12: 387-390. (SO)

512. Li, C.Y., K.C. Lu, J.M. Trappe, and W.B. Bollen
1972. Nitrate reducing capacity of roots and nodules of *Alnus rubra* and roots of *Pseudotsuga menziesii*. Plant and Soil 37(2): 409-414. (PNW)

513. McCracken, F.I.
1972. Sporulation of *Pleurotus ostreatus*0. Can. J. Bot. 50: 2111-2115. (SO)

514. McDonald, G.I., and R.J. Hoff
1972. Racial variation of *Cronartium ribicola* on *Pinus monticola*. Phytopathol. 62: 777. (INT)

515. McWain, P., and G.F. Gregory
1972. A neutral mannan from *Ceratocystis fagacearum* culture filtrate. Phytochem. 11: 2609-2612. (NE)

516. Nicholls, T.H., and G.W. Hudler
1972. Red pine—a new host for brown spot (*Scirrhia acicola*). Plant Dis. Rep. 56(8): 712-713. (NC)

517. Nicholls, T.H., D.D. Skilling, and G.W. Hudler
1973. *Scirrhia acicola* in Scotch pine Christmas tree plantations. Plant Dis. Rep. 57(1): 55-59. (NC)

518. Parker, J.
1972. Protoplasmic resistance to water deficits. In Water Deficits and Plant Growth, T.T. Kozlowski, (ed.), Vol. III, p. 125-176. Acad. Press, N.Y. (NE)

519. Parker, J.
1972. Spatial arrangements of some cryoprotective compounds in ice lattices. Cryobiol. 9: 247-250. (NE)

520. Parmeter, J.R., Jr., and R.F. Scharpf
1972. Spread of dwarf mistletoe from discrete seed sources into young stands of ponderosa and Jeffrey pines. USDA For. Serv. Res. Note PSW-269, 5 p. (PSW)

521. Peterson, G.W.
1973. Dispersal of aeciospores of *Peridermium harknessii* in central Nebraska. Phytopathol. 63(1): 170-172. (RM)

522. Peterson, G.W.
1973. Infection of *Juniperus virginiana* and *J. scopulorum* by *Phomopsis juniperovora*. Phytopathol. 63(2): 246-251. (RM)

523. Peterson, R.S.
1972. On *Coleosporium crowellii* (Uredinales). Plant Dis. Rep. 56: 474-475. (RM)

524. Peterson, R.S.
1972. Pine limb rust fungi in Mexico. Plant Dis. Rep. 56(10): 896-898. (INT)

525. Powers, H.R., Jr.
1972. Testing for pathogenic variability within *Cronartium fusiforme*0 and *C. quercuum*. In USDA For. Serv. Misc. Publ. 1221, p. 505-511. (SE)

526. Quraishi, M.A., and N. Iqbal
1972. New records of lichens in West Pakistan. Pakistan J. For. 22: 59-64. (WO)

527. Quraishi, M.A., and T. Mahmood
1971. Disease of *Juniperus polycarpos* C. Koch. Pakistan J. For. 21: 391-400. (WO)

528. Reynolds, P.E., W.H. Smith, and K.F. Jensen
1972. Effect of constant and fluctuating temperatures on the *in vitro* growth of *Ceratocystis* species. Trans. Br. Mycol. Soc. 59(1): 1-9. (NE)

529. Reynolds, P.E., W.H. Smith, and K.F. Jensen
1972. Influence of constant and fluctuating temperatures on the growth of *Macrophomina phaseoli*. Trans. Br. Mycol. Soc. 58(3): 512-515. (NE)

530. Rickman, J.T., A.E. Harvey, and C.G. Shaw
1972. SOLAR: An on-line information retrieval system for plant pathology. Wash. Agr. Expt. Stn. Bull. 758, 8 p. (INT)

531. Rier, J.P., and A.L. Shigo
1972. Some changes in red maple, *Acer rubrum*, tissues within 34 days after wounding in July. Can. J. Bot. 50(8): 1783-1784. (NE)

532. Riffle, J.W.
1972. Histopathology of *Pinus ponderosa* ectomycorrhizae infected with a *Meloidogyne* species. Phytopathol. 62: 785 (Abstr.). (RM)

533. Ross, E.W., and D.H. Marx
1972. Susceptibility of sand pine to *Phytophthora cinnamoni*. Phytopathol. 62(10): 1197-1200. (SE)

534. Rowan, S.J.
1972. Hardwood seedling production trends in southern nurseries 1965-1970. J. For. 70(3): 162-163. (SE)

535. Ruehle, J.L.
1972. Nematodes and forest trees. In Economic Nematology, J.M. Webster (ed.), Academic Press Inc. (LONDON), p. 312-334. (SE)

536. Ruehle, J.L.
1972. Response of sand pine to parasitism by lance nematode. Plant Dis. Rep. 56(8): 691-692. (SE)

537. Ruehle, J.L.
1973. Influence of plant-parasitic nematodes on longleaf pine seedlings. J. Nematol. 5(1): 7-9. (SE)

538. Scharpf, R.F.
1972. Light affects penetration and infection of pines by dwarf mistletoe. Phytopathol. 62(11): 1271-1273. (PSW)

539. Schipper, A.L., Jr.
1972. Changes in glucose-6 phosphate dehydrogenase, glyceraldehyde-3 phosphate dehydrogenase, and peroxidase activity in *Populus tremuloides* after inoculation with *Hypoxylon mammatum* or wounding. Phytopathol. 62(7): 787-788. (NC)

540. Schipper, A.L., Jr., and B.E. Haissig
1972. Quantifying enzyme activity in woody tissue. I. Enzyme extraction. In Second North Am. Forest Biol. Workshop Proc., p. 36 (Abstr.). (NC)

541. Seidler, R.J., P.E. Aho, P.N. Raju, and H.J. Evans
1972. Nitrogen fixation by bacterial isolates from decay in living white fir trees (*Abies concolor* (Gord. and Blend.) Lindl.). J. Gen. Microbiol. 73: 413-416. (PNW)

542. Seliskar, C.W., C.L. Wilson, and C.E. Bourne
1973. Mycoplasmalike bodies found in phloem of black locust affected with witches-broom. Phytopathol. 63: 30-34. (NE)

543. Sharon, E.M.
1972. Wounding induces a change in metabolic activity in cells of *Acer saccharum*. In Second North Am. Forest Biol. Workshop Proc., p. 15-16. (NE)

544. Shea, K.R.
1971. Disease and insect activity in relation to intensive culture of forests. In Proc. XV Int. Union Forest Res. Organ. (IUFRO) Congr., Gainesville, FL, March 1971, p. 109-118. (PNW)

545. Shigo, A.L.
1972. In the beginning. South. Lumberman 225: 168-169. (NE)

546. Shigo, A.L.
1972. Ring and ray shakes associated with wounds in trees. Holzforsch. 26(2): 60-62. (NE)

547. Shigo, A.L.
1972. Successions of microorganisms and patterns of discoloration and decay after wounding in red oak and white oak. Phytopathol. 62(2): 256-259. (NE)

548. Shigo, A.L.
1972. The beech bark disease today in the Northeastern U.S. J. For. 70(5): 286-289. (NE)

549. Skilling, D.D.
1972. Epidemiology of *Scleroderris lagerbergii*. Eur. J. For. Pathol. 2(1): 16-21. (NC)

550. Skilling, D.D., and T.H. Nicholls
1972. Brown spot needle disease in Scotch pine plantations. Am. Christmas Tree J. 16(2): 7-9. (NC)

551. Skilling, D.D., and J.T. O'Brien
1972. *Scleroderris* canker of northern conifers. USDA For. Serv., Forest Pest Leafl. 130, 4 p. (NC)

552. Smith, A.H., and J.M. Trappe
1972. The higher fungi of Oregon's Cascade Head Experimental Forest and vicinity. I. The genus *Phaeocollybia* (Agaricales) and notes and descriptions of other species in the Agaricales. Mycol. 64: 1138-1153. (PNW)

553. Smith, R.S., Jr.
1973. *Sirococcus* tip dieback of *Pinus* spp. in California forest nurseries. Plant Dis. Rep. 57(10): 69-73, January. (PSW)

554. Snow, G.A., A.G. Kais, and R.J. Dinus
1972. Further evidence of geographic variations in *Cronartium fusiforme*. Phytopathol. 62: 790 (Abstr.). (SO)

555. Spotts, R.A., J. Altman, and J.M. Staley
1972. Soil salinity related to ponderosa pine tipburn. Phytopathol. 62(7): 705-708. (RM)

556. Staley, J.M., and H.H. Bynum
1972. A new *Lophodermella* on *Pinus ponderosa* and *P. attenuata*. Mycol. 64(4): 722-726. (RM)

557. Trappe, J.M.
1972. Parasitism of *Helvelia lacunosa* by *Clitocybe sclerotoidea*. Mycol. 64(6): 1337-1340. (PNW)

558. Trappe, J.M., and J.W. Gerdemann
1972. *Endogone flammicorona* sp.nov., a distinctive segregate from *Endogone lactiflua*. Trans. Br. Mycol. Soc. 59(3): 403-407. (PNW)

559. Trappe, J.M., and J.W. Kimbrough
1972. *Elaphomyces viridiseptum*, a new species from Florida. Mycol. 64(3): 646-649. (PNW)

560. Walkinshaw, C.H., P.H. Johnson, and S. Venketeswaran
1972. Elemental abundances in callus tissues of carrot, pine, rice, soybean, and tobacco. In Vitro 7: 391-396. (SO)

561. Walkinshaw, C.H., S. Venketeswaran, P.S. Baur, R.H. Hall, T.E. Croley, J.D. Weete, V.E. Scholes, and R.H. Halliwell
1973. Effect of lunar materials on plant tissue cultures. Space Life Sci. 4: 78-89. (SO)

562. Wargo, P.M.
1972. Defoliation-induced chemical changes in sugar maple roots stimulate growth of *Armillaria mellea*. Phytopathol. 62(11): 1278-1283. (NE)

563. Wargo, P.M., and D.R. Houston
1973. Infection of defoliated sugar maple trees by *Armillaria mellea*. Phytopathol. 63: 209 (Abstr.). (NE)

564. Wargo, P.M., J. Parker, and D.R. Houston
1972. Starch content in roots of defoliated sugar maple. For. Sci. 18(3): 203-204. (NE)

565. Weete, J.D., C.H. Walkinshaw, and J.L. Laseter
1973. Response of tobacco tissue cultures growing in contact with lunar fines. Space Life Sci. 4: 90-96. (SO)

566. Welch, B.L., and N.E. Martin
1972. Quantitative analysis of sugars by densitometric inspection of thin layer chromatograms: Analysis of method. J. Chromatogr. 72: 359-364. (INT)

567. Young, J.L., I. Ho, and J.M. Trappe
1972. Endomycorrhizal invasion and effect on free amino acids content of corn roots. Agron. (Abstr.) 1972: 102. (PNW)

568. Zak, B.
1972. Flotation of excised root and fungal cultures on liquid media. Can. J. Microbiol. 18(4): 536-538. (PNW)

569. Zaman, M.B., S.M. Jamal, and A.S. Khan
1972. List of fungi on forest trees in West Pakistan. Pakistan Forest Instit., Biol. Sci. Res. Div., Bull. No. 4, 32 p. (WO)

Pest control techniques

570. Bingham, R.T.
1972. Artificial inoculation of large numbers of *Pinus monticola* seedlings with *Cronartium ribicola*. IN USDA For. Serv. Misc. Publ. 1221, p. 357-372. (INT)

571. Bingham, R.T., R.J. Hoff, and G.I.(Eds.) McDonald
1972. Biology of rust resistance in forest trees. USDA For. Serv. Misc. Publ. 1221, 681 p. (INT)

572. De Groot, R.C.
1972. Growth of wood-inhabiting fungi in saturated atmospheres of monoterpenoids. Mycol. 64(4): 863-870. (SO)

573. Dwinell, L.D.
1972. An inoculation system for *Cronartium fusiforme*. IN USDA For. Serv. Misc. Publ. No. 1221, p. 327-330. (SE)

574. Gregory, G.F., P. McWain, and T.W. Jones
1972. Effectiveness of benomyl as a protectant and therapeutant for oak wilt disease in red oak seedlings. Phytopathol. 62:761. (NE)

575. Hawksworth, F.G.
1972. Biological control of the mistletoes. IN Biological Control of Forest Diseases, V.J. Nordin (compiler), FROM XV Congr. Int. Union Forestry Res. Organ., Gainesville, Fla., Mar. 1971, Sect. 24, Subj. Group 2, Can. For Serv., Ottawa, p. 83-92. (RM)

576. Hoff, R.J., and G.I. McDonald
1972. Resistance of *Pinus armandii* to *Cronartium ribicola*. Can. J. For. Res. 2(3): 303-307. (INT)

577. Hoff, R.J., and G.I. McDonald
1972. Stem rusts of conifers and the balance of nature. IN USDA For. Serv. Misc. Publ. 1221, p. 525-536. (INT)

578. Jewell, F.F., and C.A. Snow
1972. Anatomical resistance to gall-rust infection in slash pine. Plant Dis. Rep. 56(6): 531-534. (SO)

579. Kuhlman, E.G.
1972. Susceptibility of loblolly and slash pine progeny to *Fomes annosus*. USDA For. Serv. Res. Note SE-176, 7 p. (SE)

580. Leaphart, C.D., and M.W. Foiles
1972. Effects of removing pole-blighted western white pine trees on growth and development of a mixed conifer stand. USDA For. Serv. Res. Note INT-161, 6 p. (INT)

581. Leaphart, C.D., R.D. Hungerford, and H.E. Johnson
1972. Stem deformities in young trees caused by snowpack and its movement. USDA For. Serv. Rep. Note INT-158, 10 p. (INT)

582. Lightle, P.C., and F.G. Hawksworth
1973. Control of dwarf mistletoe in a heavily used ponderosa pine recreation forest: Grand Canyon, Arizona. USDA For. Serv. Res. Pap. RM-106, 22 p. (RM)

583. McCluskey, R.A.
1972. Computerized mapping of blister rust epidemics in nursery seed beds. IN USDA For. Serv. Misc. Publ. 1221, p. 393-395. (INT)

584. McDonald, G.I., and R.J. Hoff
1971. Resistance of *Cronartium ribicola* in *Pinus monticola*: Genetic control of needle-spots-only resistance factors. Can. J. For. Res. 1(4): 197–202. (INT)

585. McWain, P., and G.F. Gregory
1973. A benomyl-derived fungitoxicant for tree wilt disease control. USDA For. Serv. Res. Note NE–162, 3p. (NE)

586. Marx, D.H.
1972. Biological control of root diseases in forest tree nurseries with emphasis on the role of mycorrhizae. IN Biological Control of Forest Diseases, V.J. Nordin (compiler), From: XV Congr. IUFRO, Gainesville, Fla., Mar. 1971, Sect. 24, Subj. Group 2, Can. For. Serv., Ottawa, p. 55–65. (SE)

587. Marx, D.H.
1972. Ectomycorrhizae as biological deterrents to pathogenic root infections. Annu. Rev. Phytopathol. 10: 429–454. (SE)

588. Marx, D.H.
1973. Growth of ectomycorrhizal and nonmycorrhizal shortleaf pine seedlings in soil with *Phytophthora cinnamomi*. Phytopathol. 63: 18–23. (SE)

589. Matthews, F.R., and S.J. Rowan
1972. An improved method for large-scale inoculations of pine and oak with *Cronartium fusiforme*. Plant Dis. Rep. 56(11): 931–934. (SE)

590. Miller, T.
1972. Fusiform rust in planted slash pines: Influence of site preparation and spacing. For. Sci. 18(1): 70–75. (SE)

591. Peterson, G.W.
1972. Chemical control of Phomopsis blight of junipers: A search for new methods. Tree Plant. Notes 23: 3–4. (RM)

592. Ross, E.W.
1973. *Fomes annosus* in the southereastern United States: Relation of environmental and biotic factors to stump colonization and losses in the residual stand. USDA For. Serv. Tech. Bull. 1459, 26 p. (SE)

593. Rowan, S.J.
1972. Control of Arizona cypress blight in nursery plantings. Plant Dis. Rep. 56(6): 558–560. (SE)

594. Rowan, S.J.
1972. Selected systemic fungicides provide little control of fusiform rust of loblolly pine in forest tree nurseries. Plant Dis. Rep. 56(7): 628–630. (SE)

595. Rowan, S.J., T.H. Filer, and W.R. Phelps
1972. Nursery diseases of southern hardwoods. USDA For. Serv., Forest Pest Leafl. 137, 7 p. (SE)

596. Shea, K.R., and J.L. Stewart
1972. Hemlock dwarf mistletoe. USDA Forest Serv. For. Pest Leafl. 135, 6 p. (PNW)

597. Shigo, A.L.
1971. Wound dressing research on red maples. Proc. 47th Int. Shade Tree Conf., p. 97a–98a. (NE)

598. Shigo, A.L., and R.S. Walters
1972. Decay associated with paraformaldehyde treated tapholes in sugar maple. Phytopathol. 62: 789 (Abstr.). (NE)

599. Shigo, A.L., and C.L. Wilson
1972. Discoloration associated with wounds one year after application of wound dressings. Arborist's News 37(11): 121–124. (NE)

600. Smith, R.S., Jr., A.H. McCain, and M.D. Srago
1973. Control of *Botrytis* storage rot of giant *Sequoia* seedlings. Plant Dis. Rep. 57(1): 67–69. (PSW)

601. Tarrant, R.F., and J.M. Trappe
1972. The role of *Alnus* in improving the forest environment. Plant and Soil, Spec. Vol. 1971: 335–348. (PNW)

602. Trappe, J.M.
1972. Biological control - Forest diseases. Proc. Annu. Meet. West. Forest Pest Comm., 1971: 16–19. (PNW)

603. Trappe, J.M.
1972. Regulation of soil organisms by red alder: a potential biological system for control of *Poria weirii*. In Managing Young Forests in the Douglas-Fir Region, A.B. Berg (compiler & editor), Symposium held June 15–18, 1970, Vol. 3, p. 35–51. Oregon State University, School of Forestry, Coravallis, Oregon. (PNW)

Pest management strategies

604. Paine, L.A.
1973. Administrative goals and safety standards for hazard control on forested recreation sites. USDA For. Serv. Res. Pap. PSW–88, 13 p. (PSW)

605. Powers, H.R.
1972. Diseases of shade and forest trees of the South. For. Farmer Man. 31(1): 37–39. (SE)

606. Roth, L.E., H.H. Bynum, and E.E. Nelson
1972. *Phytophthora* root rot of Port-Orford-cedar. USDA For. Serv., Forest Pest Leafl. 131, 7 p. (PNW)

Biological degradation

607. Burdsall, H.H.
1971. Notes on some lignicolous basidiomycetes of the southeastern United States. Elisha Mitchell Sci. Soc. 87(4): 239–245. (FPL)

608. De Groot, R.C.
1971. Interactions between wood decay fungi and *Streptomyces* species. Torrey Bot. Club Bull. 98(6): 336–339. (SO)

609. De Groot, R.C.
1972. Changes in sapwood of longleaf pine logs stored under water sprays for 4 months (Abstract). Phytopathol. 62(7): 753. (SO)

610. Kirk, T.K., and W.E. Moore
1972. Removing lignin from wood with white-rot fungi and digestibility of resulting wood. Wood and Fiber 4(2): 72–79. (FPL)

611. Larsen, M.J.
1972. Notes on Tomentelloid fungi IV. *Tomentella subalpina*. Mycol. 64(2): 443–446. (FPL)

612. Lombard, F.F., R.W. Davidson, and R.L. Gilbertson
1972. Studies of two species of *Phellinus* in western North America. Mycopathol. et Mycol. Appl. 46(4): 351–365. (FPL)

613. Lorenz, L.F., and L.R. Gjovik
Analyzing creosote by gas chromatography: Relationship to creosote specifications. Am. Wood-Preserv. Assoc. Proc., p. 8. (FPL)

Prevention and control of wood-destroying organisms

614. Amburgey, T.L.
1972. Preventing wood decay. Pest Control 40: 19-20, 22, 39. (SO)

615. De Groot, R.C.
1972. A practical look at wood decay. Econ. Bot. 26(1): 85-89. (SO)

616. Highley, T.L.
1973. Effect of alkaline treatment on decay resistance of wood. For. Prod. J. 23: 47-51. (FPL)

617. Highley, T.L.
Source of decay resistance in sodium hydroxide and ammonia-treated wood. Phytopathol. 63: 57-61. (FPL)

618. Springer, E.L., and W.E. Eslyn
1972. Process for controlling pulp chip deterioration with Kraft green liquor. U.S. Patent No. 3,646,196. (FPL)

547-679 O - 74 - 6

FOREST FIRE AND ATMOSPHERIC SCIENCES RESEARCH

Air pollution

619. Edinger, J.G., M.H. McCutchan, P.R. Miller, B.C. Ryan, M.J. Schroeder, and J.V. Behar
1972. Penetration and duration of oxidant air pollution in the south coast air basin of California. J. Air Pollut. Control Assoc. 22(11): 882-886. (PSW)

620. Miller, P.R., M.H. McCutchan, and H.P. Milligan
1972. Oxidant air pollution in the Central Valley Sierra Nevada foothills, and Mineral King Valley of California. Atmosph. Environ. 6:623-633, Pergamon Press. (PSW)

621. Miller, P.R., M.H. McCutchan, and B.C. Ryan
1972. Influence of climate and topography on oxidant air pollution concentrations that damage conifer forests in southern California. In Effects of Air Pollutants on Forest Trees, VII Int. Symp. of Forest Fume Damage Experts. Forstliche Bundesversuchanstalt, Wien, 1:585-608. (PSW)

Fire prevention, hazard reduction, and prescribed burning

622. Anderson, H.E.
1972. The consideration of forest fuels in fire planning. IN Symp. Planning for Fire Management, Southwestern Interagency Fire Counc. (SWIECO) Proc., p. 132-149. (INT)

623. Berg, A.R., and T.R. Plumb
1972. Bud activation for regrowth. IN Wildland shrubs - Their Biology and Utilization, USDA Forest Serv. Gen. Tech. Rep. INT-1, p. 279-286. (INT)

624. Brackebusch, A.P.
1972. Fire prescriptions for fuels and smoke management. IN Symp. Planning for Fire Management, Southwestern Interagency Fire Couc. Proc., p. 160-164. (INT)

625. Cooper, R.W.
1972. Smoke management. South. State Foresters' Annu. Meet. Proc., p. 46-48, November. (SE)

626. Cooper, R.W.
1972. Smoke and prescribed fire. R8/R9 Fire Manage. Officer Workshop Proc., 8 p., November. (SE)

627. Cooper, R.W.
1972. Status of research on air quality and forest burning. Eleventh Auburn Forest. Forum Proc. 1972: 6-8. (SE)

628. Cooper, R.W.
1972. Prescribed burning - why it is a vital forest management tool. For. Farmer Man. 31(7): 18-19 March. (SE)

629. Cooper, R.W.
1973. Fire doesn't have to smoke. Prog. Farmer 88(2): 79. (SE)

630. Dell, J.D.
1972. Preattack guide—area planning and development for forest fire protection in the Pacific Northwest Region. USDA For. Serv., PNW Reg., 67 p. (PNW)

631. Doolittle, M.L.
1972. The dimensions of man-caused forest fore risk: A systematic assessment. Ph. D. Diss., Univ. Wash., 114 p. (PNW)

632. Folkman, W.S.
1972. Children-caused fires - How to prevent them. Proc. 7th Annu. Meet. Middle Atlantic Interstate For. Fire Port. Compact, Chevy Chase, Md., 1972: 27-29. (PSW)

633. Folkman, W.S.
1972. Studying the people who cause forest fires. In Social Behavior, Natural Resources, and the Environment, p. 44-64, Harper and Row, New York. (PSW)

634. Folkman, W.S., and J. Taylor
1972. Fire prevention in California's Riverside County. Headstart Project - an evaluation. USDA For. Serv. Res. Pap. PSW-79, 29p. (PSW)

635. Green, L.R., and G.W. Benedict
1971. Herbicides to prevent forest fires. Proc. Washington State Weed Conf., Yakima, Wash., 1971: 73-79. (PSW)

636. Hough, W.A.
1973. Prescribed burning in South surveyed, analyzed. Fire Control Notes 34(1): 4-5. (SE)

637. Hough, W.A., and J.C. Turner
1972. Open burning for forestry and agricultural purposes in Georgia during 1971. Ga. For. Res. Pap. 72, 4p. (SE)

638. Howard, J.O., and F.R. Ward
1972. Measurement of logging residue—alternative applications of the line intersect method. USDA For. Serv. Res. Note PNW-183, 8 p. (PNW)

639. Main, W.A., and P.J. Roussopoulos
1972. Ignition of prescribed burns more reliable: electrical igniter test circuits. Fire Control Notes 33(3): 14-15. (NC)

640. Mobley, H.E.
1972. Tame fire can aid our forests. Macon Telegraph and News, Macon, Ga., October 24. (SE)

641. Nord, E.C., and C.M. Countryman
1972. Fire relations. IN Wildland Shrubs--Their Biology and Utilization, USDA For. Serv. Gen. Tech. Rep. INT-1, p. 88-97. (INT)

642. Phillips, C.B., L.E. Gunter, G.E. McClellan, and E.C. Nord
1972. Creeping sage—a slow burning plant useful for fire hazard reduction. California Fire Control Notes 26, 8 p. (PSW)

643. Plumb, T.R.
1973. Control of scrub oak regrowth with a broadcast application of herbicide pellets. Res. Prog. Rep. Western Soc. of Weed Science, Spokane, Wash., March 13-15. (INT)

644. Reynolds, R.R.J.
1972. Do forest protection agencies help start forest fires? J. Forest 70(10): 604-606. (SO)

645. Roussopoulos, P.J., and V.J. Johnson
1973. Estimating slash fuel loading for several Lake States tree species. USDA For. Serv. Res. Pap. NC-88, 8 p. (NC)

646. Sackett, S.S., and R.W. Cooper
1972. The role of fire in a managed forest. Planning for Fire Manage. Symp. Proc. 1971: 74-82. (SE)

647. Sando, R.W., and C.H. Wick
1972. A method of evaluating crown fuels in forest stands. USDA For. Serv. Res. Pap. NC-84, 10 p. (NC)

648. Williams, D.T.
1972. Smoke at Palm Beach during the 1971 Everglades wildfires. IN Manual of Fire-Danger and Fire-Weather Seminar, Macon, Ga., December, 9 p. (SE)

Fire management methods and systems

649. Barney, R.J.
1972. A 31-day battery-operated recording weather station. USDA For. Serv. Res. Note PNW-185, 7 p. (PNW)

650. Barney, R.J.
1972. An inexpensive meteorological radiation shield for thermistors and thermocouples. USDA For. Serv. Res. Note PNW-190, 7 p. (PNW)

651. Britton, C.M., C.M. Countryman, H.A. Wright, and A.G. Walvekar
1973. The effect of humidity, air temperature, and wind speed on fine fuel moisture content. Fire Tech. 9(1): 46-55. (PSW)

652. Fischer, W.C., and C.E. Hardy
1972. Fire-weather observer's handbook. USDA For. Serv., Intermountain Forest and Range Exp. Stn., 152 p. (INT)

653. Fosberg, M.A.
1972. Theory of precipitation effects on dead cylindrical fuels. For. Sci. 18(2): 98-108. (RM)

654. Furman, R.W., and R.S. Helfman
1973. A computer program for processing historic fire weather data for the National Fire-Danger Rating System. USDA For. Serv. Res. Note RM-234, 12 p. (RM)

655. George, C.W., and A.D. Blakely
1972. Effects of ammonium sulfate and ammonium phosphate on flammability. USDA For. Serv. Res. Pap. INT-121, 26 p. (INT)

656. Johansen, R.W., and H.K. Mikell
1972. Dehaviland Beaver succeeds as land-based air-tanker. Fire Control Notes 33(4): 3-4. (SE)

657. McCammon, B.P., and D.A. Rainey
1973. Fire-danger index generation. Keyboard 5(1) 1973: 17-18. (RM)

658. Mees, R.M.
1973. Three dimensional computer graphs. Fire Control Notes 34(3). (PSW)

659. Noste, N.V., and R.M. Percival
1972. Alaska's extra ace: water dropping. Fire Control Notes 33(2): 9. (PNW)

660. Percival, R.M., and N.V. Noste
1972. Helicopters and helibuckets used to control interior Alaska wildfires. Fire Control Notes 34(1): 16-18. (PNW)

661. USDA Forest Service
1973. The CL-215: Summary of its Performance as an air tanker. 57 p. Wash., D.C. (WO)

662. Ward, F.R., and J.D. Dell
1972. Hydraulic seeder sprays retardants. Fire Control Notes 33(4): 7-8. (PNW)

Forest fire science

663. Blackmarr, W.H.
1972. Moisture content influences ignitability of slash pine litter. USDA For. Serv. Res. Note SE-173, 7 p. (SE)

664. Broido, A.
1973. Flammable-whatever that means. Chem. Tech. 3(1): 14-17. (PSW)

665. Clements, H.B., and A. Alkidas
1973. Combustion of wood in methanol flames. Comb. Sci. and Tech. 7: 13-18. (SE)

666. Countryman, C.M.
1971. Fire whirls—why, when, and where. USDA For. Serv., PSW, 11 p. (PSW)

667. Countryman, C.M.
1972. The fire environment concept. USDA For. Serv., PSW, 12 p. (PSW)

668. Frandsen, W.H., and R.C. Rothermel
1972. Measuring the energy-release rate of a spreading fire. Combustion and Flame 19:17-24. (INT)

669. Haines, D.A., W.A. Main, and J.S. Crosby
1973. Forest fires in Missouri. USDA For. Serv. Res. Pap. NC-87, 18 p. (NC)

670. Halpern, Y., R. Riffer, and A. Broido
1973. Levoglucosenone (1,6-Anhydro-3,4-dideoxy-delta 3-beta-D-Pyranosen-2-one). A major product of the acid-catalyzed pyrolysis of cellulose and related carbohydrates. J. Org. Chem. 38:204-209. (PSW)

671. Hough, W.A.
1973. Fuels and weather influence fires in sand pine forest. USDA For. Serv. Res. Pap. SE-106, 11 p. (SE)

672. Lindenmuth, A.W., Jr., and J.R. Davis
1973. Predicting fire spread in Arizona's oak chaparral. USDA For. Serv. Res. Pap. RM-101, 11 p. (RM)

673. Nelson, R.M., Jr., and R.C. Corlett
1972. The heat of reaction dependence of energy feedback from small laminar diffusion flames. Comb. Sci. and Tech. 6:89-93. (SE)

674. Rothermel, R.C.
1972. A mathematical model for predicting fire spread in wildland fuels. USDA For. Serv. Res. Pap. INT-115, 40 p. (INT)

675. Sando, R.W., and D.A. Haines
1972. Fire weather and behavior of the Little Sioux Fire. USDA For. Serv. Res. Pap. NC-76, 6 p. (NC)

676. Shafizadeh, F., G.D. McGinnis, and C.W. Philpot
1972. Thermal degradation of xylan and related model compounds. Carbohydrate Res. 25: 23-33. (INT)

677. Stein, A.M., and B.W. Bauske
1972. Computer technique for simulating the combustion of cellulose and other fuels. USDA For. Serv. Res. Note PSW-266, 3 p. (PSW)

678. Stockstad, D.S.
1972. Modifications and test procedures for the Stockstad-Lory ignition furnace. USDA For. Serv. Res. Note INT-166, 7 p. (INT)

679. Wade, D.D., and D.E. Ward
1973. An analysis of the Air Force Bomb Range Fire. USDA For. Serv. Res. Pap. SE-105, 38 p. (SE)

Biology and understanding

680. Beckwith, R.C.
1972. Key to adult bark beetles commonly associated with white spruce stands in interior Alaska. USDA For. Serv. Res. Note PNW-189, 6 p. (PNW)

681. Beckwith, R.C.
1972. Scolytid flight in white spruce stands in Alaska. Can. Ent. 104: 1977-1983. (PNW)

Water quality

682. Martin, R.E., J.E. Green, III, and K.L. Hodge
1972. Oil sorption and water pollution by southern pine bark. In Technique of Proc. Bark and Util. of Bark Prod., For. Prod. Res. Soc., Madison, Wisc., p. 26-30. (PNW)

Natural regeneration

683. Zasada, J.C.
1972. Guidelines for obtaining natural regeneration of white spruce in Alaska. USDA For. Serv., PNW, Portland, Oreg., 16 p. (PNW)

684. Zasada, J.C.
1973. Interior Alaska hardwoods. In Silvicultural Systems for the Major Forest Types of the United States, USDA, Agric. Handb. 445, p. 20-22. (PNW)

685. Zasada, J.C., and R.A. Gregory
1972. Paper birch seed production in the Tanana Valley, Alaska. USDA For. Serv. Res. Note, PNW-117, 7 p. (PNW)

Ecological relationships

686. Van Cleve, K., and L.A. Viereck
1972. Distribution of selected chemical elements in evenaged alder (*Alnus*) ecosystems near Fairbanks, Alaska. Arctic and Alpine Res. 4(3): 239-255. (PNW)

Weather modification and weather effects

687. Baughman, R.G., and D.M. Fuquay
1972. USDA Forest Service weather modification activities. Fourteenth Interagency Conf. on Weather Modification Proc. Shenandoah National Park, Virginia. (Proceedings to be released October 1973). (INT)

688. Consul, P.C., and L.R. Shenton
1972. Use of Lagrange expansion for generating discrete generalized probability distributions. Siam J. of Applied Math., 23(2): 239-248, September. (SE)

689. Cramer, O.P.
1972. Mesosystem weather in the Pacific Northwest. A summer case study. Monthly Weather Review, 101(1): 13-23. (PNW)

690. Fosberg, M.A., A. Rango, and W.E. Marlatt
1972. Wind computations from the temperature field in an urban area. Conf. Urban Environ. and 2nd Conf. Biometerol. (Phila., Pa., Oct. 31-Nov. 2, 1972). Proc. 1972: 5-7, Am. Meteorol. Soc., Boston, Mass. (RM)

691. Fuquay, D.M., A.R. Taylor, R.G. Hawe, and C.W. Schmid
1972. Lightning discharges that caused forest fires. Geophys. Res. 77(12): 2156-2158. (INT)

692. Lamb, R.C., M.P. Waters, III, and E.V. Brender
1973. Apparent influence of weather upon seed production of loblolly pine. USDA For. Serv. Res. Note SE-183, 7 p. (SE)

693. Palmer, T.Y., and L.R. Smith
1973. Flame suppressants for propane cold fog seeding. J. Appl. Meteor. 12(2): 421-422. (PSW)

694. Paul, J.T., and M.P. Waters, III
1972. Lightning-fire occurrence statistics for southeastern Georgia. IN Manual of Fire-Danger and Fire-Weather Seminar, Macon, Ga., 8 p., December. (SE)

695. Shenton, L.R., and K.O. Bowman
1972. Further remarks on maximum likelihood estimators for the gamma distribution. Technometrics 14(3): 725-733. (SE)

696. Shenton, L.R., K.O. Bowman, and D. Sheehan
1971. Sampling moments of moments associated with univariate distributions. J. Royal Stat. Soc., Series B 33(3): 444-457. (SE)

697. Williams, D.T.
1972. The effect of moving weather systems on fire behavior. Proc. 1971 Southern Reg. Tech. Conf., Fla. Sec., SAF, May, Jacksonville, p. 69-75. (SE)

Miscellaneous

698. Martin, R.E., and J.L. Crawford
1972. Sorption of sulfate mill odors by bark. IN Techniques of Proc. Bark and Util. of Bark Prod., For. Prod. Res. Soc., Madison, Wisc. p. 42-46. (PNW)

TIMBER MANAGEMENT RESEARCH

Evaluating and improving animal habitat resources

699. Tubbs, C.H., and L.J. Verme
1972. How to create wildlife openings in northern hardwoods. USDA For. Serv., North Central For. Exp. Stn., How-to leaflet, 5p. (NC)

Managing recreational opportunities

700. Minore, D.
1973. Huckleberries, competing vegetation, and an experiment designed to save the berries. In Abstr. of papers scheduled for presentation at the 46th Annu. Meet. of Northwest Sci. Assoc. Pullman, Wash. State Univ. Press. (PNW)

701. Minore, D.
1972. The wild huckleberries of Oregon and Washington — a dwindling resource. USDA For. Serv. Res. Pap. PNW-143, 20 p. (PNW)

Trees to enhance the environment

702. Cook, D.I., and D.F. Van Haverbeke
1972. Trees and shrubs can curb noise, but with quite a few "ifs". USDA Yearb. 1972: 28-30. (RM)

703. Cook, D.I., and D.F. Van Haverbeke
1972. Trees, shrubs, and land-forms for noise control. J. Soil and Water Conserv. 27(6): 259-261. (RM)

704. Cook, D.I., and D.F. Van Haverbeke
1972. The potential value of trees, shrubs, and land form combinations for noise control. International Shade Tree Conference, Midwestern Chapter (Chicago, Illinois, February 1972) 27: 19-25. (RM)

705. Farnsworth, D.H., G.E. Gatherum, J.J. Jokela, H.B. Kriebel, D.T. Lester, C. Merritt, S.S. Pauley, R.A. Read, R.J. Sajdak, and J.W. Wright
1972. Geographic variation in Japanese larch in North Central United States plantations. Silvae Genet. 21(3-4):139-147. (RM)

706. Read, R.A.
1972. Plan before planting tree windbreaks. Colo. Rancher Farmer 26(2): 30,34. (RM)

707. Van Haverbeke, D.F., and D.I. Cook
1972. Green mufflers. Am. For. 78(11):28-31. (RM)

708. Zaman, M.B., A.A. Khan, and A. Ahmad
1972. Contribution to the knowledge of medicinal plants. Pakistan For. Inst., Med. Plant. Branch, Biol. Sci. Res. Div. Bull. 5, 13 p. (WO)

709. Zaman, M.B., and M.A. Khan
1972. List of medicinal and other flowering plants of Dir. Pakistan For. Inst., Med. Plant Branch, Biol. Sci. Res. Div. Bull. 6, 28 p., Index i-ix. (WO)

710. Zaman, M.B.: Khan, M.A., and M.A. Shah
1972. Quantitative survey of medicinal plants in Chitral forests. Pakistan For. Inst., Med. Plant Branch, Biol. Sci. Res. Div. Bull. 4, 9 p. (WO)

711. Zaman, M.B., M.A. Khan, and M.A. Shah
1972. Quantitative survey of medicinal plants in Dir forest division. Pakistan For. Insti., Med. Plant Branch, Biol. Sci. Res. Div. Bull. 3, 5 p. (WO)

712. Zaman, M.B., and M.S. Khan
1970. Hundred drug plants of West Pakistan. Pakistan For. Inst., Med. Plant Branch, Peshawar, 106 p. (WO)

Fire prevention, hazard reduction, and prescribed burning

713. Yocom, H.A.
1972. Burning to reduce understory hardwoods in the Arkansas Mountains. USDA For. Serv. Res. Note SO-145, 3 p. (SO)

Biology and understanding

714. Gara, R.I., G.G. Allan, R.M. Wilkins, and J.L. Whitmore
1973. Flight and host selection behavior of the mahogany shoot borer, *Hypsipyla grandella* Zeller (Lepid., Phycitidae). Zeitschrift fur Angewandte Entomologie 72(3): 259-266. (ITF)

Pest control techniques

715. Berntsen, C.M.
1971. Silvicultural control. West. For. Pest Comm. Annu. Meet. 1971: 19-21. (RM)

716. Jones, E.P., Jr.
1972. Fusiform rust affects planted slash pine. J. For. 70: 350-352. (SE)

Forest inventory

717. Stage, A.R., and J.R. Alley
1972. An inventory design using stand examinations for planning and programming timber management. USDA, For. Serv. Res. Pap. INT-126, 17 p. (INT)

Survey techniques

718. Amidon, E.L.
1972. Partial glossary of terms. In Geographic Data Handling, R.F. Tomlinson (Ed.), p. 1328-1351 IUG, Com. Geogr. Data Sensing and Process, Ottawa. (PSW)

Natural regeneration

719. Bonner, F.T.
1972. Maturation of sweetgum and American sycamore seeds. For. Sci. 18(3): 223-231. (SO)

720. Bonner, F.T.
1972. Measurement of moisture content in seeds of some North American hardwoods. Int. Seed Testing Assoc. Proc. 37(3): 975-983. (SO)

721. Brender, E.V., and W.H. McNab
1972. Loblolly pine seed production in the Lower Piedmont under various harvesting methods. J. For. 70: 345-349. (SE)

722. Chalupka, W.
1972. Cone crop and age of Norway spruce (*Picea abies* (L.) Karst.) trees and stands. Sylwan 4: 73-77. (WO)

723. DeBell, D.S., and A.W. Naylor
1972. Some factors affecting germination of swamp tupelo seeds. Ecol. 53(3): 504-506. (SE)

724. Harris, A.S.
1972. Natural reforestation after logging on Afognak Island. USDA For. Serv. Res. Note PNW-176, 11 p. (PNW)

725. Johnsen, T.N., Jr., G.H. Schubert, and D.P. Almas
1973. Forest regeneration competition problems: Rocky Mountain-Intermountain Region. J. For. 71: 144-147. (RM)

726. Johnston, W.F.
1972. Balsam fir dominant species under rethinned northern white-cedar. USDA For. Serv. Res. Note NC-133, 4p. (NC)

727. Minore, D.
1972. Germination and early growth of coastal tree species on organic seed beds. USDA For. Serv. Res. Pap. PNW-135, 18 p. (PNW)

728. Noble, D.L.
1973. Age of Engelmann spruce seedlings affect ability to withstand low temperature: a greenhouse study. USDA For. Serv. Res. Note RM-232, 4 p. (RM)

729. Noble, D.L.
1972. Effect of soil type and watering on germination, survival, and growth of Engelmann spruce: a greenhouse study. USDA For. Serv. Res. Note RM-216, 4 p. (RM)

730. Sander, I.L.
1972. Size of oak advance reproduction: key to growth following harvest cutting. USDA For. Serv. Res. Pap. NC-79, 6 p. (NC)

731. Van Deusen, J.L., and L.D. Beagle
1973. Judging ripeness of seeds in Black Hills ponderosa pine cones. USDA For. Serv. Res. Note RM-235, 4 p. (RM)

732. Wendel, G.W.
1972. Longevity of black cherry seed in the forest floor. USDA For. Serv. Res. Note NE-149, 4 p. (NE)

733. Zasada, J.C., and R.A. Gregory
1972. Paper birch seed production in the Tanana Valley, Alaska. USDA For. Serv. Res. Note PNW-177, 7 p. (PNW)

Silvicultural methods

734. Alexander, R.R.
1972. Partial cutting practices in old-growth lodgepole pine. USDA For. Serv. Res. Pap. RM-92, 16 p. (RM)

735. Alexander, R.R.
1972. Partial cutting practices in old-growth lodgepole pine: a field guide. USDA For. Serv. Res. Pap. RM-92A, 13 p. (RM)

736. Benzie, J.W.
1972. Ecological and environmental aspects of logging. Am. Pulpwood Assoc. Tech. Pap., p. 5-8. (NC)

737. Benzie, J.W.
1972. Harvesting patterns for red pine. In Proc. of Conf. on Biol. and Econ. Considerations in Mech. Timber Harvesting. Univ. of Minn., Agric. Exp. Stn., Misc. Rep. 116, For. Series 11, p. 21-24. (NC)

738. Benzie, J.W.
1972. Silvicultural interpretations of mechanized harvesting research results (for partial cuttings and multiple use). In Proc. of Conf. on Biol. and Econ. Considerations in Mech. Timber Harvesting. Univ. of Minn., Agric. Exp. Stn., Misc. Rep. 116, For. Series 11, p. 26-28. (NC)

739. Benzie, J.W., and Z.A. Zasada
1972. Shelterwood-strip harvesting pattern with full-tree skidding to regenerate red pine. USDA For. Serv. Res. Note NC-132, 4 p. (NC)

740. Brady, H.A.
1972. Competition control in new woody plantings. In 25th South. Weed Sci. Soc. Proc. p. 249-251. (WO)

741. Cooper, G.A., C.F. Bey, R.D. Lindmark, and R.C. Schlesinger
1972. Better hardwoods — from seeds to sawlogs and mills to market. South. Lumberman 225(2800):111-114. (NC)

742. Franklin, J.F., and D.S. DeBell
1972. Effects of various harvesting methods on forest regeneration. In Even-age Management, R.K. Herman and D.P. Lavender (eds). Proc. of Symp., Aug. 1, 1972, School of For., Oreg. State Univ., Corvallis, p. 29-57. (PNW)

743. Gysel, L.W., J.H. Cooley, and F. Stearns
1972. Timber and wildlife implication of pine conversion methods on poor oak sites. J. For. 70: 272-274. (NC)

744. McGee, C.E.
1972. From a defective hardwood stand multiple use opportunity. J. For. 70: 700-704. (SE)

745. Marquis, D.A.
1972. Effect of forest clearcutting on ecological balance. In A Perspective on Clearcutting in a Changing World. Proc. 1972 Winter Meet., New York Sect., Soc. Am. For., Applied For. Res. Inst. Misc. Rep. 4, Syracuse, NY, p. 47-58. (NE)

746. Perala, D.A.
1972. Regeneration: biotic and silvicultural factors. In Aspen Symp. Proc. USDA For. Serv. Gen. Tech. Rep. NC-1, p. 97-102. (NC)

747. Schlaegel, B.E.
1972. Growth and yield of managed stands. In Aspen Symp. Proc. USDA For. Serv. Gen. Tech. Rep. NC-1, p. 109-112. (NC)

748. U.S. Department of Agriculture, Forest Service
1973. Silvicultural systems for the major forest types of the United States. U.S. Dept. Agric. Handb. 445, 114 p. (WO)

Ecological relationships

749. Franklin, J.F.
1972. Why a coniferous forest biome? In Research on Coniferous Forest Ecosystems - A Symposium. Jerry F. Franklin, L.J. Dempster, Richard H. Waring (eds.) USDA For. Serv., Pacific Northw. For. Range Exp. Stn., Portland, Oreg. p. 3-5. (PNW)

750. Franklin, J.F., L.J. Dempster, and R.H. (eds) Waring
1972. Research on coniferous forest ecosystems — A symposium. USDA For. Serv., Pacific Northwest. For. Range Exp. Stn., Portland, Oreg. 322 p. (PNW)

751. Franklin, J.F., F.C. Hall, C.T. Dyrness, and C. Maser
1972. Federal Research Natural Areas in Oregon and Washington: A guidebook for scientists and educators. USDA For. Serv., Pacific Northwest. For. Exp. Stn., Portland, Oreg., 498 p. (PNW)

752. Leak, W.B.
1972. Competitive exclusion in forest trees. Nature 236(5348): 461-463. (NE)

753. Leak, W.B.
1972. Systems ecology. Ecol. 53: 199-200. (NE)

754. Leak, W.B.
1971. Simulation of forest succession. Summer Comp. Simulation Conf. Proc., USDA For. Serv. Northeast. For. Exp. Stn., Upper Darby, PA., p. 820-825. (NE)

755. Safford, L.O., and S. Bell
1972. Biomass of fine roots in a white spruce plantation. Can. J. For. Res. 2(3): 169-172. (NE)

756. Seidel, K.W.
1972. Drought resistance and internal water balance of oak seedlings. For. Sci. 18(1): 34-40. (NC)

757. Wenger, K.F., C.E. Ostrom, P.R. Larson, and T.D. Rudolph
1972. Potential effects of global atmospheric conditions on forest ecosystems. In Man's Impact on Terrestrial and Oceanic Ecosystems, W.H. Matthrews, F.E. Smith, and E.D. Goldberg (eds.), Chapt. 12, p. 192-202. MIT Press, Cambridge, Mass. (NC)

Site evaluation and soil improvement

758. Auchmoody, L.R.
1972. Foliar nutrient variation in four species of upland oak. In Agron. Abstr., p. 136. (NE)

759. Auchmoody, L.R.
1972. Effects of fertilizer-nutrient interactions on red oak seedling growth. USDA For. Serv. Res. Pap. NE-239, 5 p. (NE)

760. Baker, J.B., and R.H. Brendemuehl
1972. Soil phosphorus level adequate for growth of Ocala sand pine seedlings a greenhouse evaluation. Soil Sci. Soc. of Am. Proc. 36: 666-667. (SE)

761. Bjorkbom, J.C.
1973. Response of paper birch seedlings to nitrogen, phosphorus, and potassium. USDA For. Serv. Res. Note NE-157, 4 p. (NE)

762. Bjorkbom, J.C.
1973. The effects of various combinations of nitrogen, phosphorus, and potassium on paper birch seedling growth. USDA For. Serv. Res. Note NE-158, 4 p. (NE)

763. Blackmon, B.G., and E.H. White
1972. Nitrogen fertilization increases cottonwood growth on old-field soil. USDA For. Serv. Res. Note SO-143, 5 p. (SO)

764. Carmean, W.H.
1972. Site index curves for upland oaks in the Central States. For. Sci. 18: 109-120. (NC)

765. Della-Bianca, L., and C.E. McGee
1972. Reaction of natural *Rhododendron maximum* L. to liming in the Southern Appalachians. J. Elisha Mitchell Sci. Soc. 88: 109-112. (SE)

766. Dickson, R.E., T.C. Broyer, and C.M. Johnson
1972. Nutrient uptake by tupelo gum and bald cypress from saturated or unsaturated soil. Plant and Soil 37: 297-308. (NC)

767. Dickson, R.E., and T.C. Broyer
1972. Effects of aeration, water supply, and nitrogen source on growth and development of tupelo gum and bald cypress. Ecol. 53: 626-634.

768. Ferguson, E.R., and D.L. Graney
1972. Site index table for shortleaf pine in the Boston Mountains of Arkansas. USDA For. Serv. Res. Note SO-137, 4 p. (SO)

769. Graney, D.L., and E.R. Ferguson
1972. Shortleaf pine site-index relationships in the Ozark Highlands. Soil Sci. Soc. Am. Proc. 36: 495-500. (SO)

770. Hatchell, G.E., and O.G. Langdon
1972. Carolina studies offer possibilities of increasing pine growth on savannah sites. Tree Plant. Notes 23(2): 19-21. (SE)

771. Hoyle, M.C.
1972. Manganese toxicity in yellow birch (*Betula alleghaniensis* Britton) seedlings. Plant and Soil 37: 229-232. (NE)

772. Losche, C.K.
1972. Vertical tillage fails to improve black walnut seedling growth on poorly drained site. Tree Plant. Notes 23(1): 7-9. (NC)

773. Losche, C.K., W.M. Clark, E.E. Voss, and B.S. Ashley
1972. Guide to the selection of soil suitable for growing black walnut in Illinois. Spec. Publ., Northeast. Area, State and Private For., Upper Darby, Pa., 38 p. (NC)

774. Losche, C.K., and R.E. Phares
1972. Siltation damage in a black walnut plantation. J. Soil and Water Conserv. 27(5): 228-229. (NC)

775. Oliver, W.W.
1972. Height intercept for estimating site index in young ponderosa pine plantations and natural stands. USDA For. Serv. Res. Note PSW-276, 4 p. (PSW)

776. Phares, R.E., and R.F. Finn
Using foliage analysis to help diagnose nutrient deficiencies in black walnut. North. Nut Grow. Assoc. Annu. Rep. 62: 98-104. (NC)

777. Powers, R.F.
1972. Estimating site index of ponderosa pine in northern California. USDA For. Serv. Res. Note PSW-265, 8 p. (PSW)

778. Powers, R.F.
1972. Site index curves for managed stands of California black oak. USDA For. Serv. Res. Note PSW-262, 5 p. (PSW)

779. Watt, R.F., and M.E. Newhouse
1973. Some soil phases in the Missouri Ozarks have similar site indexes for oaks. USDA For. Serv. Res. Pap. NC-86, 5 p. (NC)

780. Wells, C.G., D. Whigham, and H. Leith
1972. Investigation of mineral nutrient cycling in an upland Piedmont forest. J. Elisha Mitchell Sci. Soc. 88: 66-78. (SE)

Artificial regeneration

781. Bakshi, B.K., M.A. Ram Reddy, H.S. Thapar, and S.N. Khan
1972. Studies on silver fir regeneration. Indian For. 98:135-144. (WO)

782. Barnett, J.P.
1972. Seedcoat influences dormancy of loblolly pine seeds. Can. J. For. Res. 2:7-10. (SO)

783. Barnett, J.P.
1972. Drying and storing stratified loblolly pine seeds reinduces dormancy. Tree Plant. Notes 23(3):10-22. (SO)

784. Barnett, J.P.
1972. Southern pine seeds germinate after forty years' storage. J. For. 70:629. (SO)

785. Benzie, J.W., S. Little, and R.F. Sutton
1973. Rehabilitation of forest land: the Northeast and Boreal Region. J. For. 71:154-158. (NC)

786. Bjorkbom, J.C.
1972. Ten-year growth of planted paper birch in old fields in Maine. USDA For. Serv. Res. Pap. NE-246, 6 p. (NE)

787. Bjorkbom, J.C.
1972. Stand changes in the first 10 years after seedbed preparation for paper birch. USDA For. Serv. Res. Pap. NE-238, 10 p. (NE)

788. Boyd, R.J., S. McDonald, and L.L. Mason
1972. Estimation of survival and growth potentials of nursery stock by using a "variable-moisture-stress-plot" technique. USDA For. Serv. Res. Note INT-165, 8 p. (INT)

789. Bramlett, D.L.
1973. Pollen phenology and dispersal pattern for shortleaf pine in the Virginia Piedmont. USDA For. Serv. Res. Pap. SE-104, 7 p. (SE)

790. Bramlett, D.L.
1972. Cone crop development records for six years in shortleaf pine. For. Sci. 18:31-33. (SE)

791. Bramlett, D.L.
1972. Seed and aborted ovules from cones of *Pinus echinata*. Ga. For. Res. Counc., Res. Pap. 71, 4 p. (SE)

792. Branan, J.R., and E.J. Porterfield
1971. A comparison of six species of southern pines planted in the Piedmont of South Carolina. USDA For. Serv. Res. Note SE-171, 3 p. (SE)

793. Burns, R.M., and E.A. Hebb
1972. Site preparation and reforestation of droughty, acid sands. USDA, Agric. Handb. 426, 61 p. (SE)

794. Burns, R.M., and R.D. McReynolds
1972. Scheduling and intensity of site preparation for pine in West Florida Sandhills. J. For. 70:737-740. (SE)

795. Wilson, B.C., and R.K. Campbell
1972. Seedbed density influences height, diameter, and dry weight of 3-0 Douglas-fir. Tree Plant. Notes 23(3):1-4. (PNW)

796. Cooley, J.H.
1972. Site preparation for jack pine on Grayling sands. USDA For. Serv. Res. Note NC-138, 3 p. (NC)

797. Geary, T.F., and C.B. Briscoe
1972. Tree species for plantations in the granitic uplands of Puerto Rico. USDA For. Serv. Res. Pap. ITF-14, 8 p. (ITF)

798. Geary, T.F., and J.A. Zambrana
1972. Must Honduras pine be weeded frequently in Puerto Rico? USDA For. Serv. Res. Pap. ITF-16, 16 p. (ITF)

799. Geary, T.F., G. Avila Cortes, and H.H. Hadley
1971. Germination and growth of *Pinus caribaea* directly sown into containers as influenced by shade, phosphate, and captan. Turrialba 21:336-342. (ITF)

800. Geary, T.F., R.W. Nobles, and C.B. Briscoe
1972. Hybrid mahogany recommended for planting in the Virgin Islands. USDA For. Serv. Res. Pap, ITF-15, 4 p. (ITF)

801. Gratkowski, H., D. Hopkins, and P. Lauterbach
1972. Rehabilitation of forest land: the Pacific Coast and Northern Rocky Mountain Region. J. For. 71:138-143. (PNW)

802. Grigsby, H.C.
1971. Handling prior to sticking affects rooting of loblolly pine cuttings. Int. Plant Propag. Soc. Proc. 21:398-401. (SO)

803. Huntzinger, H.J.
1972. Direct seeding black cherry: some recommendations for the Allegheny Plateau. Tree Plant. Notes 23(1):10-11. (NE)

804. Johnston, W.F.
1972. Seeding black spruce on brushy lowland successful if vegetation density kept low. USDA For. Serv. Res. Note NC-139, 4 p. (NC)

805. Kennedy, H.E., Jr.
1972. Hortizontal planting of green ash cuttings looks promising. USDA For. Serv. Res. Note SO-147, 3 p. (SO)

806. Lamb, R.C., M.P. III Waters, and E.V. Brender
1973. Apparent influence of weather upon seed production of loblolly pine. USDA For. Serv. Res. Note SE-183, 7 p. (SE)

807. Lohrey, R.E.
1972. Precommercial thinning of direct-seeded loblolly pine. USDA For. Serv. Res. Note SO-139, 4 p. (SO)

808. McAlpine, R.G., D.D. Hook, and P.P. Kormanik
1972. Horizontal planting of sycamore cuttings. Tree Plant. Notes 23(2):5-7. (SE)

809. McBride, J.R., and R.E. Dickson
1972. Gibberellic, citric acids and stratification enhance white ash germination. Tree Plant. Notes 23(3):1-2. (NC)

810. McLemore, B.F.
1972. Determining numbers of southern pine cones per bushel. J. For. 70:35-36. (SO)

811. Olson, D.F., Jr., and R.M. Hooper
1972. Northern red oak plantings survive well in Southern Appalachians. Tree Plant. Notes 23(1):16-18. (SE)

812. Owston, P.W., and W.I. Stein
1972. First-year performance of Douglas-fir and noble fir outplanted in large containers. USDA For. Serv. Res. Note PNW-174, 10 p. (PNW)

813. Phares, R.E., and G. White
1972. Large stock, deep planting improve cotton wood growth in Upper Mississippi Valley. Tree Plant. Notes 23(4):16-17. (NC)

814. Phipps, H.M.
1973. Growth response of some tree species to plastic greenhouse culture. J. For. 71:28-30. (NC)

815. Ronco, F.
1972. Planting Engelmann spruce. USDA For. Serv. Res. Pap. RM-89, 24 p. (RM)

816. Ronco, F.
1972. Planting Engelmann spruce: a field guide. USDA For. Serv. Res. Pap. RM-89A, 11 p. (RM)

817. Rowan, S.J.
1972. Hardwood seedling production trends in southern nurseries 1965 - 1970. J. For. 70:162-163. (SE)

818. Russell, T.E.
1973. Animal depredations on spot-seeded yellow-poplar in central Tennessee. USDA For. Serv. Res. Note SO-148, 4 p. (SO)

819. Schmidtline, R.C.
1972. Replacement planting with potted southern pines on research plots. USDA For. Serv. Res. Note SO-146, 4 p. (SO)

820. Schubert, G.H., and J.A. Pitcher
1973. A provisional tree seed-zone and cone-crop rating system for Arizona and New Mexico. USDA For. Serv. Res. Pap. RM-105, 8 p. (RM)

821. Shubert, G.H., and R.S. Adams
1971. Reforestation practices for conifers in California. Calif. Dep. Conserv., Div. For., 359 p. (RM)

822. Schultz, R.P.
1972. Intraspecific root grafting in slash pine. Bot. Gaz. 133:26-29. (SE)

823. Schultz, R.P.
1971. Stimulation of flower and seed production in a young slash pine orchard. USDA For. Serv. Res. Pap. SE-91, 10 p. (SE)

824. Shoulders, E.
1972. Mycorrhizal inoculation influences survival, growth, and chemical composition of slash pine seedlings. USDA For. Serv. Res. Pap. SO-74, 12 p. (SO)

825. Smalley, G.S., and K. Pierce
1972. Yellow-poplar, loblolly pine, and Virginia pine compared in Cumberland Plateau plantations. USDA For. Serv. Res. Note SO-141, 6 p. (SO)

826. Steinbeck, K., R.G. McAlpine, and J.T. May
1972. Short rotation culture of sycamore: a status report. J. For. 70:210-213. (SE)

827. Strothmann, R.O.
1972. Douglas-fir in northern California: effects of shade on germination, survival, and growth. USDA For. Serv. Res. Pap. PSW-84, 10 p. (PSW)

828. Venator, C.R., and J.A. Zambrana
1972. Extraction and germination of Kadam seed. USDA For. Serv. Res. Note ITF-14, 2 p. (ITF)

829. Walters, G.A.
1972. Packing methods studied for Australian toon and slash pine plantings. Tree Plant. Notes 23(4):7-9. (PSW)

830. Walters, G.A.
1972. Pesticide treatments on saligna eucalyptus, Australian toon seedlings affect dieback but not survival. Tree Plant. Notes 23(3):16-18. (PSW)

831. Walters, G.A.
1972. Chemical treatment of bare-root saligna eucalyptus seedlings offers no advantages. Tree Plant. Notes 23(4):4-7. (PSW)

832. Walters, G.A.
1972. Survival of tropical ash planted in tofdon-treated soils in Hawaii. USDA For. Serv. Res. Note PSW-263, 4 p. (PSW)

833. Wendel, G.W.
1972. Results of a 20-year test of hybrid poplars in West Virginia. USDA For. Serv. Res. Pap. NE-237, 5 p. (NE)

834. Williams, R.D.
1972. Storing black walnut seed. North Nut Grow. Assoc. Annu. Rep. 62:87-89. (NC)

835. Williams, R.D.
1972. Root fibrosity proves insignificant in survival, growth of black walnut seedlings. Tree Plant. Notes 23(2):22-25. (NC)

836. Works, D.W., and R.J. Boyd
1972. Using infrared irradiation to decrease germination time and increase percent germination in various species of western conifer trees. Am. Soc. Agric. Eng. Trans. 15:760-762. (INT)

837. Zaman, M.B., M.A. Khan, and M.A. Shah
1972. Cultivation prospects of *Datura metel* L. in Changa Manga plantation, Lahore. Pakistan J. For. 22:467-473. (WO)

Stand improvement

838. Barrett, J.W.
1972. Large-crowned planted ponderosa pine respond well to thinning. USDA For. Serv. Res. Note PNW-179, 12 p. (PNW)

839. Bennett, F.A.
1971. Modern thinning practice—technical and economic effects. In Southern Pine Management--Today and Tomorrow, Proc. 20th Annu. For. Symp., La. State Univ. Div. Contin. Educ., Baton Rouge, 33-45. (SE)

840. Bennett, F.A.
1971. Techniques in silvicultural operations—tending, with emphasis on mechanization. XV IUFRO Cong. Proc. Publ. No. 1:150-162. (SE)

841. Bey, C.F.
1972. Corrective pruning young black walnut trees—a new twist. 63rd Annual Report of the Northern Nut Growers Assoc., p. 26-28. (NC)

842. Brady, H.A.
1972. Drop size affects absorption of 2,4,5-T by six hardwood species. In 25th South. Weed Sci. Soc. Proc., p. 282-286. (SO)

843. Dietschman, G.H., and R.D. Pfister
1973. Growth of released and unreleased young stands in the western white pine type. USDA For. Serv. Res. Pap. INT-132, 14 p. (INT)

844. Della-Bianca, L.
1972. Screening some stand variables for post-thinning effect on epicormic sprouting in even-aged yellow poplar. For. Sci. 18:155-158. (SE)

845. Echols, R.M.
1972. Patterns of wood density distribution and growth rate in ponderosa pine. Proc. Symp. on the Effect of Growth Acceleration on the Porperties of Wood. Madison, Wisc., 1971, p. H-1 to H-16. (PSW)

846. Frank, R.M.
1973. The course of growth response in released white spruce—10-year results. USDA For. Serv. Res. Pap. NE-258, 6 p. (NE)

847. Gratkowski, H., and R. Stewart
1973. Aerial spray adjuvants for herbicidal drift control. USDA For. Serv. Gen. Tech. Rep. PNW-3, 18 p. (PNW)

848. Peevy, F.A.
1973. Bromacil and picloram under southern upland hardwoods. J. Weed Sci. Soc. Am. 21:54-56. (SO)

849. Peevy, F.A.
1972. Injection treatments for killing bottomland hardwoods. Weed Sci. 20:566-568. (SO)

850. Peevy, F.A.
1972. Injection treatments for controlling resistant hardwood species. In 25th South. Weed Sci. Soc. Proc., p. 252-256. (SO)

851. Peevy, F.A.
1972. How to kill hardwoods by injection. Weeds Today 3(1):8-9, 17. (SO)

852. Schlesinger, R.C.
1972. Sweep and crook in green ash saplings—less after 11 years. J. For. 70:687. (NC)

853. Trimble, G.R., Jr.
1973. Response to crop-tree release by 7-year-old stems of yellow-poplar and black cherry. USDA For. Serv. Res. Pa. NE-253, 10 p. (NE)

854. Walters, G.A.
1972. Coppicing to convert small cull trees to growing stock. USDA For. Serv. Res. Note PSW-272, 4 p. (PSW)

Animal damage

855. Crouch, G.L., and M.A. Radwan
1972. Arasan in endrin treatments to protect Douglas-fir seed from deer mice. USDA For. Serv. Res. Pap. PNW-136, 7 p. (PNW)

856. Heidmann, L.J.
1972. An initial assessment of mammal damage in the forests of the Southwest. USDA For. Serv. Res. Note RM-219, 7p. (RM)

857. Radwan, M.A.
1972. Differences between Douglas-fir genotypes in relation to browsing preference by black-tailed deer. Can. J. For. Res. 2:250-255. (PNW)

858. Radwan, M.A.
1972. Occurrence and genotypic differences of chlorogenic acid in Douglas-fir foliage. USDA For. Serv. Res. Note PNW-173, 6 p. (PNW)

Growth requirements

859. Anand, V.K., R.N. Chibbar, and K.K. Nanda
1972. Effects of Ga3 and IBA on rooting and on the sprouting of buds on stem cuttings of *Ipomoea fistulosa*. Plant and Cell Physiol. 13:917-921. (WO)

860. Burns, R.M.
1972. Choctawhatchee sand pine, good prospect for Georgia-Carolina sandhills. J. For. 70:741-742. (SE)

861. Cochran, P.H.
1972. Temperature and soil fertility affect lodgepole and ponderosa pine seedling growth. For. Sci. 18:132-134. (PNW)

862. Cochran, P.H.
1972. Tolerance of lodgepole and ponderosa pine seeds and seedlings to high water tables. Northwest Sci. 46:322-321. (PNW)

863. Fahn, A., R. Ben-Sasson, and T. Sachs
1972. The relation between the procambium and the cambium. In Research Trends in Plant Anatomy--K.A. Chowdhury Commemoration Volume, p. 161-170, A.K.M. Ghouse and M. Yunus (eds.), Tata McGraw-Hill, New Delhi. (WO)

864. Hacskaylo, E.
1972. Mycorrhiza: The ultimate in reciprocal parasitism. Bio Sci. 22:577-583. (NE)

865. Hacskaylo, E., and G. Bruchet
1972. Hebelomas as mycorrhizal fungi. Bull. Torrey Bot. Club 99:17-20. (NE)

866. Haissig, B.E.
1972. Meristematic activity during adventitious root primordium development. Plant Physiol. 49:886-892. (NC)

867. Haissig, B.E., and A.L. Schipper, Jr.
1972. Easy extraction of enzymes from small amounts of woody tissue. Anal. Biochem. 28:129-146. (NC)

868. Harms, W.R., and O.G. Langdon
1972. Maximizing volume and dry matter yields of loblolly pine. TAPPI Sixth For. Biol. Conf.:II-2, 2 p. (Abstr.). (SE)

869. Hoyle, M.C.
1972. Indoleacetic acid oxidase: a dual catalytic enzyme. Plant Physiol. 50:15-18. (NE)

870. Jensen, K.F., and L.S. Dochinger
1972. Gibberellic acid and height growth of white pine seedlings. For. Sci. 18:196-197. (NE)

871. Jones, J.R.
1972. Moisture stresses in Arizona mixed conifer seedlings. USDA For. Serv. Res. Pap. RM-86, 8 p. (RM)

872. Kirscher, H., and T. Sachs
1972. Correlative inhibition between strips of vascular tissue. Israel J. Bot. 21:129-134. (WO)

873. Krajicek, J.E.
1972. Spacing and pruning black walnut to produce both nuts and lumber. North. Nut Grow. Assoc. Annu. Rep. 62:95-98. (NC)

874. Larson, P.R.
1972. Interpretation of radioisotope translocation patterns in forest trees. IN Isotopes and Radiation in Soil-Plant Relationships Including Forestry, Int. Atomic Energy Agency, Vienna, Austria, p. 277-288. (NC)

875. Larson, P.R.
1972. Putting basic research into practice. Tech. Pap. Am. Pulpwood Assoc., p. 17-23. (NC)

876. Larson, P.R., and J.G. Isebrands
1972. The relation between leaf production and wood weight in first-year sprouts of two Populus clones. Can. J. For. Res. 2:98-104. (NC)

877. Larson, P.R., J.G. Isebrands, and R.E. Dickson
1972. Fixation patterns of 14/C within developing leaves of eastern cottonwood. Planta 107:301-314. (NC)

878. McKee, W.H., Jr.
1972. Light and temperature influence phosphorus absorption by slash pine. Commun. Soil. Sci. and Plant Anal. 3:223-229. (SO)

879. Nanda, K.K., and M.K. Jain
1972. Mode of action of IAA and GA3 on root and shoot growth of epiphyllous nuds of *Bryophyllum tubiflorum*. J. Exp. Bot. 23:980-986. (WO)

880. Nanda, K.K., V.K. Anand, and R.N. Chibbar
1972. The promotive effect of gibberellic acid on the production of adventitious roots on stem cuttings of *Ipomoea fistulosa*. Planta (Berl.) 105, 360-363. (WO)

881. Peters, J.K., and A.E. Squillace
1972. Genetic effects, chemistry, and foxtail growth patterns in *Pinus elliottii*. World For. Cong. Oct. 1972 (Proc. in press). (SE)

882. Reddy, M.A.R., and S.N. Khan
1972. Soil amendments and types of inocula on development of mycorrhiza. Indian For. 98:307-310. (NE)

883. Ronco, F.
1972. Overwinter food reserves of potted Engelmann spruce seedlings. Can. J. For. Res. 2:489-492. (RM)

884. Rudolph, T.D.
1972. The effect of acute gamma irradiation at various stages of seasonal growth in *Pinus banksiana* Lambert. Proc. IAEA/FAO Symp. on the Use of Isotopes and Radiation in Research on Soil-Plant Relationships Including Applications in Forestry, Vienna, Austria, Dec. 13-17, 1971, p. 289-300. (NC)

885. Benator, C.R.
1972. Effect of gibberellic acid on germination of low-vigor Honduras pine seeds. For. Sci. 18(4):331. (ITF)

886. Whitmore, J.L.
1972. *Pinus merkusii* unsuitable for plantations in Puerto Rico. Turrialba 22(3):351-353. (ITF)

887. Zelawski, W.
1972. Uptake and evolution of carbon dioxide and accumulation of organic substance in scot pine (*Pinus silvestris* L.) seedlings. Bulletin De L'Academie Polonaise Des Sciences 20: 747-753. (WO)

888. Zelawski, W., J. Kucharska, and J. Kinelska
1971. Relationship between dry matter production and carbon dioxide absorption in seedlings of Scots pine (*Pinus silvestris* L.) in their second vegetation season. (WO)

Growth and yield

889. Beck, D.E., and L. Della-Bianca
1972. Growth and yield of thinned yellow-poplar. USDA For. Serv. Res. Pap. SE-101, 20 p. (SE)

890. Bell, J.F., and A.B. Berg
1972. Levels-of-growing-stock cooperative study of Douglas-fir—Report No. 2--The Hoskins study, 1963-1970. USDA For. Serv. Res. Pap. PNW-130, 19 p. (PNW)

891. Boldt, C.E., and C.D. Markstrom
1972. Rapid growth and wood quality in Black Hills ponderosa pine. For. Prod. Res. Soc., Dallas, Tex., June 1972, Abstr. 26:6. (RM)

892. Bruce, D.
1972. Management of Douglas-fir in Europe. In Managing Young Forests in the Douglas-fir Region, Alan B. Berg (ed.), Vol. 3, Oreg. State Univ. Symp. Proc. 1970:1-15. (PNW)

893. Burton, J.D., and D.M. Smith
1972. Guying to prevent wind sway influences loblolly pine growth and wood properties. USDA For. Serv. Res. Pap. SO-80, 8 p. (SO)

894. Curtis, R.O.
1972. Yield tables past and present. J. For. 70:28-32. (PNW)

895. Dahms, W.G.
1973. Tree growth and water use response to thinning. USDA For. Serv. Res. Note PNW-194, 14 p. (PNW)

896. Dahms, W.G.
1971. Growth and soil moisture in thinned lodgepole pine. USDA For. Serv. Res. Pap. PNW-127, 32 p. (PNW)

897. Dale, M.E.
1972. Growth and yield predictions for upland oak stands 10 years after initial thinning. USDA For. Serv. Res. Pap. NE-241, 21 p. (NE)

898. Enghardt, H.G., and W.F. Mann
1972. Ten-year growth of planted slash pine after early thinnings. USDA For. Serv. Res. Pap. SO-82, 11 p. (SO)

899. Foiles, M.W.
N. Responses in western white pine stand in commercial thinning methods. USDA For. Serv. Res. Note INT-159, 8 p. (INT)

900. Geyer, W.A., and N.F. Rogers
1972. Spoils change and tree growth on coal-mined spoils in Kansas. J. Soil & Water Conserv. 27: 114-116. (NC)

901. Hatchell, G.E., K.W. Dorman, and O.G. Langdon
1972. Performance of loblolly and slash pine nursery selections. For. Sci. 18: 308-313. (SE)

902. Mann, W.F., and H.G. Enghardt
1972. Growing of planted slash pine under several thinning regimes. USDA For. Serv. Res. Pap. SO-76, 10 p. (SO)

903. Myers, C.A., and R.R. Alexander
1972. Bark thickness and past diameters of Engelmann spruce in Colorado and Wyoming. USDA For. Serv. Res. Note RM-217, 2 p. (RM)

904. Oliver, W.W.
1972. Growth after thinning ponderosa and Jeffery pine pole stands in northeastern California. USDA For. Serv. Res. Pap. PSW-85, 8 p. (PSW)

905. Reukema, D.L.
1972. Twenty-one-year development of Douglas-fir stands repeatedly thinned at varying intervals. USDA For. Serv. Res. Pap. PNW-141, 23 p. (PNW)

906. Sassaman, R.W., J.W. Barrett, and J.G. Smith
1972. Economics of thinning stagnated ponderosa pine sapling stands in the pine-grass areas of central Washington. USDA For. Serv. Res. Pap. PNW-144, 17 p. (PNW)

907. Saucier, J.R., A. Clark, and R.G. McAlpine
1972. Above ground biomass yields of short-rotation sycamore. Wood Sci. 5(1): 1-6. (SE)

908. Schultz, R.P.
1972. Root development of intensively cultivated slash pine. Soil Sci. Soc. Am. Proc. 36:158-162. (SE)

Forest measurements

909. Alban, D.H.
1972. An improved growth intercept method for estimating site index of red pine. USDA For. Serv. Res. Pap. NC-80, 7 p. (NC)

910. Alban, D.H.
1972. The relationship of red pine site index to soil phosphorus extracted by several methods. Soil Sci. Soc. Am. Proc. 36: 664-666. (NC)

911. Bennett, F.A.
1972. Cubic yields for slash pine in soil bank plantings. USDA For. Serv. Res. Note SE-182, 7 p. (SE)

912. Bennett, F.A., and B.F. Swindel
1972. Taper curves for planted slash pine. USDA For. Serv. Res. Note SE-179, 4 p. (SE)

913. Bruce, D.
1972. Some transformations of the Behre equation of tree form. For. Sci. 18(2):164-166. (PSW)

914. Cole, D.M., and A.R. Stage
1972. Estimating future diameters of lodgepole pine trees. USDA For. Serv. Res. Pap. INT-131, 20 p. (INT)

915. Ffolliot, P.F., and D.P. Worley
1973. Forest stocking equations: their development and application. USDA For. Serv. Res. Pap. RM-102, 8 p. (RM)

916. Grosenbaugh, L.R.
1973. Metrication and forest inventory. J. For. 71: 84-86. (SE)

917. Herman, F.R.
1972. Improved adapter for increment borer ratchet assembly. USDA For. Serv. Res. Note PNW-192, 4 p. (PNW)

918. Herman, F.R., C.E. Smith, and J.E. Firth
1972. Freezing decayed wood to facilitate ring counts and width measurements. USDA For. Serv. Res. Note PNW-187, 4 p. (PNW)

919. Herman, F.R., C.E. Smith, and J.E. Firth
1972. Ring count of decayed wood made easier through freezing. J. For. 70:743. (PNW)

920. Johnson, F.A., and G.B. Hartman
1972. Fall, buck, and scale cruising. J. For. 70: 566-568. (PNW)

921. Myers, C.A.
1972. Volume, taper, and related tables for southwestern ponderosa pine. USDA For. Serv. Res. Pap. RM-2 (rev.) 24 p. (RM)

922. Myers, C.A., and R.R. Alexander
1972. Bark thickness and past diameters of Engelmann spruce in Colorado and Wyoming. USDA For. Serv. Res. Note RM-217, 2 p. (RM)

923. Myers, C.A., and C.B. Edminster
1972. Volume tables and point-sampling factors for Engelmann spruce in Colorado and Wyoming. USDA For. Serv. Res. Pap. RM-95, 23 p. (RM)

924. Shepperd, W.D.
1973. An instrument for measuring tree crown width. USDA For. Serv. Res. Note RM-229, 3 p. (RM)

925. Swindel, B.F., and D.O. Yandle
1972. Allocation in stratified sampling as a game. J. Am. Stat. Assoc. 67(339): 684-686. (SE)

Management planning

926. Buckman, R.E.
1971. On research needs for future forest management. In Western Forestry Speaks, 62nd West. For. Conf. Proc., p. 63-67. (PNW)

927. Gansner, D.A., J.E. Barnard, and S.F. Gingrich
1973. Identifying regional opportunities for accelerated timber management. USDA For. Serv. Res. Pap. NE-251, 6 p. (NE)

928. Grosenbaugh, L.R.
1972. Computer systems, access, and administration. In Prod. Workshop on Computer and Information Systems in Resources Management Decisions. USDA For. Serv. and Coop. State Res. Serv., Wash., D.C., p. 22-34. (SE)

929. Myers, C.A., F.G. Hawksworth, and P.C. Lightle
1972. Simulating yields of southwestern ponderosa pine stands, including effects of dwarf mistletoe. USDA For. Serv. Res. Pap. RM-87, 16 p. (RM)

930. Neal, R.L., Jr.
1972. Cascading 9100B programs with protected stored data. Hewlett-Packard Keyboard 4(2):35. (PSW)

931. Neal, R.L., Jr.
1972. Saving a register when cumulating unordered data in 9100A/B. Hewlett-Packard Keyboard 4(3):20. (PSW)

932. Swindel, B.F.
1972. The Bayesian controversy. USDA For. Serv. Res. Pap. SE-95, 12 p. (SE)

933. Ware, K.D., J.L. Clutter, and J.E. Bethune
1972. Simulation as a tool in forest resources management. 51st Annu. Meet., Appalachian Sect., Soc. Am. For. Proc. 1972: 19-28 Durham, N. C. (SE)

Maple syrup production

934. Blum, B.M.
1972. Relationships between tree characteristics and maple sugar yields. Natl. Maple Syrup Dig. 11(3):15, 17, 19. (NE)

935. Gabriel, W.J., R.S. Walters, and D.W. Seegrist
1972. The relationship between sap-flow rate and sap volume in dormant sugar maples. USDA For. Serv. Res. Note NE-153, 4 p. (NE)

936. Gibbs, C.B., and H.C. Smith
1973. Cambial dieback and taphole closure in sugar maple after tapping. USDA For. Serv. Res. Note NE-155, 4 p. (NE)

937. Ruth, R.H., and J.C. Underwood
1971. Maple syrup in Oregon. Proc. Eighth Conf. Maple Prod. Oct. 19-20 Boyne Falls, Mich. ARS 73-73 (Abstr.). (PNW)

938. Ruth, R.H., J.C. Underwood, C.E. Smity, and H. Y. Yang
1972. Maple syrup production from bigleaf maple. USDA For. Serv. Res. Note PNW-181. (PNW)

939. Smith, H.C., and A.G. Snow, Jr.
1972. Pumping during the fall season. Natl. Maple Syrup Dig. 11(2):10-11. (NE)

940. Smith, H.C., and A.G. Snow, Jr.
1972. Installing plastic tubing to collect sugar maple sap. USDA For. Serv. Res. Pap. NE-240, 16 p. (NE)

941. Walters, R.S., and H.C. Smith
1972. Taphole numbers and tubing size. Natl. Maple Syrup Dig. 11(3):10-13. (NE)

942. Walters, R.S., and H.C. Smith
1972. Fall tapping. Natl. Maple Syrup Dig. 11(2):12-13. (NE)

Naval stores

943. Clements, R.W.
1973. Wages earned, number of faces chipped, and production per man-day in the gum naval stores industry. USDA For. Serv. Res. Note SE-187, 4 p. (SE)

944. Gansel, C.R.
1971. Effects of several levels of inbreeding on growth and oleoresin yield in slash pine. Proc. Eleventh South. Conf. For. Tree Improv. 1971:173-177. (SE)

945. Roberts, D.R.
1972. Practical progress in mecahnization, gum collection. Nav. Stores Rev. 82(3):11, 18. (SE)

946. Zamski, E.
1972. Temperature and photoperiodic effects on xylem and vertical resin duct formation in *Pinus halepensis* Mill. Israel J. Bot. 21:99-107. (WO)

947. Zamski, E., and A. Fahn
1972. Observations on resin secretion from isolated portions of resin ducts of *Pinus haleporis* Mill. Israel Journal of Botany 21: 35-38. (WO)

Distribution and classification of forest trees

948. Griffin, J.R., and W.B. Critchfield
1972. The distribution of forest trees in California. USDA For. Serv. Res. Pap. PSW-82, 114 p. (PSW)

949. Little, E.L., Jr.
1972. New tree species from Esmeraldas, Ecuador (concluded). Phytologia, 25(2):59-64. (WO)

950. Liu, T.
1971 (1972). A monograph of the genus *Abies*. Natl. Taiwan Univ., Taipei, Taiwan, China. 608 p. (WO)

951. Steinhoff, R.J.
1972. White pines of western North America and Central America. USDA Misc. Publ. 1221, p. 215-232. (INT)

Inherent variation

952. Becker, W.A., and M.A. Marsden
1972. Estimation of heritability and selection gain for blister rust resistance in western white pine. In Biology of Rust Resistance in Forest Trees. USDA Misc. Publ. 1221, p. 397-409. (INT)

953. Bey, C.F.
1972. Trends in growth of black walnut originating in various geographic areas. North. Nut Grow. Assoc. Annu. Rep. 62:83-86. (NC)

954. Bey, C.F.
1972. Leaf flush on black walnut at several midwest locations. In 19th Northeastern Forest Tree Improvement Conference Proceedings, University of Maine, August 2-4, 1971, p. 47-51. (NC)

955. Bramlett, D.L.
1971. Correlations between reproductive and vegetative growth in a 6-year-old Virginia pine plantation. USDA For. Serv. Res. Pap. SE-88, 6 p. (SE)

956. Campbell, R.K.
1972. Genetic variability in juvenile height growth of Douglas-fir. Silvae Genet. 21:126-129. (PNW)

957. Carl, C.M., Jr., and H.W. Yawney
1972. Multiple seedlings in *Acer saccharum*. Bull. Torrey Bot. Club 99:142-144. (NE)

958. Copes, D.L.
1972. Inheritance of megastrobili color in Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco). Can. J. Bot. 50(10): 2045-2048. (PNW)

959. Coyne, J.F., and G.C. Keith
1972. Geographic survey of monoterpenes in loblolly and shortleaf pines. USDA For. Serv. Res. Pap. SO-79, 12 p. (SO)

960. Franklin, E.C.
Some perspectives on progeny testing in a recurrent seelction program for tree improvement. Working Party on Progeny Test, IUFRO, Macon, Ga., Oct. 25-27, 1972 (Proc. in press). (SO)

961. Gabriel, W.J.
1972. One-parent progeny test of sugar maple selected for superior sap-sugar production—early results. Proc. 19th Northeast. For. Tree Impr. Conf. p. 12-16. (NE)

962. Gowin, T.
1972. Development of apical buds of Scots pine (*Pinus silvestris* L.) seedlings of different provenience. Ekologia Polska 20:772-779. (WO)

963. Gregory, R.A., and J.A. Romberger
1972. The shoot apical ontogeny of the *Picea abies* seedling. I. Anatomy, spical dome diameter, and plastochron duration. Am. J. Bot. 59: 587-597. (NE)

964. Gregory, R.A., and J.A. Romberger
1972. The shoot apical ontogeny of the *Picea abies* seedling. II. Growth rates. Am. J. Bot. 59: 598-606. (NE)

965. Hejnowicz, Z.
1972. The orientation of the plane of cell division in cambia of trees and factors controlling it. Final Tech. Rep. PL-480 Res. Proj. E21-FS-40, 73p. (WO)

966. Jeffers, R.M.
1972. Cone characteristics and seed yield in jack pine. Proc. Northeast. For. Tree Improv. Conf. 19: 35-43. (NC)

967. Johnson, L.C., and L.C. Saylor
1972. El Dorado pine: An aneuploid Monterey pine cultivar. J. Hered. 63(5): 293-296. (PSW)

968. Lafarge, T.
1973. Relationships among 3rd-, 5th, and 15th-year measurements in a study of stand variation of loblolly pine in Georgia. IUFRO, Working Party on Progeny Testing, Macon, Ga., Oct. 1972. (SE)

969. Mohn, C.A., and W.K. Randall
1971. Inheritance and correlation of growth characters in *Populus deltoides*. Silvae Genet. 20:182-184. (SO)

970. Nalborczyk, E., B. Zelawska, and M. Kolakowska
1971. Effects of X-rays on seed germination and growth of Scots pine (*Pinus sylvestris* L.) seedlings of different provenience. Acta Soc. Bot. Pol. 40:403-412. (WO)

971. Namkoong, G., R.A. Usanis, and R.R. Silen
1972. Age-related variation in genetic control of height growth in Douglas-fir. Theor. and Appl. Genet. 42:151-159. (PNW)

972. Nienstaedt, H., and A. Tiech
1972. The genetics of white spruce. USDA For. Serv. Res. Pap. WO-15, 24 p. (NC)

973. Schultz, R.P.
1972. Inherent vigor influences growth of slash pine more than intensive cultural treatments. Silvae Genetica 21(1-2): 26-28. (SE)

974. Sluder, E.R.
1973. Open-pollinated progenies of slash pine: their performance at fifteen years and the relationships among third-year data and eight- and fifteenth-year volumes per plot. IUFRO Working Party on Progeny Testing, Macon, Ga., Oct. 1972. (SE)

975. Sluder, E.R.
1972. Variation in specific gravity of yellow-poplar in the Southern Appalachians. Wood Sci. 5:132-138. (SE)

976. Whitmore, J.L.
1971. *Cedrela* provenance trial in Puerto Rico and St. Croix: nursery phase assessment. Turrialba 21:343-349. (ITF)

977. Wilkinson, R.C., and J.W. Hanover
1972. Geographical variation in the monoterpene composition of red spruce. Phytochem. 11:2007-2010. (NE)

978. Winstead, J.E.
1972. Fiber tracheid length and wood specific gravity of seedlings as ecotypic characters in *Liquidambar styraciflua* L. Ecol. 53:165-172. (WO)

979. Zelawski, W., and E. Nalborczyk
1971. Productivity of photosynthesis in Scots pine (*Pinus sylvestris* L.) grown from seed irradiated by X-rays. Acta Soc. Bot. Pol. 40:413-421. (WO)

980. Zelawski, W., T. Gowin, and A. Lotocki
1972. Preliminary study of the ecotype differentiation on three plantations of Scots pine (*Pinus sylvestris* L.) in Poland. Ekol. Pol. 20:763-770. (WO)

Insect-disease resistance

981. Bingham, R.T.
1972. Taxonomy, crossability, and relative blister rust resistance of 5-needled white pines. In Biol. Rust Resist. For. Trees, USDA Misc. Publ. 1221, p. 271-280. (INT)

982. Bingham, R.T., R.J. Hoff, and R.J. Steinhoff
1972. Genetics of western white pine. USDA For. Serv. Res. Pap. WO-12, 18 p. (WO)

983. Boyer, W.D.
1972. Brown-spot resistance in natural stands of longleaf pine seedlings. USDA For. Serv. Res. Note SO-142, 4 p. (SO)

984. Demeritt, M.C., Jr., W.M. Chang, J.D. Murphy, and H.D. Gerhold
1972. Selection system for evaluating resistance of scotch pine seedlings to ozone and sulfur dioxide. NEFTIC Proc. 19:87-97. (NE)

985. Dinus, R.J.
1972. Testing for fusiform rust resistance in slash pine. In Biol. Rust Resist. For. Trees USDA, Misc. Pub. 1221, p. 331-339. (SO)

986. Garrett, P.W.
1972. Resistance of eastern white pine (*Pinus strobus* L.) provenances to the white-pine weevil (*Pissodes strobi* Peck.). Silvae Benet. 21:119-121. (NE)

987. Garrett, P.W.
1973. Geographic variation in resistance to white-pine weevil (*Pissodes strobi* Peck) by eastern white pine (*Pinus strobus* L.). Can. Entomol. 105(3). (NE)

988. Hare, R.C.
1972. Physiology and biochemistry of resistance to pine rusts. In Biol. Rust Resist. in For. Trees USDA, Misc. Publ. 1221, p. 465–478. (SO)

989. Kinloch, B.B. Jr.
1972. Genetic variation in resistance to *Cronartium* and *Peridermium* rusts in hard pines. In Biol. Rust Resist. in For. Trees USDA, Misc. Pub. 1221, p. 445–463. (PSW)

990. Snyder, E.B., and H.J. Derr
1972. Breeding longleaf pines for resistance to brown spot needle blight. Phytopath. 62:325–329. (SO)

Tree breeding methodology

991. Bey, C.F., A.S. Mickelson, and M. Gerardo
1972. Balck walnut seedling seed orchard development—a case history. Proc. Northeast. Area Nurserymen's Conf., p. 48–58. (NC)

992. Bramlett, D.L., and T.W. Popham
1971. Model relating unsound seed and embryonic lethal alleles in self-pollinated pines. Silvae Genet. 20: 192–193. (SE)

993. Clausen, K.E.
1973. The effect of pollen irradiation on reproductive capacity, seedling growth, and variation of *Betula nigra*. I. Seed yield, germination, and germinant abnormalities. Radia. Bot. 13: 37–54. (NC)

994. Conkle, M.T.
1972. Forest Tree Improvement in California—1970. USDA For. Serv. Res. Note, PSW-275, 4 p. (PSW)

995. Conkle, M.T.
1972. Analyzing genetic diversity in conifers, isozyme resolution by starch gel electrophoresis. USDA For. Serv. Res. Note, PSW-264, 5 p. (PSW)

996. Copes, D.L.
1973. Inheritance of graft compatibility in Douglas-fir. Bot. Gaz. 134: 49–52. (PNW)

997. Franklin, E.C., and A.E. Squillace
Short-term progeny tests and second generation breeding in slash pine. Sixth TAPPI For. Biol. Conf., Appleton, Wisc., May 1-3, 1972. Canad. J. For. Sci. (In press). (SE)

998. Gibbs, C.B.
1973. The New England Spruce-Fir Seed Orchard Program. In Joint Proceedings of the 10th Lake States Forest Tree Improv. Conf. and 7th Central States For. Tree Improv. Conf., September 22–24, 1971, Gen. Tech. Rep. NC-3, p. 53–56. (NE)

999. Jain, M.K., and K.K. Nanda
1972. Effect of temperature and some antimetabolites on the interaction effects of auxin and nutrition in rooting etiolated stem segments of *Salix tetrasperma*. Physiol. Plant. 27: 169–172. (WO)

1000. Johnson, L.C., and S.M.F. Ward
1972. Male cone production in *Metasequoia glyptostreboides* growing at the Dominican College of San Rafael, California. Calif. Hort. J. 33: 98–100, 119. (PSW)

1001. Kaeiser, M., and D.T. Funk
1972. Structural changes in walnut grafts. Nut Grow. Assoc. Annu. Rep. 62: 90–94. (NC)

1002. Kamienska, A.
1972. Endogenous auxins and growth inhibitors in the developing inflorescences and infructescences of black poplar (*Populus nigra* L.). Acta Soc. Bot. Pol. 41:393–400. (WO)

1003. Konar, R.N.
1972. Tissue and cell culture of pines and allied conifers. PL-480 Final Tech. Rep. A7-FS-36, 35 p. (WO)

1004. McAlpine, R.G., and P.P. Kormanik
1971. Rooting cuttings from select yellow-poplar trees. Proc. Eleventh South. Conf. For. Tree Improv., p. 241–243. (SE)

1005. McAlpine, R.G., and P.P. Kormanik
1972. Rooting yellow-poplar cuttings from girdled trees. USDA For. Serv. Res. Note SE-180, 4 p. (SE)

1006. Marquis, D.A.
1973. Factors affecting financial returns from hardwood tree improvement. J. For. 71:79–83. (NE)

1007. Michniewicz, M., and E. Galoch
1972. Dynamics of endogenous inhibitor of abscisic acid properties in the development of buds, newly formed shoots and adventitous roots of willow cuttings (*Salix viminalis* L.). Bull. Acad. Pol. Sci. Ser. Biol. 20: 333–337. (WO)

1008. Kopcewicz, J., M. Michniewicz, and K. Kriesel
1972. Dynamics of gibberellin-like substances in the development of buds, newly formed shoots and adventitous roots of willow cuttings. (*Silix viminalis* L.). Acta Soc. Bot. Pol. 41:301–310. (WO)

1009. Namkoong, G.
1972. Persistence of variances for stochastic, discrete-time, population growth models. Theor. Pop. Biol. 3:507–518. (WO)

1010. Nanda, K.K.
1972. Investigations on the use of auxins in vegetative reproduction of forest plants. PL-480 Final Tech. Rep. A7-FS-11, 215 p. (WO)

1011. Nanda, K.K., and M.K. Jain
1972. Utilization of sugars and starch as carbon sources in the rooting of etiolated stem segments *Populus nigra*. New Phytol. 71:825–828. (WO)

1012. Rudolph, T.D.
1972. Mutation breeding in forest trees. In Mutation Breeding Workshop, January 17–18, 1972, Univ. of Tenn. Agric. Campus, Knoxville, Tenn. (NC)

1013. Schmidtling, R.C.
1971. Cultivating and fertilizing stimulate precocious flowering in loblolly pines. Silvae Genet. 20: 220–221. (SO)

1014. Schreiner, E.J.
1972. Procedure for selection of hybrid poplar clones for commercial trials in the Northeastern Region. NEFTIC Proc. 19: 108–116. (NE)

1015. Snyder, E.B.
1972. Glossary for forest tree improvement workers. USDA For. Serv. South. For. Exp. Stn., 22 p. (SO)

1016. Snyder, E.B.
1972. Five-year performance of self-pollinated slash pines. For. Sci. 18: 246. (SO)

1017. Sorensen, F.C.
1972. The seed orchard tree as a pollen sampler: A model and example. USDA For. Serv. Res. Note PNW-175, 11 p. (PNW)

1018. Squillace, A.E., K.W. Dorman, and R.E. McNees
1972. Breeding slash pine at Olustee, Florida: A success story. Agric. Sci. Review 10(3): 25-32. (SE)

1019. Steinhoff, R.J., and R.J. Hoff
1972. Chilling requirements for breaking dormancy of western white pine seedlings. USDA For. Serv. Res. Note INT-153, 6 p. (INT)

1020. Tucovic, A.
1972. Develop breeding techniques for oaks. PL-480 Final Tech. Rep. E30-FS-6, 105 p. (WO)

1021. Tucovic, A.
1972. Breed walnuts for high quality wood products. PL-480 Final Tech. Rep. E30-FS-7, 80 p. (WO)

1022. Valanne, T.
1972. Colchicine effects and colchicine-induced polyploidy in *Betula*. Annu. Acad. Fenn., Biologica IV: 191, 28 p. (WO)

1023. Vidakovic, M., and B. Jurkovic-Bevilacqua
1970. Observation on the ovule development following cross pollination between Austrian and Scots pines using irradiated and non-irradiated pollen. Proc. Working Group on the Sexual Reprod. of For. Trees. Varparanta, Finland, 8 p. (WO)

Utilization potential and processing of wood

1024. Echols, R.M.
1972. Product suitability of wood determined by density gradients across growth rings. USDA For. Serv. Res. Note PSW-273, 6 p. (PSW)

1025. Hejnowicz, Z., and J.A. Romberger
1973. Migrating cambial domains and the origin of wavy grain in xylem of broadleaved trees. Am. J. Bot. 60:209-222. (WO)

1026. Isebrands, J.G.
1972. Proportion of wood elements within eastern cottonwood. Wood Sci. 5:139-146. (NC)

1027. Isebrands, J.G., and D.W. Bensend
1972. Incidence and structure and gelatinous fibers within rapid growing eastern cottonwood. Wood and Fiber 4:61-71. (NC)

Miscellaneous

1028. Bingham, R.T.
1972. Station publications in forest genetics and related fields. USDA For. Serv. Res. Note INT-157, 10 p. (INT)

1029. Darwin, W.N. Jr.
1972. Tupelo (*Nyssa* spp.). USDA, Am. Woods FS-269, 8 p. (SO)

1030. Franklin, J.F., R.E. Jenkins, and R.M. Romancier
1972. Research Natural Areas: Contributors to environmental quality programs. J. Environ. Qual. 1:133-139. (PNW)

1031. Haissig, B.E.
1972. Near total recovery of small tissue samples after milling. For. Sci. 18:261-272. (NC)

1032. Jones, J.R., and D.C. Markstrom
1972. Aspen an American wood. USDA, Am. Woods FS-217, 8 p. (RM)

1033. Kallio, E., and R.M. Godman
1973. American basswood an American wood. USDA, Am. Woods FS-219, 8 p. (NC)

1034. Krochmal, A., L. Wilken, and M. Chien
1972. Lobeline content of four Appalachian lobelias. Lloydia 35:303-304. (SE)

1035. Krochmal, A., L. Wilken, and M. Chien
1972. Plant and lobeline harvest of *Lobelia inflata* L. Econ. Bot. 26:216-220. (SE)

1036. Landt, E.F., and R.E. Phares
1973. Black walnut an American wood. USDA, Am. Woods FS-270, 7 p. (NC)

1037. Stein, W.J.
1972. 1971 listing of selected publications on reforestation. IN West. For. Conserv. Assoc. West. Reforestation Coord. Comm. Proc. 1971:61-71. (PNW)

1038. Wodzicki, T.J., and S. Zajaczkowski
1970. Methodical problems in studies on seasonal production of cambial xylem derivatives. Acta Soc. Bot. Pol. 39:509-520. (WO)

FOREST ECONOMICS AND MARKETING RESEARCH

FOREST SURVEY

Forest inventory

1039. Barnard, J.E., and T.M. Bowers
1973. A preview of Delaware's timber resource. USDA Forest Serv. Resour. Note NE-159, 3 p. (NE)

1040. Barnard, J.E., and C.E. Mayer
1972. Penn's Sylvania—the current forest resources of Pennsylvania. Pennsylvania Forests 63(3): 59-64. (NE)

1041. Cathey, R.A.
1972. Forest statistics for Central Georgia, 1972. USDA For. Serv. Resour. Bull. SE-22, 34 p. (SE)

1042. Christopher, J.F., and H.S. Sternitzke
1972. Data and services available from the forest survey and future plans. Paper presented at the Southwest Tech. Div. Am. Pulp. Assn., 4 p. (SO)

1043. Hedlund, A., and J.M. Earles
1972. Forest statistics for central Alabama counties. USDA For. Serv. Resour. Bull. SO-37, 23 p. (SO)

1044. Hedlund, A., and J.M. Earles
1972. Forest statistics for north Alabama counties. USDA For. Serv. Resour. Bull. SO-34, 13 p. (SO)

1045. Hedlund, A., and J.M. Earles
1972. Forest statistics for southwest Alabama counties. USDA For. Serv. Resour. Bull. SO-33, 13 p. (SO)

1046. Knight, H.A.
1972. Forest statistics for north central Georgia. USDA For. Serv. Resour. Bull. SE-24, 34 p.

1047. Knight, H.A.
1972. Forest resources in Southeast survey monitors from decades of change. South. Lumberman 225(2800): 147-159. (SE)

1048. McClure, J.P.
1972. Customized forest information retrieval. For. Farmer 31(10): 6-7. (SE)

1049. Murphy, P.A.
1972. Forest resources of Tennessee. USDA For. Serv. Resour. Bull. SO-35, 33 p. (SO)

1050. Murphy, P.A.
1972. Timber volume increases in Tennessee. For. Farmer 31(12): 13-14. (SO)

1051. Ostrom, A.J.
1972. Timber volume in Wisconsin counties. USDA For. Serv. Res. Note NC-145, 4 p. (NC)

1052. Oswald, D.D.
1972. Timber resources of Mendocino and Sonoma counties, California. USDA For. Serv. Resour. Bull. PNW-40, 75 p. (PNW)

1053. Spencer, J.S., Jr., and H.W. Thorne
1973. Wisconsin's 1968 timber resource—a perspective. USDA For. Serv. Resour. Bull. NC-15, 80 p. (NC)

1054. Sternitzke, H.S.
1972. Updated Southern forest resource situation. In the 11th Auburn Forest Forum Proc., p. 44-47, Auburn Univ. (SO)

1055. Sternitzke, H.S.
1972. Bald cypress: endangered or expanding species? Econ. Bot. 26: 130-134. (SO)

1056. Sternitzke, H.S., and J.F. Christopher
1972. Southern timber supply trends and outlook. For. Prod. J. 22(7): 13-16. (SO)

1057. U.S. Department of Agriculture Forest Service
1972. Forest statistics for the United State, by state and region, 1970. USDA For. Serv. 96 p. (WO)

Timber utilization and production

1058. Bellany, T.R.
1972. Southern pulpwood production, 1971. USDA For. Serv. Resour. Bull. SE-23, 20 p. (SE)

1059. Beltz, R.C.
1972. Midsouth pulpwood movement. Pulpwood Prod. 20(4): 14, 16. (SO)

1060. Bertelson, D.F.
1972. Alabama's forest industries: Preview of a survey report. Ala. For. Prod. 15(7): 69-70. (SO)

1061. Bertelson, D.F.
1972. Tennessee forest industries modernize. For. Farmer 31(5): 6-7. (SO)

1062. Bertelson, D.F.
1972. Alabama forest industries. USDA For. Serv. Resour. Bull. SO-36, 29 p. (SO)

1063. Bertelson, D.F.
1972. Midsouth pulpwood prices, 1971. Southern Pulp and Paper Manuf. 35(10-A): 15. (SO)

1064. Blyth, J.E.
1973. Lake States pulpwood production dips to 4 million cords in 1972. USDA For. Serv. Res. Note NC-148, 3 p. (NC)

1065. Blyth, J.E., and R. Massengale
1972. Missouri's primary forest products output and industries, 1969. USDA For. Serv. Resour. Bull. NC-16, 15 p. (NC)

1066. Bones, J.T., and N.P. Kingsley
1972. Future use of primary wood fiber for papermaking in the Northeast. Pulp and Paper, April 1972: 100-101. (NE)

547-679 O - 74 - 7

1067. Bones, J.T., and C.J. Lohr
1972. Primary wood-product industries of Kentucky—1969. USDA For. Serv. Resour. Bull. NE-15, 38 p. (NE)

1068. Bones, J.T., and J.K. Sherwood
1972. Primary wood-product industries of Pennsylvania - 1969. USDA For. Serv. Resour. Bull. NE-27, 34 p. (NE)

1069. Choate, G.A.
1972. Flows of mill residues to pulp and board manufactures, Washington, Oregon, and California, 1968. USDA For. Serv. Resour. Bull. PNW-39, 49 p. (PNW)

1070. Gardner, R.B., and D.W. Hann
1972. Utilization of lodgepole pine logging residues in Wyoming increases fiber yield. USDA For. Serv. Res. Note INT-160, 4 p. (INT)

1071. Hahn, J.T.
1973. Local net timber volume equations for Wisconsin. USDA For. Serv. Res. Note NC-149, 4 p. (NC)

1072. Hutchin, C.C.
1972. Pulpwood prices in the Southeast, 1971. USDA For. Serv. Res. Note SE-178, 2 p. (SE)

1073. Setzer, T.S.
1973. Logging residues on harvesting operations in South Dakota, Wyoming, Utah, and Colorado. USDA For. Serv. Res. Paper INT-135, 12p. (INT)

1074. Wall, B.R.
1972. Oregon timber harvest. USDA For. Serv. Resour. Bull. PNW-43, 2p. (PNW)

1075. Wall, B.R.
1972. 1971 Washington timber harvest. USDA For. Serv. Resour. Bull. PNW-41, 2 p. (PNW)

1076. Wall, B.R.
1972. Relationship of log production in Oregon and Washington to economic conditions. USDA For. Serv. Res. Pap. PNW-147, 13 p. (PNW)

1077. Wall, B.R.
1972. Log production in Washington and Oregon—an historical perspective. USDA For. Serv. Resour. Bull. PNW-42, 89 p. (PNW)

Survey techniques

1078. Aldrich, R.C., and W.J. Greentree
1972. Forest and nonforest land classification using aircraft and space imagery. In Monitoring forest land from high altitude and from space, final report. Available in microfiche from NASA, Wash., DC. (PSW)

1079. Aldrich, R.C., J. von Mosch, and W.J. Greentree
1972. Projection-viewer for microscale aerial photography. USDA For. Serv. Res. Note PSW-277, 5 p. (PSW)

1080. Dana, R.W.
1973. Digital sensitometry of color infrared film as an aid to pattern recognition studies. In Proc. 2nd annu. remote sensing of earth resources Conf., Univ. Tenn. Space Instit., Tullahoma, Tenn., March 26-28, 1973. (PSW)

1081. Dana, R.W., and B. Marx
1972. Calibration of focal plane shutters. In Monitoring Forest Land from High Altitudes and from Space, Final Report. Available in microfiche from NASA, Wash., DC. (PSW)

1082. Dana, R.W., and N.X. Norick
1972. Mathematical modeling of film characteristic curves. In Monitoring Forest Land from High Altitudes and from Space, Final Report. Available in microfiche from NASA, Wash., DC. (PSW)

1083. Hazard, J.W., and J.M. Berger
1972. Volume tables versus dendrometers for forest surveys. J. For. 70(4): 216-219. (PNW)

1084. Heller, R.C.
1972. Remote sensing in forestry—promises and problems. In Proc. Soc. Am. For. Nat. Conv., Hot Springs, Ark., Oct. 1-5, 1972. (PSW)

1085. Heller, R.C., and R.V. Bega
1973. Detection of forest diseases by remote sensing. J. For. 71(1): 18-21. (PSW)

1086. Heller, R.C., B. Spada, and A.M. Woll
1972. Remote sensing in resource evaluation, planning, protection and management. Proc. 7th World Forestry Congress, Buenos Aires, Argentina, Oct. 4-18, 1972, 11 p. (PSW)

1087. Hildebrandt, G., and R.C. Heller
1973. The implications of remote sensors for forestry research and practices. Proc. XV IUFRO Congress, Gainsville, Fl., Mar. 14-20, 1971. (PSW)

1088. Howard, J.O., and F.R. Ward
1972. Measurement of logging residue—alternative applications of the line intersect method. USDA For. Serv. Res. Note PNW-183, 8 p. (PNW)

1089. MacLean, C.D.
1972. Improving inventory volume estimation by double sampling on aerial photographs. J. For. 70: 748-749. (PNW)

1090. MacLean, C.D.
1972. Photo stratification improves Northwest timber volume estimates. USDA For. Serv. Res. Pap. PNW-150, 10 p. (PNW)

1091. Norick, N.X., and M. Wilkes
1972. Classification of land use by automated procedures. In Monitoring Forest Land from High Altitudes and from Space, Final Report. Available in microfiche from NASA, Wash. DC. (PSW)

1092. Pope, R.B., B. Payondeh, and D.P. Paine
1972. Photo plot bias. USDA For. Serv. Res. Pap. PNW-145, 8 p. (PNW)

1093. Van Hooser, D.D.
1972. Evaluation of two-stage 3P sampling for forest surveys. USDA For. Serv. Res. Pap. SO-77, 9 p. (WO)

1094. Van Hooser, D.D., and R.C. Biesterfeldt
1972. 3P sampling to update forest surveys. South. Lumberman 225(2800): 120-122. (SO)

1095. Weber, F.P.
1973. The use of airborne spectrometers and multispectral scanners for previsual detection of ponderosa pine trees under stress from insects. In Monitoring Forest Land from High Altitude and from Space, Final Report. Available in microfiche from NASA, Washington, DC. (PSW)

1096. Weber, F.P., R.C. Aldrich, F.G. Sadowski, and F.J. Thomson
1972. Land use classification in the Southeastern forest region by multispectral scanning and computerized mapping. In Monitoring Forest Land from High Altitude and from Space, Final Report. Available in microfiche from NASA, Wash., DC. (PSW)

1097. Weber, F.P., T.H. White, and R.C. Heller
1973. Trend and spread of bark beetle infestations in the Black Hills. In Monitoring Forest Land from High Altitudes and from Space Final Report. Available in microfiche from NASA, Wash., DC. (PSW)

FOREST ECONOMICS RESEARCH

Methods of financial evaluation and planning

1098. Leuschner, W.A.
1972. A new approach for projecting long term timber supply and inventory. For. Prod. J.22(5): 49-53. (NC)

1099. Lundgren, A.L., and E.F. Thompson
1972. Uncertainty in forestry investment decisions regarding timber growing. Va. Polytechnic Inst. and State Univ., Blacksburg, Va. Publ. FWS-1-72, 31 p. (NC)

1100. McCay, R.E., and P.S. DeBald
1973. A probability approach to sawtimber tree-value projections. USDA For. Serv. Res. Pap. NE-254, 8 p. (NE)

1101. Randall, R.M.
1972. An operations research approach to Douglas-fir thinning. USDA For. Serv. Res. Pap. PNW-148, 23 p. (PNW)

1102. Schweitzer, D.L., R.W. Sassaman, and C.H. Schallau
1972. Allowable cut effect—some physical and economic implications. J. For. 70(7): 415-418. (PNW)

Timber growing economics

1103. Anderson, W.C.
1972. Southern forestry investments in an era of environmental concern. For. Prod. J. 22(6): 14-16. (SO)

1104. Anderson, W.C.
1972. Growing pine saw-timber fast. South. Lumberman 225(2800): 156-158. (SO)

1105. Anderson, W.C.
1972. Economically oriented production functions for slash and loblolly pine plantations. USDA For. Serv. Res. Note SO-144, 3 p. (SO)

1106. Gansner, D.A., J.E. Barnard, and S.F. Gingrich
1972. Identifying regional opportunities for accelerated timber management. USDA For. Serv. Res. Pap. NE-251, 6 p. (NE)

1107. Guttenberg, S.
1972. To lease or not to lease your timberlands. For. Farmer 31(7): 41-42. (SO)

1108. Guttenberg, S.
1971. Timberland leases by southern forest industry. Soc. Am. For. South. Reg. Tech. Conf. Proc. 1971: 147-151. (SO)

1109. Larsen, D.N., and D.A. Gansner
1972. Explaining the forest product selling behavior of private woodland owners. USDA For. Serv. Res. Pap. NE-257, 4 p. (NE)

1110. Leuschner, W.A.
1972. Aspen supplies in the Lake States. Am. Pulpwood Assoc. Tech. Pap., p. 73-76. (NC)

1111. Leuschner, W.A.
1972. Projecting the aspen resource in the Lake States. USDA For. Serv. Res. Pap. NC-81, 32 p. (NC)

1112. Leuschner, W.A.
1972. Projections of inventories in the Lakes States. USDA For. Serv. Gen. Tech. Rep. NC-1., p. 10-15. (NC)

1113. McCauley, O.D., and D.A. Marquis
1972. Investment in precommercial thinning of northern hardwoods. USDA For. Serv. Res. Pap. NE-245, 12 p. (NE)

1114. McCauley, O.D., and G.R. Trimble
1972. Forestry returns evaluated for uneven-aged management in two Appalachian woodlots. USDA For. Serv. Res. Pap. NE-244, 12 p. (NE)

1115. Mendel, J.J., T.J. Grisez, and G.R. Trimble
1973. The rate of value increase for sugar maple. USDA For. Serv. Res. Pap. NE-250, 19 p. (NE)

1116. Miller, R.L., and F.R. Lason
1973. A cost analysis of ponderosa pine watershed. USDA For. Serv. Res. Note RM-231, 7 p. (RM)

1117. Payne, B.R.
1972. Accelerated roadbuilding on the North Umpqua—an economic analysis. USDA For. Serv. Res. Pap. PNW-137, 32 p. (PNW)

1118. Sassaman, R.W., J.W. Barrett, and J.G. Smith
1972. Economics of thinning stagnated ponderosa pine sapling stands in the pine-grass areas of central Washington. USDA For. Serv. Res. Pap. PNW-144, 17 p. (PNW)

1119. Sassaman, R.W., E. Holt, and K. Bergsvik
1972. User's manual for a computer program for simulating intensively managed allowable cut. USDA For. Serv. Gen. Tech. Rep. PNW-1, 50 p. (PNW)

1120. Schweitzer, D.L., and R.W. Sassaman
1973. Harvest volume regulation affects investment value. For. Ind. 100(3): 35. (PNW)

1121. Siegel, W.C.
1972. Environmental law and forest management. J. For. 70: 682-686. (SO)

1122. Siegel, W.C.
1972. Timber loans by U.S. life insurance companies. USDA For. Serv. Res. Pap. SO-73, 20 p. (SO)

Multiple-use economics

1123. Alston, R.M.
1972. FOREST—Goals and decisionmaking in the Forest Service. USDA For. Serv. Res. Pap. INT-138, 84 p. (INT)

1124. Duran, G., and H.F. Kaiser
1972. Range management proactices investment costs, 1970. USDA Agric. Handbk. 435, 38 p. (WO)

1125. Irland, L.C., and J.W. Tarver
1972. Problems of water resource development in the Gulf Coast estuarine zone. Proc. Symp. on Uncertainties in Hydrologic and Water Resour. Syst. 1972: 121-126. (SO)

1126. Kaiser, H.F., K. DeBower, R. Lockard, and J.W. Putman
1972. Forest-range environmental production analytical system (FREPAS). USDA Handbk. 430, 211 p. (WO)

1127. Leary, R.A., and K.E. Skog
1972. A computational strategy for system identification in ecology. Ecol. 53(5): 969-973. (NC)

1128. Navon, D.I., and H.W. Anderson
1972. Modern techniques in planning the management of wildland resources. Proc. Joint FAO/USSR Int. on For. Influences and Watershed Manage. 1972: 359-368. (PSW)

1129. O'Connell, P.F.
1972. Valuation of timber, forage, and water from National Forest lands. Annu. Reg. Sci. 6(2): 1-14. (RM)

1130. O'Connell, P.F., and H.E. Brown
1972. Economics of chaparral management in the Southwest. Proc. Am. Water Resour. Assoc. 1972: 260-266. (RM)

1131. O'Connell, P.F.
1972. Use of production functions to evaluate multiple-use treatments on forested watersheds. Water Resour. Res. 8(5): 1188-1198. (RM)

1132. Sassaman, R.W.
1972. Economic returns from planting forage in National Forests. J. For. 70(8): 487-488. (PNW)

Impacts on forest industry and regional economies

1133. Boster, R.S., and W. Martin
1972. The value of primary versus secondary data in interindustry analysis: a study in the economics of the economic models. Annu. Reg. Sci. 6(2): 35-44. (RM)

1134. Dutrow, G.F.
1972. Shift-share analysis of southern forest industry. For. Prod. J. 22(12): 10-14. (SO)

1135. Granskog, J.E.
1972. Timber producers associations—new bargaining force? For. Ind. 99(5): 30-31. (SO)

1136. Guttenberg, S.
1972. Managerial ingenuity is key to southern pine pulpwood supply. For. Ind. 99(7): 83-84. (SO)

1137. Irland, L.C.
1972. Labor trends in southern forest industries, 1950 to 1969. USDA For. Serv. Res. Pap. SO-81, 20 p. (SO)

1138. Kaiser, H.F.
1972. Multiregional input-output model for forest resource analysis. For. Sci. 18: 46-53. (SO)

1139. Kaiser, H.F., and W.C. Anderson
1972. Gains from joint lumber and chip production at pulpwood chipping yards. J. For. 70: 230-234. (SO)

FOREST PRODUCTS MARKETING RESEARCH

Marketing underused species and residues

1140. Applefield, M.
1972. The challenge of marketing fine wood residues. Ala. For. Prod. 15(7): 25-28, 30. (SE)

1141. Applefield, M.
1973. Researching wood residues. South. Lumberman 226(2805): 15-16. (SE)

1142. Austin, J.W.
1973. Price trends in fiberwood used by mills in the Northwest. Pulp and Paper 47(3). (PNW)

1143. Cuppett, D.G., D.M. Emanuel, and P.S. Wysor
1972. Evaluation of a new sawing system for low-grade hardwood logs. South. Lumberman 225(2799): 11-14. (NE)

1144. Harpole, G.B.
1972. The local markets for hardwood lumber products produced from forests of Maui, Hawaii. Pac. Southwest For. and Range Exp. Stn., Berkeley, CA, 13 p. (PSW)

1145. Sarles, R.L.
1973. Bark—a Cinderella story. Bark mulches and soil conditioner: new products for the wood industry. North. Logger and Timber Processer 21(9): 14-15. (NE)

Supply, demand, and price analysis

1146. Applefield, M.
1972. Regulations affecting wood use in building construction. For. Prod. J. 22(4): 12-16. (SE)

1147. Bond, R.S.
1972. Labor supply—a critical problem in certain Northeastern primary wood-processing industries. North. Logger and Timber Processer 20(10): 18, 20-43. (NE)

1148. Bradley, D.P.
1972. Improve forest inventory with access data—measure transport distance and cost to markets. USDA For. Serv. Res. Pap. NC-82, 21 p. (NC)

1149. Bradley, D.P.
1972. General print routine. The Wang Laboratories Programmer 6(5): 17. (NC)

1150. Bradley, D.P., and F.E. Biltonen
1972. Economic operability—factors affecting harvest and transport costs. Proc. Aspen Symp. Gen. Tech. Rep. NC-1, p. 16-20. (NC)

1151. Brock, S.M., W.G. Hogan, and T.H. Pendleton
1972. Workmen's compensation laws: Their application to logging and sawmilling in the Northeast. Northeast J. For. 70(7): 408-410. (NE)

1152. Carpenter, E.M.
1972. Canadian forest products shipped into the North Central region. USDA For. Serv. Res. Pap. NC-83, 22 p. (NC)

1153. Darr, D.R.
1972. Production, prices, employment and trade in Northwest forest industries, fourth quarter 1971. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg., 55 p. (PNW)

1154. Darr, D.R.
1972. Production, prices, employment and trade in Northwest forest industries, first quarter 1972. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg., 29 p. (PNW)

1155. Gill, T.G., and R.B. Phelps
1972. Wood preservation statistics, 1971. USDA For. Serv. in cooperation with Am. Wood Preservers Assoc. 34 p. (WO)

1156. Glover, R.P., and B.G. Hansen
1972. The secondary forest industry of West Virginia—1972. W. Va. Dep. Nat. Resour., Resourc. Bull. (NE)

1157. Goho, C.D., and A.J. Martin
1973. Sawlog sizes: a comparison in two Appalachian areas. USDA For. Serv. Res. Note NE-160. (NE)

1158. Hair, D., and A.H. Ulrich
1972. The demand and price situation for forest products, 1971-72. USDA Misc. Pub. 1231, 88 p. (WO)

1159. Holt, E.
1972. Production, prices, employment and trade in Northwest forest industries, second quarter 1972. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg., 55 p. (PNW)

1160. Holt, E.
1972. Production, prices, employment and trade in Northwest forest industries, third quarter 1972. Pac. Northwest For. and Range Exp. Stn., Portland, Oreg., 54 p. (PNW)

1161. Host, J.R.
1973. Productivity in Montana logging operations. Symp. Proc. Planning and Decision Making as Applied to Forest Harvesting, Oreg. State Univ. (INT)

1162. Lawrence, J.D.
1972. Potential role of mobile homes in interregional housing production. Bus. and Econ. Dimensions 8(6): 8-14. (SE)

1163. Lawrence, J.D.
1972. Timber availability from private lands. Fla. For. Reporter 15(1): 2. (SE)

1164. Lawrence, J.D., and R.K. Jones
1972. Mobile home shipment patterns. Atlanta Econ. Rev. (4): 8-14. (SE)

1165. Lawrence, J.D., H.E. Dickerhoof, and C.A. Fasick
1972. Mobile home production in the South may cause changes in wood products marketing. South. Lumberman 225(2800): 115-116. (SE)

1166. Lege, F.O., and J.D. Lawrence
1972. Availability of stumpage for the open market. South. Lumberman 225(2800): 134-135. (SE)

1167. Lindell, G.R.
1972. The changing market for hardwood plywood stock panels. USDA For. Serv. Res. Pap. NC-78, 7 p. (NC)

1168. Lindell, G.R., and E. Kallio
1972. Paneling outlook appears good through 1974. Plywood and Panel XIII (3): 20-21, 25. (NC)

1168a. Lindell, G.R., and E. Kallio
1973. Use of paneling will increase in apartment construction. Plywood Panel XIII (8):23-24. (NC)

1169. Marcin, T.C.
1972. Projections of demand for housing by type of unit and region. USDA Agric. Hdbk. 428, 76 p. (NC)

1170. Martin, A.J.
1972. The aggressive skidder operator and his impact on logging costs. South. Lumberman 225(2800): 129. (NE)

1171. Miller, W.C.
1972. Distribution of parquet flooring during 1969. USDA For. Serv. Res. Pap. NE-218, 11 p. (NE)

1172. Sampson, G.R.
1972. Determining maximum stumpage values for wood product mills with alternative proceses. For. Prod. J. 22(8): 26-30. (SE)

1173. Sendak, P.E.
1972. Why the tariff failed. National Maple Syrup Digest. 2(2): 18-22. (NE)

1174. Sendak, P.E.
1972. The effect of the tariff on the maple industry. USDA For. Serv. Res. Note NE-148, 5p. (NE)

1175. Timson, F.G., and T.W. Church
1972. Appalachian loggers wrestle very few heavyweights! North. Logger and Timber Processer 29(10): 26, 28. (NE)

1176. Wright, M.G., and W.H. Reid
1972. Wood materials used in apartment construction. USDA For. Serv. Res. Pap. WO-16, 12 p. (WO)

Marketing—general

1177. Anderson, R.B., and W.C. Miller
1973. Wooden beverage cases cause little damage to bottle caps. USDA For. Serv. Res. Note NE-161. (NE)

1178. Applefield, M.
1972. The fire district scene from the Southern region. Southern Building, p. 13-14. (SE)

1179. Applefield, M.
1972. Use of fire zones in the Western region. Building Standards, Int'l. Conf. Bldg. Officials, bimonthly, 41(4): 6-9. (SE)

1180. Applefield, M.
1973. Fire limits statistics for cities in the Northeast. Int. Fire Chief, 38(8): 16-19. (SE)

1181. Applefield, M.
1973. Fire district use in North Central region cities. Fire J. 69(1): 28-31, 33. (SE)

1182. Baumgras, J.E.
1972. The third challenge—management's attitude. North. Logger and Timber Processer 21(2): 17, 18, 41. (NE)

1183. Church, T.W., Jr.
1972. The CHUBALL—a unique new device for selectively logging steep slopes. North. Logger and Timber Processer 21(2): 8, 9, 27, 33. (NE)

1184. Cooper, G.A., F.B. Calvin, R.D. Lindmark, and R.C. Schlesinger
1972. Better hardwoods—from seeds to sawlogs and mills to markets. South. Lumberman 225(2800): 111-114. (NC)

1185. Cuppett, D.G., and E.P. Craft
1972. Kiln drying of presurfaced 4/4 Appalachian oak. For. Prod. J. 22(6): 36-41. (NE)

1186. Kallio, E., and R. Bodman
1973. Basswood—an American wood. USDA For. Serv. Am. Woods FS-219. (NC)

1187. Kallio, E., and R.D. Lindmark
1972. Kentuckians fashion better living from handcrafts. News of Farmer Cooperatives 39(6): 12-14. (NC)

1188. Martin, A.J.
1972. MULGRES user's manual. Northeast. For. Exp. Stn., Upper Darby, PA, Unnumbered Res. Pap., 137 p. (NE)

1189. Nolley, J.W.
1972. Christmas tree marketing: a bibliography. Am. Christmas Tree J. 16(2): 23-28. (NE)

1190. Nyland, R.D. et al
1972. Volume related to weight in tree-length hardwoods. AFRI State Univ. NY, Res. Note 4, 1 p. (NE)

1191. Pendleton, T.H.
1973. Research needs in workmen's compensation. For. Prod. J. 23(3): 11-12. (NE)

1192. Stern, E.G.
1972. Southern pine pallets assembled with stiff-stock and hardened-steel pallet nails. Va. Poly. Tech. Inst. Bull. 106, 12 p. (NE)

1193. Stern, E.G.
1972. MIBANT test criteria for pallet nails. Va. Poly. Tech. Inst. Bull. 115, 26 p. (NE)

FOREST PRODUCTS AND ENGINEERING RESEARCH

FOREST ENGINEERING RESEARCH

Improving engineering systems

1194. Arola, R.A.
1973. Compression debarked chips from a whole-tree chipper. USDA For. Serv. Res. Note NC-147, 4 p. (NC)

1195. Arola, R.A., and J.R. Erickson
1973. Compression debarking of wood chips. USDA For. Serv. Res. Pap. NC-85, 11 p. (NC)

1196. Arola, R.A., and W.A. Hillstrom
1972. Compression debarking of branchwood chips from Finland. USDA For. Serv. Res. Note NC-143, 4 p. (NC)

1197. Biller, C.J.
1972. Are you overloading your skidder tires? USDA For. Serv. Res. Pap. NE-247, 7 p. (NE)

1198. Biller, C.J., and L.R. Johnson
1972. Comparing logging systems through simulation. ASAE Pap. No. 72-654, Winter Mtg., Chicago. (NE)

1199. Bradley, D.P., and F.E. Biltonen
1972. Economic operability—factors affecting harvest and transport costs. USDA For. Serv. Gen. Tech. Rep. NC-1 (Also Proc. Aspen Symp.), p. 16-20. (NC)

1200. Burke, J.D.
1972. Road and landing criteria for mobile-crane yarding systems. USDA For. Serv. Res. Note PNW-186, 13 p. (PNW)

1201. Carter, M.R., R.B. Gardner, and D.B. Brown
1973. Optimum economic layout of forest harvesting work roads. USDA For. Serv. Res. Pap. INT-133, 13 p. (INT)

1202. Erickson, J.R.
1972. The status of methods for debarking wood chips. J. TAPPI 55(8): 1216-1220. (NC)

1203. Erickson, J.R.
1972. The moisture content and specific gravity of the bark and wood of northern pulpwood species. USDA For. Serv. Res. Note NC-141, 3 p. (NC)

1204. Erickson, J.R.
1972. Bark removal after chipping challenges industry. North. Logger and Timber Processor 21(2): 10, 11, 30. (NC)

1205. Erickson, J.R.
1972. Field chipping: problems and solutions. FAO/NORAD Symp. on Production, Handling and Transport of Wood Chips, Hurdal, Norway. (NC)

1206. Gardner, R.B., and D.W. Hann
1972. Utilization of lodgepole pine logging residues in Wyoming increases fiber yield. USDA For. Serv. Res. Note INT-168, 6 p. (INT)

1207. Gardner, R.B., W.S. Hartsog, and G.L. Jacobsen
1973. Balloon logging. Agr. Engr. 54(2), p. 14-17. (INT)

1208. Hartsog, W.S., and M.J. Gonsior
1973. Analysis of construction and initial performance of the China Glen Road, Warnen District, Payette National Forest. USDA For. Serv. Gen. Tech. Rep. INT-5, 22 p. (INT)

1209. Johnson, L.R., D.L. Gochenour, Jr., and C.J. Biller
1972. Simulation analysis of timber-harvesting systems. Tech. Papers, 23rd Annual AIIE Conf. & Conv. Anaheim, CA, p. 353-361. (NE)

1210. Julien, L.M., J.C. Edgar, and T.M. Conder
1972. Segregation of aspen, balsam, and spruce wood and bark chips based on density differences. For. Prod. J. 22(6): 56-59. (NC)

1211. Lysons, H.H.
1973. Running skyline yarding system has merit. West. Conserv. J., January-February, p. 28-31. (PNW)

1212. Mann, C.N., and H.H. Lysons
1972. A method of estimating log weights. USDA For. Serv. Res. Pap. PNW-138, 75 p. (PNW)

1213. Peters, P.A., and J.D. Burke
1972. Average yarding distance on irregular-shaped timber harvest settings. USDA For. Serv. Res. Note PNW-178, 13 p. (PNW)

1214. Schillings, P.L.
1972. Engineering research in forest mechanization. Proc. La. State U. Annu. For. Symp., p. 13-17. (SO)

1215. Schillings, P.L.
1972. Equipment development: Highlights and hangups. Proc. Professional Affairs Program, Annu. Meet. SE Sect., Soc. Am. Foresters, January 26-28, 1972, Birmingham, Alabama, p. 60-63. (SO)

1216. Steinhilb, H.M., and J.R. Erickson
1972. Weights and centers of gravity for red pine, white spruce, and balsam fir. USDA For. Serv. Res. Pap. NC-75, 7 p. (NC)

1217. Sturos, J.A.
1973. Predicting segregation of wood and bark chips by differences in terminal velocity. USDA For. Serv. Res. Pap. NC-91. (NC)

1218. Sturos, J.A.
1973. Segregation of foliage from chipped tree tops and limbs. USDA For. Serv. Res. Note NC-146, 4p. (NC)

FOREST PRODUCTS UTILIZATION RESEARCH

Utilization potential and processing of wood

1219. Atherton, G.H., and J.R. Welty
1972. Drying rates of Douglas-fir veneer in superheated steam at temperatures to 800 degrees F. Wood Sci. 4(4): 209-218. (SO)

1220. Barger, R.L., and P.F. Ffolliott
1972. Physical characteristics and utilization of major woodland tree species in Arizona. USDA For. Serv. Res. Pap. RM-83, 80 p. (RM)

1221. Boone, R.S., and M. Chudnoff
1972. Compression wood formation and other characteristics of plantation-grown *Pinus caribaea*. USDA For. Serv. Res. Pap. ITF-13, 16 p. (ITF)

1222. Brisbin, R.L.
1972. A performance test of the log and tree grades for eastern white pine. USDA For. Serv. Res. Note NE-151, 5 p. (NE)

1223. Burton, J.D., and D.M. Smith
1972. Guying to prevent wind sway influences loblolly pine growth and wood properties. USDA For. Serv. Res. Pap. SO-80, 8p. (SO)

1224. Chudnoff, M., and E. Goytia
1972. Preservative treatments and service life of fence posts in Puerto Rico (1972 Progress Report). USDA For. Serv. Res. Pap. ITF-12, 28 p. (ITF)

1225. Clark, A., and D.R. Phillips
1972. Slash pine logs lose weight in storage. South. Lumberman 225(2795): 15-17. (SE)

1226. Conrad, L.A., and R.C. Koeppen
1972. An analysis of charcoal from the Brewster Site, Iowa. Plains Anthropologist 17(55): 52-54. (FPL)

1227. Cooper, G.A.
1972. Prefreezing reduces shrinkage and alters sorption in black walnut. For. Prod. J. 22(5): 54-60. (NC)

1228. Cooper, G.A., and E.F. Landt
1973. Horses and hardwoods. The West. Horseman 38(2): 73. (NC)

1229. Cooper, G.A., C.F. Bey, R.D. Lindmark, and R.C. Schlesinger
1972. Better hardwoods—from seeds to sawlogs and mills to markets. South. Lumberman 225(2800): 111-114. (NC)

1230. Davidson, H.L., and L.R. Gjovik
1972. Dual treatment of marine piles: Predrying and treatment. Amer. Wood- Preserv. Assoc. Proc. 68: 162-168. (FPL)

1231. Erickson, B.J., and D.C. Markstrom
1972. Predicting softwood cutting yields by computer. USDA For. Serv. Res. Pap. RM-98, 15 p. (RM)

1232. Fahey, T.D., and D.L. Hunt
1972. Lumber recovery from Douglas-fir thinnings at a bandmill and two chipping canters. USDA For. Serv. Res. Pap. PNW-131, 9 p. (PNW)

1233. Foulger, A.N., and J. Harris
1973. Volume of wood, bark, and needles after clearcutting a lodgepole pine stand. J. For. 71(2): 93-95. (FPL)

1234. Foulger, A.N., F. Freese, and J. Harris
1972. Effect of nutrient concentration on the stem anatomy of white ash seedlings. Wood and Fiber 4(2): 112-114. (FPL)

1235. Gillespie, R.H., and W.C. Lewis
1972. Evaluating adhesives for building construction. USDA For. Serv. Res. Pap. FPL-172, 12 p. (FPL)

1236. Gjovik, L.R., H.G. Roth, and H.L. Davidson
1972. Treatment of Alaskan species by double-diffusion and modified double-diffusion methods. USDA For. Serv. Res. Pap. FPL-182, 20 p. (FPL)

1237. Hallock, H., and E.H. Bulgrin
1972. Stabilization of hard maple flooring with polyethylene glycol 1000. USDA For. Serv. Res. Pap. FPL-187, 8 p. (FPL)

1238. Hanks, L.F.
1973. Green lumber grade yields for subfactory class hardwood logs. USDA For. Serv. Res. Pap. NE-256, 8 p. (NE)

1239. Hann, R.A., and J.C. Ward
1972. A look at and a result of reorganizing drying research at Forest Products Laboratory. Proc. West. Dry Kiln Clubs Ann. Meet., May 8-9, 1972, Redding, Calif., 6 p. (FPL)

1240. Heebink, B.G.
1972. Some views on large structural particleboard panels. USDA For. Serv. Res. Note FPL-0220, 8 p. (FPL)

1241. Heebink, B.G.
1972. Irreversible dimensional changes in panel materials. For. Prod. J. 22(5): 44-48. (FPL)

1242. Heebink, B.G., W.F. Lehmann, and F.V. Hefty
1972. Reducing particleboard pressing time: Exploratory Study. USDA For. Serv. Res. Pap. FPL-180, 12 p. (FPL)

1243. Henley, J.W.
1972. Grading sugar pine saw logs in trees. USDA For. Serv. Res. Pap. PNW-132, 8 p. (PNW)

1244. Hiller, C.H., F. Freese, and D.M. Smith
1972. Relationships in black walnut heartwood between color and other physical and anatomical characteristics. Wood and Fiber 4(1): 38-42. (FPL)

1245. Hse, C.Y.
1972. Influence of resin formulation variables on bond quality of southern pine plywood. For. Prod. J. 22(9): 104-108. (SO)

1246. Hse, C.Y.
1972. Method for computing a roughness factor for veneer surfaces. Wood Sci. 4(4): 230-233. (SO)

1247. Hse, C.Y.
1972. Surface tension of phenol formaldehyde wood adhesives. Holzforsch. 26(2): 82-85. (SO)

1248. Hse, C.Y.
1972. Wettability of southern pine veneer by phenol formaldehyde wood adhesives. For. Prod. J. 22(1): 51-56. (SO)

1249. Johnson, E.J., and P. Koch
1972. Thermal reactions of small loblolly pine cubes heated on one face in an air atmosphere. Wood Sci. 4(3): 154-162. (SO)

1250. Jokerst, R.W.
1972. Feasibility of producing a high-yield laminated structural product: Residual heat of drying accelerates adhesive cure. USDA For. Serv. Res. Pap. FPL-179, 10 p. (FPL)

1251. Jokerst, R.W.
1972. Evaluation of adhesive-bond quality in telephone crossarms after 16 to 23 years of exterior exposure. USDA For. Serv. Res. Pap. FPL-171, 8 p. (FPL)

1252. Jung, P., W. Tanner, and K. Wolter
1972. The fate of *myo*—inositol in *Fraxinus* tissue cultures. Phytochem. 11(5): 1655-1659. (FPL)

1253. Kinney, R.E., E.L. Burmeister, and I.B. Sachs
1972. A simple device for fiber stretching in the scanning electron microscope. Wood and Fiber 4(1): 53-55. (FPL)

1254. Koch, P.
1972. Utilization of the southern pines. USDA, Agric. Hanb. 420, 2 Vols., 1663 p. (SO)

1255. Koch, P.
1972. The three rings-per-inch dense southern pine—should it be developed? Proc. Symp. on the Effect of Growth Acceleration on Properties of Wood, 1971 (Also in J. Forest. 70(6): 332, 334, 336). (SO)

1256. Koch, P.
1972. Changing raw material supplies and their effect upon wood processing technology. Preprint, 7th World Forest. Congr., Buenos Aires., Argentina 9 p. (SO)

1257. Koch, P.
1972. Process for steam straightening and kiln drying lumber. U.S. Pat. 3,680,219, U.S. Pat. Off., Wash., DC. (SO)

1258. Koch, P.
1972. Drying southern pine at 240 degrees F — Effects of air velocity and humidity, board thickness and density. For. Prod. J. 22(9): 62-67. (SO)

1259. Lane, P.H., R.O. Woodfin, J.W. Henley, and M.E. Plank
1972. Lumber yield from Sitka spruce in southeastern Alaska. USDA For. Serv. Res. Pap. PNW-134, 44 p. (PNW)

1260. Lane, P.H., R.O. Woodfin, J.W. Henley, and M.E. Plank
1973. New timber cruising grades for coast Douglas-fir. USDA For. Serv. Res. Pap. PNW-151, 12 p. (PNW)

1261. Lane, P.H., R.O. Woodfin, J.W. Henley, and M.E. Plank
1973. Timber cruising grades for coast Douglas-fir. USDA For. Serv. Misc. Publ., 4 p. (PNW)

1262. Lehmann, W.F.
1972. Moisture-stability relationships in wood-base composition boards. For. Prod. J. 22(7): 53-59. (FPL)

1263. Lorenz, L.F., and L.R. Gjovik
1972. Analyzing creosote by gas chromatography: Relationship to creosote specification. Proc. Am. Wood- Preserv. Assoc. 68: 32-34. (FPL)

1264. Lowery, D.P.
1972. Predicting moisture content in Douglas-fir and ponderosa pine boards. For. Prod. J. 22(12): 35-39. (INT)

1265. Lowery, D.P.
1972. Predicting moisture content in ponderosa pine boards. Proc. West. Dry Kiln Clubs 23rd Annu. Meet. (INT)

1266. Lowery, D.P.
1972. Vapor pressures generated in wood during drying. Wood Sci. 5(1): 73-80. (INT)

1267. Lutz, J.F.
1972. Veneer species that grow in the United States. USDA For. Serv. Res. Pap. FPL-167, 127 p. (FPL)

1268. McMillen, J.M., and R.C. Baltes
1972. New kiln schedule for presurfaced oak lumber. For. Prod. J. 22(5): 19-26. (FPL)

1269. McMillin, C.W.
1972. Laser machining—A status report. South. Lumberman 225(2795): 19-20. (SO)

1270. McMillin, C.W., and G.E. Woodson
1972. Moisture content of southern pine as related to thrust, torque, and chip formation in boring. For. Prod. J. 22(11): 55-59. (SO)

1271. Maeglin, R.R., and H.E. Wahlgren
1972. Western wood density survey—Report No. 2. USDA For. Serv. Res. Pap. FPL-183, 24 p. (FPL)

1272. Malcolm, F.B., and H. Hallock
1972. Effects of three sawing methods on warp of hard maple dimension cuttings. For. Prod. J. 22(4): 57-60. (FPL)

1273. Manwiller, F.G.
1972. Tracheid dimensions in rootwood of southern pine. Wood Sci. 5(2): 122-124. (SO)

1274. Manwiller, F.G.
1972. Volumes, wood properties, and fiber dimensions of fast- and slow-grown spruce pine. Proc. Symp. on the Effect of Growth Acceleration on Properties of Wood, 1971. (SO)

1275. Manwiller, F.G.
1972. Wood and bark properties of spruce pine. USDA For. Serv. Res. Pap. SO-78, 25 p. (SO)

1276. Markstrom, D.C., and R.A. Hann
1972. Seasonal variation in wood permeability and stem moisture content of three Rocky Mountain species. USDA For. Serv. Res. Note RM-212. (RM)

1277. Markstrom, D.C., and V.P. Yerkes
1972. Specific gravity variation with height in Black Hills ponderosa pine. USDA For. Serv. Res. Note RM-213, 4 p. (RM)

1278. Miniutti, V.P.
1973. Contraction in softwood surfaces during ultraviolet irradiaiton and weathering. J. of Paint Tech. 45(577): 27-34. (FPL)

1279. Mueller, L.A., R.L. Barger, A. Bourke, and D.C. Markstrom
1972. Roll laminating fiber overlays on low-grade ponderosa pine lumber. USDA For. Serv. Res. Pap. RM-97, 28 p. (RM)

1280. Murmanis, L.L., and I.B. Sachs
1972. Filamentous component (slime) in coniferous sieve elements. Proc. Electron Microscopy Soc. Am. XIII: 214-215. (FPL)

1281. Nelson, N.D.
1973. Effects of amounts of parenchyma on quantity of phenolic extractives produced during formation of heartwood in *Juglans nigra* and *Quercus rubra*. Ph.D. Dissertation (Forestry and Forest Prod.), Univ. Wis., Madison. (FPL)

1282. Okkonen, E.A., H.E. Wahlgren, and R.R. Maeglin
1972. Relationships of specific gravity to tree height in commercially important species. For. Prod. J. 22(7): 37-42. (FPL)

1283. Peters, C.C., A.F. Mergen, and H.R. Panzer
1972. Thick-slicing of wood: Effects of wood and knife inclination angle. For. Prod. J. 22(9): 84-91. (FPL)

1284. Phillips, D.R., and J.G. Schroeder
1972. Some physical characteristics of bark from plantation-grown slash and loblolly pine. For. Prod. J. 22(10): 30-33. (SE)

1285. Quirk, J.T., D.M. Smith, F. Freese, and K.E. Wolter
1972. Stimulatory Effect of torque on the radial growth of red pine trees. Plant Physiol. Vol. 49 Supplement, p. 5 (Abstract No. 28), Annual Mtg, August 27-31, 1972, Univ. of Minn., Minneapolis, Minn. (FPL)

1286. Rast, E.D., D.L. Sonderman, and G.L. Gammon
1973. A guide to hardwood log grading (revised). USDA For. Serv. Gen. Tech. Rep. NE-1, 32 p. (NE)

1287. Rietz, R.C.
1972. News and views of this kiln-drying business: A calendar for air-drying lumber in the Upper Midwest. USDA For. Serv. Res. Note FPL-0224, 3 p. (FPL)

1288. Rosen, H.N.
1973. Continuous measurement of the swelling of wood. For. Prod. J. 23(3): 55-57. (NC)

1289. Sachs, I.B., and R.E. Kinney
1972. An "ultrafilter" in the margo of mature bordered pits? Proc. Electron Micros. Soc. of Am. XIII: 206-207. (FPL)

1290. Saucier, J.R., A. Clark, and R.G. McAlpine
1972. Above ground biomass yields of short-rotation sycamore. Wood Sci. 5(1): 1-6. (SE)

1291. Schaffer, E.L., J.L. Tschernitz, C.C. Peters, R.C. Moody, R.W. Jokerst, and J.J. Zahn
1972. FPL Press-Lam process: Fast efficient conversion of logs into structural products. For. Prod. J. 22(11): 11-18. (FPL)

1292. Schaffer, E.L., J.L. Tschernitz, C.C. Peters, R.C. Moody, R.W. Jokerst, and J.J. Zahn
1972. Feasibility of producing a high-yield laminated structural product General summary. USDA For. Serv. Res. Pap. FPL-175, 20 p. (FPL)

1293. Schumann, D.R.
1973. Dimension stock yields from lumber of three hardwood species. For. Prod. J. 23(3): 17-21. (FPL)

1294. Schumann, D.R.
1972. Predicting dimension parts cost. Woodr. and Furn. Digest, p. 26-28. (FPL)

1295. Simpson, W.T., and R.C. Baltes
1972. Accelerating oak air drying by presurfacing. USDA For. Serv. Res. Note FPL-0223, 12 p. (FPL)

1296. Skolmen, R.G.
1972. Paintability of four woods in Hawaii. USDA For. Serv. Res. Note PSW-267, 4 p. (PSW)

1297. Skolmen, R.G.
1972. Specific gravity variation in *Robusta eucalyptus* grown in Hawaii. USDA For. Serv. Res. Pap. PSW-78, 7 p. (PSW)

1298. Stewart, H.A.
1972. Abrasive or knife planing? Furn. Prod. 35(241): 20-23. (NC)

1299. Stewart, H.A.
1972. Abrasive vs. knife planing. Wood and Wood Prod. 77(8): 73-76. (NC)

1300. Tschernitz, J.L.
1973. Enzyme mixture improves creosote treatment of kiln-dried Rocky Mountain Douglas-fir. For. Prod. J. 23(3): 30-38. (FPL)

1301. Wahlgren, H.E.
1972. Research on solid waste recycling at the Forest Products Laboratory. Ames For., Vol. 59, Iowa State U., Ames, Iowa. (FPL)

1302. Wahlgren, H.E., and D.R. Schumann
1973. Properties of major southern pines: Part I, Wood density surveys. USDA For. Serv. Res. Pap. FPL-176-177, Part I 58p. - Part II 10p. (FPL)

1303. Ward, J.C.
1972. Anaerobic bacteria associated with honeycomb and ring failure in red and black oak lumber. Phyta 62(7): 796, (Abstr.). (FPL)

1304. Ward, J.C., R.A. Hann, R.C. Baltes, and E.H. Bulgrin
1972. Honeycomb and ring failure in bacterially infected red oak lumber after kiln drying. USDA For. Serv. Res. Pap. FPL-165, 36 p. (FPL)

1305. Wengert, E.M.
1972. Review of high-temperature kiln- drying of hardwoods. South. Lumberman, Sept. 15, p. 17-19. (FPL)

1306. Wilcox, W.W.
1972. Apparent defects and hidden defects. Proc. 20th Western Int. For. Dis. Work Conf. (PNW)

1307. Woodfin, R.O., and W.Y. Pong
1972. Dry veneer volume losses in production of red and white fir plywood. USDA For. Serv. Res. Note PNW-191, 12 p. (PNW)

1308. Woodson, G.E., and C.W. McMillin
1972. Boring deep holes in southern pine. For. Prod. J. 22(4): 49-53. (SO)

1309. Wu, K.Y.T., and E.A. McGinnes
1972. Variability of intra-increment chemical properties in some hardwood species. 163rd Am. Chem. Soc. Mgt., Boston, Abstracts. (NE)

1310. Zicherman, J.B., and R.J. Thomas
1971. Distribution of loblolly pine ash residue. TAPPI 54(10): 1727-1730. (SO)

1311. Zicherman, J.B., and R.J. Thomas
1972. Analysis of loblolly pine ash materials. Holzforsch. 26(4): 150-152. (SO)

1312. Zicherman, J.B., and R.J. Thomas
1972. Scanning electron microscopy of weathered coatings on wood. J. Paint Technol. 44(570): 88-94. (SO)

Wood chemistry and fiber products

1313. Auchter, R.J.
1972. Pulping without barking increases fiber yield. Pulb and Paper 46(6): Addendum p. 6-7. (FPL)

1314. Auchter, R.J.
1973. Recycling forest products retrieved from urban waste. For. Prod. J. 23(2): 12-16. (FPL)

1315. Baker, A.J.
1973. Effect of lignin on the *in vitro* digestibility of wood pulp. J. Animal Sci. 36(4): 768-771. (FPL)

1316. Byrd, V. L.
1972. Effect of relative humidity changes on compressive creep response of paper. TAPPI 55(11). (FPL)

1317. Caulfield, D.F.
1973. Small-angle X-ray scattering by paper: A new method for investigating inter fiber bonding. TAPPI 56(3): 102-106. (FPL)

1318. Deshpande, V.H., A.V. Rama Rao, R. Srinivasan, and K. Venkataraman
1972. Wood phenolics of *Morus* spp: Part II - The action of DDQ on mulberrin, mulberrochromene, and cyclomulberrin. Indian J. Chem. 10:681-682. (FPL)

1319. Dhruva, B.R., A.V. Rama Rao, R. Srinivasan, and K. Venkataraman
1972. Structure of a quinone from teak tissue culture. Indian J. Chem. 10: 683-685. (FPL)

1320. Fahey, D.J., and D.S. Pierce
1973. The role of phenolic resin in imparting properties to dry-formed hardboards. TAPPI 56(3): 53-56. (FPL)

1321. Feist, W.C., E.L. Springer, and G.J. Hajny
1972. Determining loss of wood substance in outside chip storage: A comparison of two methods. USDA For. Serv. Res. Pap. FPL-189, 9 p. (FPL)

1322. Godshall, W.D., and J.W. Koning
1972. Method for measuring and controlling web tension of corrugating medium during singlefacing process. USDA For. Serv. Res. Note FPL-0219, 7 p. (FPL)

1323. Harkin, J.M.
1972. Wood and pulp chromophores: A critical review. TAPPI 55(7): 991-996. (FPL)

1324. Harris, J.F.
1972. Alkaline decomposition of D-xylose-1-^{14}C, D-glucose-1-^{14}C, and D-glucose-6-^{14}C. Carbohyd. Res. 23:207-215. (FPL)

1325. Horn, R.A.
1972. How fiber morphology affects pulp characteristics and properties of paper. Chem 26 8(5): 39-44. (FPL)

1326. Horn, R.A., and R.J. Auchter
1972. Kraft pulping of pulpwood chips containing bark. Pap. Trade J. 156(45): 55-59. (FPL)

1327. Kirk, T.K., and W.E. Moore
1972. Removing lignin from wood with white-rot fungi and digestibility of resulting wood. Wood and Fiber 4(2): 72-79. (FPL)

1328. Koning, J.W., E.W. Kuenzi, R.C. Moody, and W.D. Godshall
1972. Improving comparability of paperboard test results: Using flexible and rigid type testing machines. TAPPI 55(5): 757-760. (FPL)

1329. Laundrie, J.F., and J.G. Berbee
1972. High yields of kraft pulp from rapid-growth hybrid poplar trees. USDA For. Serv. Res. Pap. FPL-186, 22 p. (FPL)

1330. Laundrie, J.F., and J.G. Berbee
1973. The disappearing forest—Boosting pulp yield per acre with short- rotation corps. Chem 26 Paper Processing 9(2): 48-50. (FPL)

1331. Laundrie, J.F., and D.J. Fahey
1973. Kraft pulps and hardboards from short-rotation sycamores. Pap. Trade J. 157(6): 26-27. (FPL)

1332. Laundrie, J.F., and J.H. Klungness
1973. Dry methods of separating plastic films from waste paper. Pap. Trade J. 157(6): 34-36. (FPL)

1333. Laundrie, J.F., and J.H. Klungness
1973. Effective dry methods of separating thermoplastic films from waste papers. USDA Forest Serv. Res. Pap. FPL-200, 11 p. (FPL)

1334. Mohaupt, A.A., and J.W. Koning
1972. A practical method for recycling wax-treated corrugated. Boxboard Containers 79(6) 60-62. (FPL)

1335. Moore, W.E., M.J. Effland, and M.A. Millett
1972. Hydrolysis of wood and cellulose with cellulytic enzymes. J. Agr. Food Chem. 20(6): 1173-1175. (FPL)

1336. Mujumdar, R.B., R. Srinivasan, and K. Venkataraman
1972. Taxiresinol, a new lignan in the heartwood of *Taxus baccata*. Indian J. Chem. 10: 677-680. (FPL)

1337. Myers, G.C.
1972. FPL studies recovery of wood fiber from household refuse. Pulp and Paper 46(11): 128-130. (FPL)

1338. Myers, G.C.
1972. Separating household wastepaper: Survey results. TAPPI 55(3): 389-392. (FPL)

1339. Rama Rao, A.V., S.S. Rathi, and K. Venkataraman
1972. Chaplashin, a flavone containing an oxepine ring from the heartwood of *Artocarpus chaplasha* Roxb. Indian J. Chem. 10: 905-907. (FPL)

1340. Rama Rao, A.V., S.S. Rathi, and K. Venkataraman
1972. Electron-impact spectra of flavones with nuclear isoprenoid substituents. Indian J. Chem. 10: 989-1001. (FPL)

1341. Ravindranath, B., and T.R. Seshadri
1972. Chemistry of the santalin pigments: I. Structure of santalin permethyl ether. Tetrahedron Let. No. 13: 1201-1204. (FPL)

1342. Rowe, J.W., M.K. Seikel, D.N. Roy, and E. Jorgensen
1972. Chemotaxonomy of *Ulmus*. Phytochem. 11: 2513-2517. (FPL)

1343. Rowell, R.M.
1972. Acyl migrations in the synthesis of ethyl 4-0-methyl-beta-D-glucopyranoside. Carbohydrate Res. 23: 417-424. (FPL)

1344. Rowell, R.M., and J. Green
1972. Effects of trans alpha-hydroxyl groups in alkaline degradation of glycosidic bonds. USDA For. Serv. Res. Pap. FPL-188, 8 p. (FPL)

1345. Rowell, R.M., and J. Green
1972. Kinetics of oxidative alkaline degradation of end group stabilized cellulose models. TAPPI 55(9): 1326-1327. (FPL)

1346. Scott, R.W., and J. Green
1972. A rapid analysis for total carbohydrate in wood or pulp: Dehydrating to furans in concentrated sulfuric acid. TAPPI 55(7): 1061-1063. (FPL)

1347. Seshadri, T.R.
1972. Polyphenols of *Pterocarpus* and *Dalbergia* woods (Review article). Phytochem. 11: 881-898. (FPL)

1348. Smith, W.E., and V.L. Byrd
1972. Fiber bonding and tensile stress-strain properties of earlywood and latewood handsheets. USDA For. Serv. Res. Pap. FPL-193, 9 p. (FPL)

1349. Stockmann, V.E.
1972. Developing a hypothesis: Native cellulose elementary fibrils are formed with metastable structure. Biopolymers 11: 251-270. (FPL)

1350. Venkataraman, K.
1972. Wood phenolics in the chemotaxonomy of the *Moraceae* (Review article). Phytochemistry 11: 1571-1586. (FPL)

1351. Wangaard, F.F., and G.E. Woodson
1973. Fiber length-fiber strength interrelationship for slash pine and its effect on pulp-sheet properties. Wood Sci. 5(3): 235-240. (SO)

1352. Zinkel, D.F., and B.B. Evans
1972. Terpenoids of *Pinus strobus* cortex tissue. Phytochem. 11: 3387-3389. (FPL)

Wood engineering

1353. Anderson, L.O.
1972. Condensation problems: Their prevention and solution. USDA For. Serv. Res. Pap. FPL-132, 37 p. (FPL)

1354. Anderson, L.O., T.B. Heebink, and A.E. Oviatt
1972. Construction guides for exposed wood decks. USDA, Agric. Handb. 432, 78 p. (FPL)

1355. Bendtsen, B.A.
1972. Important structural properties of four wester softwoods: white-pine, sugar pine, western redcedar, and Port-Orford cedar. USDA For. Serv. Res. Pap. FPL-191, 17p. (FPL)

1356. Bendtsen, B.A.
1973. Mechanical properties and specific gravity of randomly sampled subalpine fir. USDA For. Serv. Res. Pap. FPL-197, 12 p. (FPL)

1357. Bendtsen, B.A., and R.L. Ethington
1972. Properties of major southern pines: Part I, Wood density survey; Part II, Structural properties and specific gravity. USDA For. Serv. Res. Pap. FPL-176-177, Part I 58p. - Part II 10p. (FPL)

1358. Bohannan, B.
1971. Interaction of structural elements. Proc. Meet. Int. Union of Forestry Res. Organ.; Working Group on Struc. Util., p. 141-147. (FPL)

1359. Bohannan, B.
1972. FPL timber bridge deck research. J. Struct. Div. Am. Soc. Civil Eng. 98(ST3): 729-740. (FPL)

1360. Bohannan, B.
1972. Glued-laminated timber bridges—reality or fantasy. Proc. 1972 Annu. Meet. of Amer. Inst. of Timber Construction. (FPL)

1361. Bohannan, B.
1972. Glu-lam development and current FPL research. South. Lumberman 225(2800): 170-171, 174. (FPL)

1362. Bohannan, B.
1972. Strength criteria of glued-laminated timber. Natl. Bur. of Standards Special Pub.. 361, Vol. 1. Performance concept in Buildings, Proc. of Joint RILEM-ASTM-CIB Symp., May 2-5, 1972 Philadelphia, Pa. p. 625-632. (FPL)

1363. Boller, K.H.
1972. A facility to evaluate three-dimensional performance of house modules. USDA For. Serv. Res. Note FPL-0225, 12 p. (FPL)

1364. Boller, K.H.
1972. Effect of exposure on a proposed glass-reinforced plastic laminate to be used in building constructions. Natl. Bur. of Standards Rep. PB 211-314 available from NTIS. (FPL)

1365. Chern, J., and E.W. Kuenzi
1972. Development of basic information for the design of paper shipping sacks. TAPPI 55(10): 1477-1481. (FPL)

1366. Duff, J.E.
1972. Vapor barrier decreases moisture conditions in wood walls exposed to air conditioning and heating. USDA For. Res. Pap. SE-98, 8 p. (SE)

1367. Ethington, R.L., and H.O. Fleischer
1973. A perspective on structural lumber grading. The Building Official and Code Administrator VII(3): 10-11. (FPL)

1368. U.S.D.A. Forest Service, Forest Products Laboratory
1972. FPL Homes—Quality on a budget. Unnumbered Rep., 36 p. (FPL)

1369. Gaby, L.I.
1972. Abrasive planing up-grades pine surfaces. South. Lumberman 225(2800): 131-133. (SE)

1370. Gaby, L.I.
1972. Warping in southern pine studs. USDA For. Serv. Res. Pap. SE-96, 8 p. (SE)

1371. Gerhards, C.C.
1972. Relationship of tensile strength of southern pine dimension lumber to inherent characteristics. USDA For. Serv. Res. Pap. FPL-174, 32 p. (FPL)

1372. Gillespie, R.H., and W.C. Lewis
1972. Evaluating adhesives for building construction. USDA For. Serv. Res. Pap. FPL-172, 13 p. (FPL)

1373. Godshall, W.D., and J.W. Koning
1972. Method for measuring and controlling web tension of corrugating medium during single facing process. USDA For. Serv. Res. Note FPL-0219, 7 p. (FPL)

1374. Grantham, J.B., and T.B. Heebink
1972. Insuring noise control in wood-framed buildings. For. Prod. J. 22(5): 36-43. (PNW)

1375. Grantham, J.B., and T.B. Heebink
1972. Sound insulation in a modular motel. USDA For. Serv. Res. Note PNW-193, 8 p. (PNW)

1376. Jones, R.E.
1972. Airborne noise control for vertically stacked bathrooms with mechanical exhaust systems. USDA For. Serv. Res. Note FPL-0222, 10 p. (FPL)

1377. Koning, J.W., E.W. Kuenzi, R.C. Moody, and W.D. Godshall
1972. Improving comparability of paperboard test results: Using flexible and rigid type testing machines. TAPPI 55(5): 757-760. (FPL)

1378. McNatt, J.D.
1973. Buckling due to linear expansion of hardboard. For. Prod. J. 23(1): 37-43. (FPL)

1379. Millett, M.A., and C.C. Gerhards
1972. Accelerated aging: Residual weight and flexural properties of wood heated in air at 115 degrees to 175 degrees C. Wood Sci. 4(4): 193-201. (FPL)

1380. Montrey, H.M., and E.W. Kuenzi
1972. Design parameters for torsion of sandwich strips having trapezoidal, rectangular, and triangular cross sections. USDA For. Serv. Res. Pap. FPL-156, 40 p. (FPL)

1381. Moody, R.C.
1972. Tensile strength of lumber laminated from 1/8-inch-thick veneers. USDA For. Serv. Res. Pap. FPL-181, 28 p. (FPL)

1382. Moody, R.C., and C.C. Peters
1972. Feasibility of producing a high-yield laminated structural product: Strength properties of rotary knife-cut, laminated southern pine. USDA For. Serv. Res. Pap. FPL-178, 12 p. (FPL)

1383. Oviatt, A.E.
1973. Wood is ideal in pool natatorium roof construction. Data and Reference Annual Edition of Swimming Pool Weekly and Swimming Pool Age, p. 178, 180, 182, 183, 186-188, 190. (PNW)

1384. Ramaker, T.J., and M.D. Davister
1972. Predicting performance of hardboard in I-beams. USDA For. Serv. Res. Pap. FPL-185, 12 p. (FPL)

1385. Schaffer, E.L.
1972. Modeling the creep of wood in a changing moisture environment. Wood and Fiber 3(4): 232-235. (FPL)

1386. Sherwood, G.E.
1972. Nu-frame roof joints. USDA For. Serv. Res. Pap. FPL-88, 9 p. (Supl.). (FPL)

1387. Sherwood, G.E.
1972. FPL designs meet family housing needs. USDA For. Serv. Res. Pap. FPL-173, 28 p. (FPL)

1388. Sherwood, G.E.
1972. Wood structures can resist hurricanes. Civil Eng. 42(9): 91-94. (FPL)

1389. Stern, R.K., and D.E. Dunmire
1972. Appalachian hardwoods for pallets: Correlation between service and laboratory testing. USDA For. Serv. Res. Pap. FPL-169, 16 p. (FPL)

1390. Tuomi, R.L.
1972. Advancements in timber bridges through research and engineering. Proc. of 13th Annu. Colo. State Univ. Bridge Eng. Conf., p. 34-61. (FPL)

1391. Vick, C.B.
1972. Short fasteners for control of nail popping under resilient tile flooring. USDA For. Serv. Res. Pap. SE-99, 15 p. (SE)

1392. Vick, C.B., and R.M. Walker
1973. Underlayment fastening device. (U.S. Pat. No. 3,717,067) U.S. Pat. Off., Washington, D.C. (SE)

1393. Vick, C.B., and R.M. Walker
1972. A new nail to eliminate nail popping and dimpling under resilient tile flooring. USDA For. Serv. Res. Pap. SE-102, 14 p. (SE)

1394. Wilkinson, T.L.
1972. Analysis of nailed joints with dissimilar members. J. Struct. Div., ASCE (98) ST9 Proc. Pap. 9189, p. 2005-2013. (FPL)

1395. Wilkinson, T.L.
1972. Effect of deformed shanks, prebored lead holes, and grain orientation on the elastic bearing constant for laterally loaded nail joints. USDA For. Serv. Res. Pap. FPL-192, 13 p. (FPL)

Miscellaneous

1396. Claxton, H.D., and G. Riensi
1972. An analytical procedure to assist decision-making in a government research organization. USDA For. Serv. Res. Pap. PSW-80, 20 p. (PSW)

1397. George, R.P., R.M. Albrecht, K.B. Raper, I.B. Sachs, and A.P. MacKenzie
1972. Scanning electron microscopy of spore germination in *Dictyostellium discodeum*. J. Bacter. 112(3): 1383-1386. (FPL)

1398. Henderson, J.H., R.N. Clayton, M.L. Jackson, J.K. Syers, R.W. Rex, J.L. Brown, and I.B. Sachs
1972. Cristobalite and quartz isolation from soils and sediments by hydrofluosilicic acid treatment and heavy liquid separation. Proc. Soil Sci. Soc. Am. 36(5): 830-835. (FPL)

1399. Landis, M.W.
1972. Wood. Paper presented at the Products and Materials (Safety and Potential Liability) Eng. Short Course, Univ. Wis. (FPL)

1400. Landt, E.F., and R.E. Phares
1973. Black walnut: An American wood. USDA For. Serv., Am. Woods FS-270, 7 p. (NC)

1401. Pomeranz, Y., and I.B. Sachs
1972. Determining the structure of the barley kernel by scanning electron microscopy. Cereal Chem. 49(1): 1-7. (FPL)

1402. Pomeranz, Y., and I.B. Sachs
1972. Scanning electron microscopy of the oak kernel. Cereal Chem. 49(1): 20-25. (FPL)

1403. Pomeranz, Y., and I.B. Sachs
1972. Scanning electron microscopy of the buckwheat kernel. Cereal Chem. 49(1): 23-29. (FPL)

1404. Riensi, G., and H.D. Claxton
1972. A data collection and processing procedure for evaluating a research program. USDA For. Serv. Res. Pap. PSW-81, 18 p. (PSW)

1405. Saucier, J.R.
1973. American chestnut— An American wood. USDA For. Serv., Am. Woods FS-230, 6 p. (SE)

1406. Schroeder, J.G.
1972. Butternut—an American wood. USDA For. Serv. Am. Woods FS-223, 6 p. (SE)

1407. Vick, C.B.
1973. Yellow-poplar—an American wood. USDA For. Serv., Am. Woods FS-272, 7 p. (SE)

AUTHORS

Numbers refer to the Publications List which begins on p. 47.

A

Aarhus, D.G. 356
Abrahamson, L.P. 314, 315
Adams, R.S. 821
Ahmad, A. 708
Aho, P.E. 541
Alban, D.H. 909, 910
Albrecht, R.M. 1397
Alcaniz, R. 195
Aldon, E.F. 49, 50
Aldrich, R.C. 1078, 1079, 1096, 423, 424, 431
Alexander, L.T. 385
Alexander, R.R. 734, 735, 903, 922
Alkidas, A. 665
Allan, G.G. 714
Alley, J.R. 717
Almas, D.P. 725
Alston, R.M. 1123
Altman, J. 555
Amburgey, T.L. 614
Amidon, E.L. 718
Amidon, P.H. 291
Amman, C.D. 316
Amman, G.D. 317, 422
Anand, V.K. 859, 880
Anderson, A.E. 148, 149, 150, 151, 213, 214
Anderson, G.W. 473, 474
Anderson, H.E. 622
Anderson, H.W. 1128, 126, 133, 15, 51
Anderson, L.O. 1353, 1354
Anderson, N.G. 395
Anderson, R.B. 1177
Anderson, W.C. 1103, 1104, 1105, 1139
Andregg, M.J. 245
Andrews, T.L. 318
Anonymous 52, 53, 54, 55
Applefield, M. 1140, 1141, 1146, 1178, 1179, 1180, 1181
Arnold, J.F. 1
Arola, R.A. 1194, 1195, 1196
Asher, W.C. 319
Ashley, B.S. 773
Atherton, G.H. 1219
Aubertin, G.M. 140
Auchmoody, L.R. 758, 759
Auchter, R.J. 1313, 1314, 1326
Austin, J.W. 1142
Avila Cortes, G. 799

B

Baker, A.J. 1315
Baker, B.H. 422
Baker, J.B. 760
Baker, M.F. 152, 162
Bakshi, B.K. 475, 476, 477, 781
Ballard, T.M. 16
Balls, L.D. 191
Baltes, R.C. 1268, 1295, 1304
Barger, J.H. 393
Barger, R.L. 1220, 1279
Barnard, J.E. 1039, 1040, 1106, 927
Barnes, A.H. 76
Barnett, J.P. 782, 783, 784
Barney, R.J. 649, 650
Barnhisel, R.I. 143
Barras, S.J. 320, 321
Barrett, J.W. 1118, 248, 838, 906
Batch, D.L. 141
Batzer, H.O. 300, 322
Baughman, R.G. 687
Baumgras, J.E. 1182
Baur, P.S. 478, 488, 561
Bauske, B.W. 677
Baxter, C.C. 436
Beagle, L.D. 731
Beal, R.H. 443, 444, 445, 446, 447
Beaton, J.A. 460, 479, 480
Beck, D.E. 889
Becker, W.A. 952
Beckwith, R.C. 323, 324, 680, 681
Bega, R.V. 1085, 449, 461, 509
Behar, J.F. 453
Behar, J.V. 619
Bell, J.F. 890
Bell, S. 755
Bellany, T.R. 1058
Beltz, R.C. 1059
Ben-Sasson, R. 863
Benator, C.R. 885
Bender, F.W. 452, 484
Bendtsen, B.A. 1355, 1356, 1357
Benedict, G.W. 635
Bennett, F.A. 839, 840, 911, 912
Bennett, W.H. 325, 394
Bensend, D.W. 1027
Benzie, J.W. 736, 737, 738, 739, 785
Berbee, J.G. 1329, 1330
Berg, A.B. 890
Berg, A.R. 623
Berger, J.M. 1083
Bergsvik, K. 1119
Berndt, H.W. 122
Berntsen, C.M. 715
Beroza, M. 376
Berry, F.H. 460, 479, 480, 481
Berry, L.R. 247
Bertelson, D.F. 1060, 1061, 1062, 1063
Bertolin, G.E. 82
Bethlahmy, N. 56
Bethune, J.E. 933
Bey, C.F. 1229, 741, 841, 953, 954, 991
Biesterfeldt, R.C. 1094
Biller, C.J. 1197, 1198, 1209
Biltonen, F.E. 1150, 1199
Bingham, R.T. 1028, 482, 570, 571, 981, 982
Birch, M.C. 351
Bjorkbom, J.C. 761, 762, 786, 787
Blackmarr, W.H. 663
Blackmon, B.G. 763
Blair, R.M. 207
Blaisdell, J.P. 202
Blakely, A.D. 655
Blank, R.W. 464
Blum, B.M. 934
Blyth, J.E. 1064, 1065
Bodman, R. 1186
Boeker, E.L. 171, 215
Boelter, D.H. 27, 57
Bohannan, B. 1358, 1359, 1360, 1361, 1362
Bohart, G.E. 326
Bohning, J.W. 174
Boldt, C.E. 891
Bollen, W.B. 17, 22, 26, 512
Boller, K.H. 1363, 1364
Bond, R.S. 1147
Bones, J.T. 1066, 1067, 1068
Bonner, F.T. 719, 720
Books, D.J. 153, 216, 259
Boone, R.S. 1221
Bormann, F.H. 24
Boster, R.S. 1133
Bourke, A. 1279
Bourne, C.E. 542
Bowden, D.C. 148, 149, 150, 213, 214
Bowers, T.M. 1039
Bowman, K.O. 695, 696
Boyd, R.J. 788, 836
Boyer, W.D. 983
Brackebusch, A.P. 624
Bradley, D.P. 1148, 1149, 1150, 1199
Brady, H.A. 740, 842
Bramlett, D.L. 789, 790, 791, 955, 992
Branan, J.R. 792
Brander, R.B. 216, 246
Branson, B.A. 141
Brantley, J.N. 395
Breillatt, J.P. 395
Brendemuehl, R.H. 760
Brender, E.V. 692, 721, 806
Brewer, J.W. 327
Brewster, S.F., Jr. 166
Brink, G.E. 33, 88, 89
Brisbin, R.L. 1222
Briscoe, C.B. 797, 800
Britton, C.M. 651
Brock, S.M. 1151
Broido, A. 664, 670
Brown, D.B. 1201
Brown, H.D. 468
Brown, H.E. 1130, 97
Brown, J.L. 1398
Brown, J.M. 127, 128, 58, 59
Brown, S.J. 407
Brownlee, R.G. 419
Broyer, T.C. 766, 767
Bruce, D. 892, 913
Bruchet, G. 865
Brush, R.O. 183
Buchanan, H 154
Buchanan, H. 222
Buckman, R.E. 926
Bulgrin, E.H. 1237, 1304
Bultena, G.L. 267
Bultman, J.D. 444, 447
Burdge, R. 268
Burdsall, H.H. 607
Burdsall, H.H., Jr. 485
Burgess, R.L. 269, 270
Burkart, L. 207
Burke, J.D. 1200, 1213
Burmeister, E.L. 1253
Burns, R.M. 793, 794, 860
Burroughs, E.R., Jr. 105
Burton, J.D. 1223, 893
Bynum, H.H. 556, 606
Byrd, V. L. 1316
Byrd, V.L. 1348

C

Calvin, F.B. 1184
Campbell, C.J. 60
Campbell, R.E. 61

Campbell, R.K. 795, 956
Campbell, R.W. 301, 370
Campbell, W.A. 483
Carl, C.M., Jr. 957
Carmean, W.H. 764
Carolin, V.M. 302
Carolin, V.M., Jr. 328
Carpenter, E.M. 1152
Carpenter, L.H. 185
Carter, F.L. 432, 433, 434, 435
Carter, M.R. 1201
Cathey, R.A. 1041
Caulfield, D.F. 1317
Chalupka, W. 722
Chang, W.M. 984
Chase, T. 472
Chern, J. 1365
Chibbar, R.N. 859, 880
Chien, M. 1034, 1035
Choate, G.A. 1069
Christopher, J.F. 1042, 1056
Chudnoff, M. 1221, 1224
Church, T.W. 1175
Church, T.W., Jr. 1183
Clark, A. 1225, 1290, 907
Clark, E.W. 329, 385
Clark, K.A. 245
Clark, R.N. 269, 270, 271, 290
Clark, W.M. 773
Clausen, K.E. 993
Claxton, H.D. 1396, 1404
Clayton, J.L. 1, 129
Clayton, R.N. 1398
Clements, H.B. 665
Clements, R.W. 943
Clutter, J.L. 933
Cochran, P.H. 861, 862
Cole, D.M. 914
Conder, T.M. 1210
Conkle, M.T. 994, 995
Conrad, L.A. 1226
Conrad, P.W. 186
Consul, P.C. 688
Cook, D.I. 702, 703, 704, 707
Cooley, J.H. 743, 796
Cooper, G.A. 1184, 1227, 1228, 1229, 741
Cooper, R.W. 625, 626, 627, 628, 629, 646
Copeland, O.L. 62
Copes, D.L. 958, 996
Cordell, H.K. 272
Corlett, R.C. 673
Cormier, P.L. 283
Coulter, W.K. 302
Countryman, C.M. 641, 651, 666, 667
Coyne, J.F. 959
Craft, E.P. 1185
Cramer, O.P. 689
Crawford, J.L. 698
Cremmen, J. 482
Critchfield, W.B. 948
Croley, T.E. 561
Crosby, J.S. 669
Crouch, G.L. 855
Cupal, J. 243
Cupal, J.J. 242
Cuppett, D.G. 1143, 1185
Currie, P.O. 156
Currier, P.O. 194
Curtis, R.O. 894
Curtis, W.R. 18, 2, 63
Cuthbert, R.A. 393

D

Dahms, W.G. 895, 896
Dale, M.E. 897
Dana, R.W. 1080, 1081, 1082, 425, 426, 427
Darr, D.R. 1153, 1154
Darwin, W.N. 440
Darwin, W.N. Jr. 1029
Daterman, G.E. 328, 330, 396
Daves, G.D., Jr. 396
Davidson, H.L. 1230, 1236
Davidson, R.W. 493, 612
Davidson, W.H. 64
Davis, C.J. 363
Davis, G. 64
Davis, J.R. 172, 672
Davister, M.D. 1384
De Graaf, R.M. 244
De Groot, R.C. 572, 608, 609, 615
Dealy, J.E. 187
DeBald, P.S. 1100
DeBano, L.F. 3
DeBarr, G.L. 397
DeBell, D.S. 723, 742
DeBower, K. 1126
DeByle, N.V. 19
DeGraaf, R.M. 183
Dell, J.D. 630, 662
Dell, T.R. 69
Della-Bianca, L. 765, 844, 889
Demeritt, M.C., Jr. 984
Dempster, L.J. 750
Denton, R.E. 310, 398
Derr, H.J. 990
Deshpande, V.H. 1318
Dhruva, B.R. 1319
Dickerhoof, H.E. 1165
Dickerson, A.L. 273
Dickerson, B.P. 65
Dickinson, J.O. 20
Dickson, R.E. 766, 767, 809, 877
Dietschman, G.H. 843
Dietz, D.R. 188, 238
Dinus, L.A. 433, 434
Dinus, R.J. 554, 985
Dixon, R.A. 258
Dobhinger, L.S. 451
Dochinger, L.S. 450, 452, 484, 495, 870
Donaldson, B.A. 215
Doolittle, M.L. 631
Doolittle, R.E. 376
Dorman, K.W. 1018, 901
Doty, R.D. 66, 84
Douglass, J.E. 67, 68, 81
Driscoll, R.S. 155, 156, 157, 158, 159, 161, 227
Drooz, A.T. 331, 332
Duckles, C.K. 299
Duff, J.E. 1366
Duffy, I.T. 217
Duffy, P.D. 12, 47, 69
Duncan, D.A. 218
Dunmire, D.E. 1389
Duran, G. 1124
Dutrow, G.F. 1134
Dwinell, L.D. 573
Dyrness, C.T. 4, 70, 73, 751,

E

Earles, J.M. 1043, 1044, 1045
Ebel, B.H. 303, 313, 333, 350
Echols, R.M. 1024, 845
Edgar, J.C. 1210
Edinger, J.G. 453, 619
Edminster, C.B. 923
Effland, M.J. 1335
Elsner, G.H. 260, 279
Emanuel, D.M. 1143
Engelken, J.H. 287
Enghardt, H.G. 898, 902
Erdmann, G.G. 498
Erickson, B.J. 1231
Erickson, J.R. 1195, 1202, 1203, 1204, 1205, 1216
Eslyn, W.E. 618
Ethington, R.L. 1357, 1367
Evans, B.B. 1352
Evans, H.J. 541
Evans, K.E. 160
Evans, L.S. 454

F

Fahey, D.J. 1320, 1331
Fahey, T.D. 1232
Fahn, A. 863, 947
Farmer, E.E. 71
Farnsworth, D.H. 705
Fasick, C.A. 1165
Fatzinger, C.W. 334
Fedde, G.F. 335, 336, 399
Fedde, V.H. 331
Feddema, C. 189
Federer, C.A. 28, 72, 75, 92
Feist, W.C. 1321
Fellin, D.G. 337, 400
Ferguson, E.R. 768, 769
Ferguson, R.B. 190, 203
Ffolliot, P.F. 915
Ffolliott, P.F. 1220
Filer, T.H. 595
Finn, R.F. 776
Firth, J.E. 918, 919
Fischer, V.L. 285
Fischer, W.C. 652
Fleischer, H.O. 1367
Fletcher, J.E. 71
Florez, A 191
Foiles, M.W. 580, 899
Folkman, W.S. 632, 633, 634
Fosberg, M.A. 653, 690
Foulger, A.N. 1233, 1234
Fowler, R.F. 401, 402
Fowler, W.B. 85
Francis, R.E. 157, 158, 161, 176
Frandsen, W.H. 668
Frank, R.M. 846
Franklin, E.C. 960, 997
Franklin, J.E. 395
Franklin, J.F. 1030, 73, 742, 749, 750, 751
Fredriksen, R.L. 142
Freese, F. 1234, 1244, 1285
Frischknecht, N.C. 152, 162, 163, 184
Frye, R.H. 338
Funk, D.T. 1001
Fuquay, D.M. 687, 691
Furman, R.W. 654
Furniss, M.M. 339, 340, 341, 351, 403, 415
Furniss, R.L. 339

G

Gabriel, W.J. 935, 961
Gaby, L.I. 1369, 1370
Galford, J.R. 342, 348, 404
Galoch, E. 1007
Gammon, G.L. 1286
Gansel, C.R. 944
Gansner, D.A. 1106, 1109, 927
Gara, R.I. 714
Garcia, G. 50
Gardner, R.B. 1070, 1201, 1206, 1207
Garrett, P.W. 986, 987
Garrison, G.A. 192
Gartner, F.R. 164, 249
Gary, H.L. 29, 74
Gatherum, G.E. 705
Geary, T.F. 797, 798, 799, 800
Gee, G.W. 75
Geist, J.M. 193
George, C.W. 655
George, R.P. 1397
Gerardo, M. 991
Gerdemann, J.W. 558
Gerhards, C.C. 1371, 1379
Gerhold, H.D. 984
Geyer, W.A. 900
Gibbs, C.B. 936, 998
Gilbert, C.G. 261
Gilbertson, R.L. 485, 612
Gill, T.G. 1155
Gillespie, R.H. 1235, 1372
Gillette, N.L. 413
Gingrich, S.F. 1106, 927
Gjovik, L.R. 1230, 1236, 1263, 613
Glover, R.P. 1156
Gochenour, D.L., Jr. 1209
Godman, R.M. 1033

Godshall, W.D. 1322, 1328, 1373, 1377
Goebel, N.B. 113
Goho, C.D. 1157
Gonsior, M.J. 1208
Goodin, J.R. 202
Goolsby, R.P. 405
Gowin, T. 962, 980
Goytia, E. 1224
Graney, D.L. 768, 769
Granskog, J.E. 1135
Grantham, J.B. 1374, 1375
Gratkowski, H. 801, 847
Grechius, Y.A. 250
Green, J. 1344, 1345, 1346
Green, J.E., III 682
Green, L.R. 635
Greentree, W.J. 1078, 1079, 423, 424
Gregory, G.F. 515, 574, 585
Gregory, R.A. 685, 733, 963, 964
Grelen, H.E. 165
Griffin, J.R. 948
Grigsby, H.C. 802
Grimble, D.G. 343
Grisez, T.J. 1115
Grosenbaugh, L.R. 916, 928
Gunter, L.E. 642
Guthrie, W.A. 219
Guttenberg, S. 1107, 1108, 1136
Gysel, L.W. 743

H

Hacskaylo, E. 864, 865
Hadley, H.H. 799
Haeffner, A.D. 76
Haferkamp, M.R. 194
Hafley, W.L. 381
Hahn, J.T. 1071
Haines, D.A. 669, 675
Hair, D. 1158
Haissig, B.E. 1031, 486, 487, 540, 866, 867
Hajny, G.J. 1321
Hall, F.C. 122, 73, 751
Hall, R.H. 488, 561
Halliwell, R.H. 561
Halliwell, R.S. 478
Hallock, H. 1237, 1272
Halls, L.K. 195, 196
Halpern, Y. 670
Halverson, H.G. 30, 36, 42, 43, 44
Hamilton, J.E. 83
Hamlen, R.A. 421
Hanks, L.F. 1238
Hann, D.W. 1070, 1206
Hann, R.A. 1239, 1276, 1304
Hanover, J.W. 977
Hansen, B.G. 1156
Hansen, E.A. 112, 292
Hardy, C.E. 652
Hare, R.C. 988
Harkin, J.M. 1323
Harlan, J.C. 228
Harlow, R.F. 196, 219
Harms, W.R. 868
Harpole, G.B. 1144
Harris, A.R. 31, 39, 77
Harris, A.S. 724
Harris, J. 1233, 1234
Harris, J.F. 1324
Harris, L.E. 163
Hartman, G.B. 920
Hartsog, W.S. 1207, 1208
Harvey, A.E. 489, 490, 530
Hatchell, G.E. 770, 901
Haupt, H.F. 40
Hawe, R.G. 691
Hawksworth, F.G. 467, 575, 582, 929
Hay, C.J. 344, 345
Hazard, J.W. 1083
Healy, W.M. 220
Hebb, E.A. 793
Hedlund, A. 1043, 1044, 1045
Heebink, B.G. 1240, 1241, 1242
Heebink, T.B. 1354, 1374, 1375
Heede, B.H. 78, 79
Hefty, F.V. 1242
Heidmann, L.J. 856
Heisler, G.M. 83
Hejnowicz, Z. 1025, 965
Helfman, R.S. 654
Heller, R.C. 1084, 1085, 1086, 1087, 1097, 304, 312, 428, 429, 461
Helvey, J.D. 113, 80, 81
Hendee, J. 268
Hendee, J.C. 267, 269, 270, 271, 274, 289, 290
Henderson, J.H. 1398
Hendrix, F.F. 483
Hendrix, J.W. 491
Henley, J.W. 1243, 1259, 1260, 1261
Hepting, G.H. 492
Herman, F.R. 917, 918, 919
Hermann, F.J. 197, 198, 199
Herring, J.L. 346
Herrington, L.P. 82, 83
Hewlett, J.D. 81
Hibbert, A.R. 32
Highley, T.L. 616, 617
Hildebrandt, G. 1087, 429
Hiller, C.H. 1244
Hillstrom, W.A. 1196
Hinds, T.E. 493
Ho, I. 567
Hodge, K.L. 682
Hodges, C.S. 494
Hodgson, E. 380
Hoff, R.J. 1019, 514, 571, 576, 577, 584, 982
Hogan, W.G. 1151
Holmgren, R.C. 166, 200
Holt, E. 1119, 1159, 1160
Hook, D.D. 808
Hooper, R.M. 811
Hopkins, D. 801
Horn, R.A. 1325, 1326
Hornbeck, J.W. 28, 34, 72
Host, J.R. 1161
Hough, W.A. 636, 637, 671
Houston, D.R. 462, 463, 563, 564
Howard, D.A. 230
Howard, J.O. 1088, 638
Howe, V.K. 352
Hoyle, M.C. 771, 869
Hse, C.Y. 1245, 1246, 1247, 1248
Hubbard, H.B., Jr. 406
Hudler, G.W. 516, 517
Hungerford, R.D. 340, 581
Hunt, D.L. 1232
Huntzinger, H.J. 803
Hutchin, C.C. 1072
Hutchings, S.S. 200
Hutchinson, J.G. 275

I

Ingebo, P.A. 32
Iqbal, N. 526
Irland, L.C. 1125, 1137
Isebrands, J.G. 1026, 1027, 876, 877

J

Jackson, M.L. 1398
Jacobsen, G.L. 1207
Jacobson, M. 396
Jain, M.K. 1011, 879, 999
Jamal, S.M. 569
James, G.A. 272, 276, 277, 278
Jameson, D.A. 167
Jeffers, R.M. 966
Jenkins, R.E. 1030
Jennings, D.T. 347
Jensen, K.F. 495, 528, 529, 870
Jewell, F.F. 578
Johansen, R.W. 656
Johnsen, T.N., Jr. 725
Johnson, C.M. 766
Johnson, E.J. 1249
Johnson, F.A. 920
Johnson, H.E. 581
Johnson, L.C. 1000, 967
Johnson, L.R. 1198, 1209
Johnson, P.H. 560
Johnson, V.J. 645
Johnston, H.R. 445, 446, 448
Johnston, R.S. 84
Johnston, W.E. 279
Johnston, W.F. 726, 804
Jokela, J.J. 705
Jokerst, R.W. 1250, 1251, 1291, 1292
Jones, E.P., Jr. 716
Jones, J.R. 1032, 871
Jones, R.A. 43, 44
Jones, R.E. 1376
Jones, R.K. 1164
Jones, R.R. 371
Jones, T.W. 371, 496, 574
Jorgansen, C.D. 184
Jorgensen, E. 1342
Julien, L.M. 1210
Jung, P. 1252
Juntunen, E.T. 21
Jurkovic-Bevilacqua, B. 1023

K

K.E. Severson, K.E. 237
Kaeiser, M. 1001
Kais, A.G. 497, 554
Kaiser, H.F. 1124, 1126, 1138, 1139
Kallio, E. 1033, 1168, 1168a, 1186, 1187
Kamensky, S.A. 498
Kamienska, A. 1002
Kaplan, R. 253, 255
Kaplan, S. 254, 255
Keith, G.C. 959
Kennedy, B.H. 348
Kennedy, H.E., Jr. 805
Kennedy, P.C. 337
Kerbs, R.R. 160
Kessler, K.J., Jr. 464, 499, 500
Khan, A.A. 708
Khan, A.S. 569
Khan, M.A. 709, 711, 837
Khan, M.S. 712
Khan, S.N. 781, 882
Kidd, W.J. 8
Kimbrough, J.W. 559
Kinelska, J. 888
King, J.G. 262
Kingsley, N.P. 1066
Kinloch, B.B. Jr. 989
Kinney, R.E. 1253, 1289
Kirk, T.K. 1327, 610
Kirscher, H. 872
Klemmedson, J.O. 120, 121
Kline, L.N. 403, 415
Klock, G.O. 122, 5, 85
Klungness, J.H. 1332, 1333
Knight, F.B. 343, 354, 364, 365, 366, 367, 368
Knight, H.A. 1046, 1047
Knipe. O.D. 86
Knipe, O.D. 87
Knutson, D.M. 465, 501, 502
Koch, P. 1249, 1254, 1255, 1256, 1257, 1258
Koenigs, J.W. 503, 504
Koeppen, R.C. 1226
Koerber, T.W. 326
Kojima, M. 256
Kolakowska, M. 970
Konar, R.N. 1003
Koning, J.W. 1322, 1328, 1334, 1373, 1377
Kopcewicz, J. 1008
Koplin, J.R. 349
Kormanik, P.P. 1004, 1005, 808
Kovner, J. 90

547-679 O - 74 - 8

Kowal, R.J. 303, 350
Krajicek, J.E. 873
Krammes, J.S. 137
Krebill, R.G. 341, 505, 506, 507
Kriebel, H.B. 705
Kriesel, K. 1008
Krochmal, A. 1034, 1035
Krueger, W.C. 201, 222
Kucharska, J. 888
Kuenzi, E.W. 1328, 1365, 1377, 1380
Kuhlman, E.G. 508, 579
Kundaeli, J.N. 221

L

Laemmlen, F.F. 509
Lafarge, T. 968
Lamb, R.C. 692, 806
Landis, M.W. 1399
Landt, E.F. 1036, 1228, 1400
Lane, P.H. 1259, 1260, 1261
Langdon, O.G. 770, 868, 901
Lanier, C.N. 351
LaPage, W.F. 280, 281, 282, 283
Larsen, D.N. 1109
Larsen, M.J. 611
Larson, F.R. 96
Larson, J.S. 244
Larson, P.R. 757, 874, 875, 876, 877
Laseter, J.L. 510, 511, 565
Lason, F.R. 1116
Laundrie, J.F. 1329, 1330, 1331, 1332, 1333
Lauterbach, P. 801
Lautz, W. 481
Lawler, G.W. 510
Lawrence, J.D. 1162, 1163, 1164, 1165, 1166
Laycock, W.A. 154, 222
Leaf, A.L. 25
Leaf, C.F. 33, 88, 89
Leaf, C.G. 90
Leak, W.B. 752, 753, 754
Leaphart, C.D. 580, 581
Leary, R.A. 1127
Lege, F.O. 1166
Lehmann, W.F. 1242, 1262
Leiser, A.T. 286
Leith, H. 780
Lennartz, M.R. 168
Leonard, R.E. 25, 82, 83, 91, 92
Leonard, R.G. 180
Lessard, R.A. 470
Lester, D.T. 705
Leuschner, W.A. 1098, 1110, 1111, 1112
Lewis, C.E. 169
Lewis, F.B. 406, 421
Lewis, W.C. 1235, 1372
Li, C.Y. 22, 512
Lightle, P.C. 467, 582, 929
Likens, G.E. 24
Lime, D.W. 261, 263, 275, 284
Lindell, G.R. 1167, 1168, 1168a
Lindenmuth, A.W., Jr. 672
Lindmark, R.D. 1184, 1187, 1229, 741
Little, E.L., Jr. 949
Little, S. 785
Litton, R.B., Jr. 257
Liu, T. 950
Lloyd, R.D. 285
Lockard, R. 1126
Lohr, C.J. 1067
Lohrey, R.E. 165, 807
Lombard, F.F. 612
Look, M. 297, 299
Loper, B.R. 26
Lorenz, L.F. 1263, 613
Lorio, P.L., Jr. 352
Losche, C.K. 772, 773, 774
Lotocki, A. 980
Lowery, D.P. 1264, 1265, 1266
Lu, K.C. 22, 512
Lucas, B.A. 353
Lucas, R.C. 264
Lull, H.W. 130, 93
Lundgren, A.L. 1099
Lutz, J.F. 1267
Lyon, R.L. 299, 353, 407, 409, 413, 414
Lysons, H.H. 1211, 1212

M

Mace, A.C., Jr. 262
MacKenzie, A.P. 1397
MacLean, C.D. 1089, 1090
Maeglin, R.R. 1271, 1282
Magill, A.W. 286
Mahmood, T. 527
Main, W.A. 639, 669
Malcolm, F.B. 1272
Mann, C.N. 1212
Mann, J.F. 230
Mann, W.F. 898, 902
Manwiller, F.G. 1273, 1274, 1275
Marburger, R.G. 245
Marcin, T.C. 1169
Markstrom, C.D. 891
Markstrom, D.C. 1032, 1231, 1276, 1277, 1279
Marlatt, W.E. 690
Marquis, D.A. 1006, 1113, 745
Marsden, M.A. 952
Martignoni, M.E. 308, 395
Martin, A.J. 1157, 1170, 1188
Martin, C.N. 352
Martin, C.W. 24, 34
Martin, N.E. 491, 566
Martin, R.E. 682, 698
Martin, S.C. 210, 223, 224, 225
Martin, W. 1133
Martinelli, M., Jr. 131, 132, 35, 94
Marx, B. 1081, 426
Marx, D.H. 533, 586, 587, 588
Maser, C. 73, 751
Mason, L.L. 788
Mason, R.R. 358, 420
Massengale, R. 1065
Massey, H.F. 143, 23
Mata, S.A., 309
Matthews, F.R. 589
Mauldin, J.K. 436, 438
Maurice, S.C. 283
Maxwell, R.K. 246
May, J.T. 826
May, L.C. 494
Mayer, C.E. 1040
Mazzone, H.M. 395
McAlpine, R.G. 1004, 1005, 1290, 808, 826, 907
McBride, J.R. 449, 809
McCain, A.H. 466, 600
McCambridge, W.F. 354, 355, 408
McCammon, B.P. 657
McCauley, O.D. 1113, 1114
McCay, R.E. 1100
McClellan, G.E. 642
McClure, J.P. 1048
McCluskey, R.A. 583
McConnen, R.J. 234
McCracken, F.I. 314, 513
McCutchan, M.H. 453, 456, 457, 619, 620, 621
McDonald, G.I. 514, 576, 577, 584
McDonald, G.I.(Eds.) 571
McDonald, S. 788
McDonough, W.T. 186, 191
McGee, C.E. 744, 765
McGinnes, E.A. 1309
McGinnis, G.D. 676
McKee, W.H., Jr. 878
McKell, C.M. 202
McKnight, M.E. 305, 356
McLane, W.H. 411
McLemore, B.F. 810
McManus, M.L. 306, 307, 357, 369
McMillen, J.M. 1268
McMillin, C.W. 1269, 1270, 1308
McMinn, J.W. 168
McNab, W.H. 721
McNatt, J.D. 1378
McNees, R.E. 1018
McReynolds, R.D. 794
McWain, P. 515, 574, 585
Medin, D.E. 148, 149, 150, 203, 213, 214
Medin, D.E., Bowden, D.C. 151
Mees, R.M. 658
Megaham, W.F. 8
Megahan, W.F. 10, 104, 6, 7, 95
Mendel, J.J. 1115
Mergen, A.F. 1283
Merkel, E.P. 303, 397
Merritt, C. 705
Mersereau, R.C. 9
Merton, J.R. 218
Messner, H.E. 238
Michniewicz, M. 1007, 1008
Mickelson, A.S. 991
Mikell, H.K. 656
Miller, P.R. 449, 453, 454, 455, 456, 457, 458, 459, 619, 620, 621
Miller, R.L. 1116, 96
Miller, T. 590
Miller, W.C. 1171, 1177
Millers, I. 401
Millett, M.A. 1335, 1379
Milligan, H.P. 456, 620
Milligan, R.P. 458
Miniutti, V.P. 1278
Minore, D. 700, 701, 727
Miskus, R.P. 318
Mobley, H.E. 640
Moeller, G.H. 287, 288
Mohaupt, A.A. 1334
Mohn, C.A. 969
Monroe, R.J. 381
Montrey, H.M. 1380
Moody, R.C. 1291, 1292, 1328, 1377, 1381, 1382
Moore, D.G. 144, 145, 26
Moore, G.E. 359, 360, 361
Moore, W.E. 1327, 1335, 610
Morris, M.J. 156, 185
Morrison, D.A. 288
Moser, J.C. 362, 417, 419
Mueggler, W.F. 170
Mueller, L.A. 1279
Mujumdar, R.B. 1336
Murmanis, L.L. 1280
Murphy, J.D. 984
Murphy, P.A. 1049, 1050
Myers, C.A. 467, 903, 921, 922, 923, 929
Myers, G.C. 1337, 1338

N

Nalborczyk, E. 970, 979
Namkoong, G. 1009, 971
Nanda, K.K. 1010, 1011, 859, 879, 880, 999
Navon, D.I. 1128, 133
Naylor, A.W. 723
Neal, R.L., Jr. 930, 931
Neel, W.W. 377
Nelson, E.E. 606
Nelson, N.D. 1281
Nelson, R.E. 363
Nelson, R.M., Jr. 673
Newhouse, M.E. 779
Newsome, L. 315, 440
Nicholls, T.H. 468, 469, 516, 517, 550
Nienstaedt, H. 972
Noble, D.L. 728, 729
Nobles, R.W. 800
Nolley, J.W. 1189

Nord, E.C. 641, 642
Nord, J.C. 343, 364, 365, 366, 367, 368, 369
Norick, N.X. 1082, 1091, 427, 430
Norris L.A. 145
Norris, L.A. 21
Noste, N.V. 659, 660
Nyland, R.D. et al 1190

O

O'Brien, J.T. 551
O'Connell, P.F. 1129, 1130, 1131, 97
O'Gwynn, C.H. 397
Okkonen, E.A. 1282
Oliver, W.W. 775, 904
Olson, D.F., Jr. 811
Orr, H.K. 125, 134, 98
Ostmark, H.E. 325
Ostrom, A.J. 1051
Ostrom, C.E. 757
Oswald, D.D. 1052
Oviatt, A.E. 1354, 1383
Owston, P.W. 36, 812
Ozment, A.D. 118

P

Packer, P.E. 135, 19, 62
Page, M. 409, 414
Paine, D.P. 1092
Paine, L.A. 604
Palmer, T.Y. 693
Panzer, H.R. 1283
Parker, H.D., Jr. 226, 227, 228
Parker, J. 518, 519, 564
Parmeter, J.R., Jr. 520
Pase, C.P. 60, 99
Patric, J.H. 100
Patton, D.R. 171
Paul, J.T. 694
Pauley, S.S. 705
Payne, B.R. 1117, 251
Payondeh, B. 1092
Pearson, H.A. 172, 229, 230, 232
Pearson, H.A., Whitaker, L.B. 231
Peevy, F.A. 848, 849, 850, 851
Pendleton, T.H. 1151, 1191
Pennington, R.E. 179
Perala, D.A. 746
Percival, R.M. 659, 660
Perla, R.E. 136
Perry, T. 321
Peters, C.C. 1283, 1291, 1292, 1382
Peters, J.K. 881
Peters, P.A. 1213
Petersen, L.E. 237
Peterson, G.L. 261
Peterson, G.W. 521, 522, 591
Peterson, R.S. 523, 524
Pfister, R.D. 843
Phares, R.E. 1036, 1400, 774, 776, 813
Phelps, R.B. 1155
Phelps, W.R. 496, 595
Phillips, C.B. 642
Phillips, D.R. 1225, 1284
Philpot, C.W. 676
Phipps, H.M. 814
Pieper, G.R. 298
Pierce, D.S. 1320
Pierce, K. 825
Pierce, R.S. 24, 28, 72
Pitcher, J.A. 820
Plank, M.E. 1259, 1260, 1261
Plass, W.T. 101, 102
Plumb, T.R. 623, 643
Plummer, A.P. 204
Podgwaite, J.D. 370
Pomeranz, Y. 1401, 1402, 1403
Pond, F.W. 173, 174, 232
Pong, W.Y. 1307
Pope, R.B. 1092
Popham, T.W. 992
Porterfield, E.J. 792
Potter, D.R. 289, 290
Powell, W.M. 483
Powers, H.R. 605
Powers, H.R., Jr. 525
Powers, R.F. 777, 778
Price, D.A. 154
Puri, Y.N. 477
Putman, J.W. 1126

Q

Quinkert, A.K. 276
Quirk, J.T. 1285
Quraishi, M.A. 526, 527

R

Radwan, M.A. 855, 857, 858
Ragain, D.P. 282
Rainey, D.A. 657
Raju, P.N. 541
Ram Reddy, M.A. 477, 781
Rama Rao, A.V. 1318, 1319, 1339, 1340
Ramaker, T.J. 1384
Randall, R.M. 1101
Randall, W.K. 969
Rango, A. 690
Raper, K.B. 1397
Rasmussen, L.A. 410
Rast, E.D. 1286
Rathi, S.S. 1339, 1340
Ratliff, R.D. 175, 205, 233, 234
Ravindranath, B. 1341
Read, R.A. 705, 706
Reardon, R.W. 411
Reddy, M.A.R. 882
Reeves, C.C. 24
Regelin, W.L. 252
Reichert, D.W. 252
Reid, W.H. 1176
Reinhart, K.G. 130, 146, 93
Reppert, J.N. 161, 176, 234
Reukema, D.L. 905
Rex, R.W. 1398
Rexrode, C.O. 371
Reynolds, H. G. 177
Reynolds, H.G. 215, 221, 235, 236
Reynolds, P.E. 528, 529
Reynolds, R.R.J. 644
Rice, R.M. 10, 103, 104, 137, 3
Rich, L.R. 37
Richardson, B.Z. 105
Richmond, C.E. 353, 412
Rickman, J.T. 530
Riensi, G. 1396, 1404
Rier, J.P. 531
Rietz, R.C. 1287
Riffer, R. 670
Riffle, J.W. 532
Ringger, D. 178
Roberts, D.R. 945
Roberts, R.B. 299
Roberts, R.W. 205
Robertson, J.L. 353, 407, 413, 414
Robinson, R.M. 245
Rogers, J.J. 106
Rogers, L.L. 246
Rogers, M. 206
Rogers, N.F. 900
Romancier, R.M. 1030
Romberger, J.A. 1025, 963, 964
Ronco, F. 815, 816, 883
Rosen, H.N. 1288
Ross, E.W. 533, 592
Roth, H.G. 1236
Roth, L.E. 606
Rothacher, J. 107
Rothacher, J.S. 10, 104
Rothermel, R.C. 668, 674
Roton, L.M. 362
Roussopoulos, P.J. 639, 645
Rowan, S.J. 534, 589, 593, 594, 595, 817
Rowe, J.W. 1342
Rowell, R.M. 1343, 1344, 1345
Roy, D.N. 1342
Rudinsky, J.A. 403, 415
Rudolph, T.D. 1012, 757, 884
Ruehle, J.L. 405, 535, 536, 537
Russell, T.E. 818
Ruth, J.M. 419
Ruth, R.H. 937, 938
Ryan, B.C. 453, 457, 619, 621

S

Sachs, I.B. 1253, 1280, 1289, 1397, 1398, 1401, 1402, 1403
Sachs, T. 863, 872
Sackett, S.S. 646
Sadowski, F.G. 1096, 431
Safford, L.O. 755
Sajdak, R.J. 705
Sampson, G.R. 1172
Sander, I.L. 730
Sanderson, H.R. 247
Sando, R.W. 647, 675
Sanford, G.R., Searcy, A. 278
Sarles, R.L. 1145
Sartz, R.S. 38, 39
Sassaman, R.W. 1102, 1118, 1119, 1120, 1132, 248, 906
Satterlund, D.R. 40
Saucier, J.R. 1290, 1405, 907
Saylor, L.C. 967
Schaffer, E.L. 1291, 1292, 1385
Schallau, C.H. 1102
Scharpf, R.F. 386, 520, 538
Schier, G.A. 108, 109, 110, 111
Schillings, P.L. 1214, 1215
Schipper, A.L., Jr. 486, 487, 539, 540, 867
Schlaegel, B.E. 747
Schlesinger, R.C. 1184, 1229, 741, 852
Schmid 338
Schmid, C.W. 691
Schmid, J.M. 372, 416
Schmidt, R.A., Jr. 41, 46
Schmidtline, R.C. 819
Schmidtling, R.C. 1013
Schmitz, R.F. 351, 373, 403, 415
Schneeweis, J.C. 237
Scholes, V.E. 478, 561
Schreiber, L.R. 404
Schreiner, E.J. 1014
Schreuder, H.T. 277, 332
Schroeder, J.G. 1284, 1406
Schroeder, M.J. 453, 619
Schubert, G.H. 172, 725, 820
Schultz, R.P. 822, 823, 908, 973
Schumann, D.R. 1293, 1294, 1302
Schweitzer, D.L. 1102, 1120
Scott, R.W. 1346
Scott, V.E. 171, 215
Seegrist, D.G. 393
Seegrist, D.W. 452, 935
Segelquist, C.A. 179, 180, 206
Seidel, K.W. 756
Seidler, R.J. 541
Seikel, M.K. 1342
Seliskar, C.W. 542
Sendak, P.E. 1173, 1174
Seshadri, T.R. 1341, 1347
Setzer, T.S. 1073
Severinghaus, C.W. 291
Severson, K.E. 164, 238, 249, 250
Shafer, E.L., Jr. 291
Shafizadeh, F. 676
Shah, M.A. 710, 711, 837
Sharon, E.M. 543

Sharpe, K.M. 289, 290
Shaw, C.G. 530
Shea, K.R. 544, 596
Sheehan, D. 696
Shenton, L.R. 688, 695, 696
Shepperd, W.D. 924
Sherwood, G.E. 1386, 1387, 1388
Sherwood, J.K. 1068
Shigo, A.L. 470, 471, 472, 492, 531, 545, 546, 547, 548, 597, 598, 599
Shon, F.L. 413, 414
Short, H.L. 180, 207, 239
Shoulders, E. 824
Shubert, G.H. 821
Siegel, W.C. 1121, 1122
Silen, R.R. 971
Silverstein, R.M. 419
Simpson, W.T. 1295
Singh, S. 476, 477
Skilling, D.D. 469, 517, 549, 550, 551
Skog, K.E. 1127
Skolmen, R.G. 1296, 1297
Skovlin, J.M. 208
Skutt, H.R. 470
Sluder, E.R. 974, 975
Smalley, G.S. 825
Smith H.R. 357
Smith, A.H. 552
Smith, C.E. 918, 919
Smith, D.M. 1223, 1244, 1285, 893
Smith, D.W. 140
Smith, F.W. 138
Smith, H.C. 936, 939, 940, 941, 942
Smith, J.G. 1118, 248, 906
Smith, J.L. 36, 42, 43, 44
Smith, L.R. 693
Smith, R.H. 374
Smith, R.S., Jr. 466, 553, 600
Smith, V.K. 443, 445, 446
Smith, W.E. 1348
Smith, W.H. 528, 529
Smity, C.E. 938
Smythe, R.V. 432, 433, 434, 435, 436, 437, 438, 439
Snow, A.G., Jr. 939, 940
Snow, C.A. 578
Snow, G.A. 554
Snyder, E.B. 1015, 1016, 990
Solomon, J.D. 375, 376, 377, 440
Solomon, M.J. 112, 292
Sommerfield, R.A. 45
Sonderman, D.L. 1286
Sonnet, P.E. 417
Sorensen, F.C. 1017
Southwell, C.R. 444, 447
Spada, B. 1086, 428
Spencer, J.S., Jr. 1053
Spencer, M.M. 159
Spotts, R.A. 555
Springer, E.L. 1321, 618
Springfield, H.W. 87
Squillace, A.E. 1018, 881, 997
Srago, M.D. 600
Srinivasan, R. 1318, 1319, 1336
Stage, A.R. 717, 914
Staley, J.M. 555, 556
Stankey, G.H. 265
Statler, M.W. 411
Stearns, F. 178, 743
Stein, A.M. 677
Stein, W.I. 812
Stein, W.J. 1037
Steinbeck, K. 826
Steinhilb, H.M. 1216
Steinhoff, R.J. 1019, 951, 982
Stern, E.G. 1192, 1193
Stern, R.K. 1389
Sternitzke, H.S. 1042, 1054, 1055, 1056
Stevens, R.E. 327, 378, 418
Stewart, H.A. 1298, 1299
Stewart, J.L. 596
Stewart, R. 847
Stockmann, V.E. 1349
Stockstad, D.S. 678
Stringer, C.A. 435
Strothmann, R.O. 827
Struble, G.R. 379
Sturos, J.A. 1217, 1218
Stutzbach, S.J. 25
Sutton, R.F. 785
Swain, A.M. 266
Swank, W.T. 113, 68
Swanson, R.H. 114
Swanston, D.N. 11
Swindel, B.F. 381, 912, 925, 932
Syers, J.K. 1398

T

Tabler, R.D. 115, 46
Tanner, W. 1252
Tarrant, R.F. 116, 117, 147, 26, 601
Tarver, J.W. 1125
Tattar, T.A. 471, 472
Taylor, A.R. 691
Taylor, J. 634
Taylor, J.F. 380
Thapar, H.S. 781
Thatcher, R.C. 361
Thilenius, J. F. 182
Thilenius, J.F. 181
Thill, R.E. 250
Thomas, H.A. 381
Thomas, J.W. 183, 245, 251, 258
Thomas, R.J. 1310, 1311, 1312
Thompson, C.G. 420
Thompson, E.F. 1099
Thompson, J.R. 118
Thomson, F.J. 1096, 431
Thornbury, J.R. 310
Thorne, H.W. 1053
Tiech, A. 972
Tiedemann, A.R. 119, 120, 121, 122
Tigner, T.C. 358
Timson, F.G. 1175
Tingey, W.M. 184
Toevs, W.J. 502
Torgersen, T.R. 382
Townsend, A.M. 452
Trappe, J.M. 117, 22, 512, 552, 557, 558, 559, 567, 601, 602, 603
Trimble, G.R. 100, 1114, 1115
Trimble, G.R., Jr. 853
Trostle, G.C. 355
Tschernitz, J.L. 1291, 1292, 1300
Tu, C.M. 17
Tubbs, C.H. 699
Tucovic, A. 1020, 1021
Tumlinson, J.H. 419
Tuomi, R.L. 1390
Turkowski, F. 236
Turner, G.T. 209, 240
Turner, J.C. 637
Tyre, G.L. 293, 294

U

U.S. Department of Agriculture Forest Service 1057
U.S. Department of Agriculture, Forest Service 748
U.S.D.A. Forest Service, Forest Products Laboratory 1368
Ulrich, A.H. 1158
Underwood, J.C. 937, 938
Urness, P.J. 241
Ursic, S.J. 12, 47
Usanis, R.A. 971
USDA Forest Service 661
USDA Forest Service, Northeastern Forest Experiment Station 295

V

Valanne, T. 1022
Valentine, H.T. 301
Van Cleve, K. 686
Van Deusen, J.L. 731
Van Haverbeke, D.F. 702, 703, 704, 707
Van Hooser, D.D. 1093, 1094
VanderHeide, T. 98
Veal, D.L. 115
Venator, C.R. 828
Venkataraman, K. 1318, 1319, 1336, 1339, 1340, 1350
Venketeswaran, S. 560, 561
Verme, L.J. 699
Verry, E.S. 48
Vick, C.B. 1391, 1392, 1393, 1407
Vidakovic, M. 1023
Viereck, L.A. 686
Vogel, W.G. 13, 14

v

von Mosch, J. 1079, 424

V

Voss, E.E. 773

W

Wade, D.D. 679
Wagar, J.A. 256, 296
Wahlgren, H.E. 1271, 1282, 1301, 1302
Waite, T.H. 312
Walker, R.M. 1392, 1393
Walkinshaw, C.H. 478, 488, 510, 511, 560, 561, 565
Walkotten, W.J. 123
Wall, B.R. 1074, 1075, 1076, 1077
Wallmo, O.C. 185, 252
Walters, G.A. 829, 830, 831, 832, 854
Walters, R.S. 598, 935, 941, 942
Walvekar, A.G. 651
Wangaard, F.F. 1351
Ward, A.L. 242, 243
Ward, D.E. 210, 225, 679
Ward, F.D. 206
Ward, F.R. 1088, 638, 662
Ward, J.C. 1239, 1303, 1304
Ward, S.M.F. 1000
Ware, K.D. 933
Wargo, P.M. 562, 563, 564
Waring, R.H. (eds) 750
Washburne, R.F. 271
Waters, M.P. III 806
Waters, M.P., III 692, 694
Watkins, R.K. 241
Watt, R.F. 779
Weber, F.P. 1095, 1096, 1097, 311, 312, 431
Weeks, R.W. 242, 243
Weete, J.D. 510, 511, 561, 565
Welch, B.L. 566
Welch, G.R. 294
Wells, C.G. 780
Welty, J.R. 1219
Wendel, G.W. 732, 833
Wendt, J.S. 255
Wenger, K.F. 757
Wengert, E.M. 1305
Werner, R.A. 383, 384
Westfall, S.E. 175, 205
Whigham, D. 780
Whitaker, L.B. 165
White, E.H. 763

White, G. 813
White, J.D. 385
White, L.R. 297
White, T.H. 1097
White, W.B. 460
Whitmore, J.L. 714, 886, 976
Wick, C.H. 647
Wicker, E.F. 340
Wickman, B.E. 386, 420
Wilcox, W.W. 1306
Wilken, L. 1034, 1035
Wilkes, M. 1091, 430
Wilkins, R.M. 714
Wilkinson, R.C. 977
Wilkinson, T.L. 1394, 1395
Williams, D.T. 648, 697
Williams, G.P., Jr. 124
Williams, K. 139
Williams, L.H. 439, 441, 442, 448
Williams, R.D. 834, 835
Wilson, B.C. 795
Wilson, C.L. 542, 599
Wilson, L.F. 387, 388, 389, 390, 401, 402
Winstead, J.E. 978
Wodzicki, T.J. 1038
Wolfe, F., Jr. 45
Woll, A.M. 1086, 428
Wolter, K. 1252
Wolter, K.E. 1285
Wolters, G.L. 211, 212
Woodfin, R.O. 1259, 1260, 1261, 1307
Woodson, G.E. 1270, 1308, 1351
Works, D.W. 836
Worley, D.P. 915
Wright, H.A. 651
Wright, J.W. 705
Wright, M.G. 1176
Wu, K.Y.T. 1309
Wysor, P.S. 1143

Y

Yamamoto, T. 125, 15
Yandle, D.O. 925
Yang, H. Y. 938
Yates, H.O. 405
Yates, H.O., III 313, 333, 391, 392
Yawney, H.W. 957
Yendol, W.G. 421
Yerkes, V.P. 1277
Yocom, H.A. 713
Yoshiyama, R.M. 459
Young, J.L. 567

Z

Zahn, J.J. 1291, 1292
Zajaczkowski, S. 1038
Zak, B. 568
Zaman, M.B. 569, 708, 709, 711, 712, 837
Zaman, M.B.: Khan, M.A. 710
Zambrana, J.A. 798, 828
Zamski, E. 946, 947
Zasada, J.C. 683, 684, 685, 733
Zasada, Z.A. 739
Zelawska, B. 970
Zelawski, W. 887, 888, 979, 980
Zicherman, J.B. 1310, 1311, 1312
Zinkel, D.F. 1352
Zoerb, M.H., Jr. 397

INDEX

The first number(s) following the index term refers to the narrative statement in the *Research Accomplishments 1973* report. Numbers in parentheses refer to the publication list.

A

Abies **225** (544)
  lasiocarpa **156** (507)
Acacia **223** (569)
acacia koa **216** (509)
Acer saccharum **189, 200, 270, 271** (499, 500, 562-564, 597-599)
acid, chlorogenic **444** (858)
  gibberellic (859, 866, 870, 880, 885)
  indoleacetic **451** (869, 879)
  indolebutyric (859)
acids, fatty **619** (433-435)
adhesion **564** (1245)
adhesives **562** (1235)
advance regeneration (730)
adventitious roots (880)
aerial application **349** (145)
  logging **316** (1207, 1211)
  photographs **26** (157, 158)
  photography **119, 121, 372, 374** (312, 424, 426, 461-463)
  photos **334** (76)
  spraying **16** (298)
  spraying, malathion **255** (401)
Afognak Island **353** (724)
Agathis pumila **249** (398)
air pollution, **19-25, 119, 368** (450, 451, 454-456, 458, 459, 461-463, 619-621, 757)
  oxidant **20-22, 25** (455, 456, 458, 459, 619-621)
  particulate **24** (451)
air pollution chamber **22** (459)
  quality **96, 97** (625-627, 629, 637, 648)
  tanker **98, 99** (656, 661)
airplane, CL-215 **98** (661)
Alabama, forest inventory **280** (1043-1045)
Alaska **102, 146, 226, 353, 356, 357, 369, 375** (323, 324, 659, 660, 680, 681, 683-687, 724, 733)
alder **333** (117)
aldrin **268** (402)
allowable cut effect **300** (1102)
Alnus rubra **263** (601-603)
amelioration **24** (451)
ammonia **19** (450)
analyzing recreation resources and use **75** (700, 701)
anatomy **23, 497** (454, 946, 947)
animal damage **442-444** (855-858)
  feed **597** (1315, 1327, 1335)
  nutrition **43, 45** (215, 231)
anobiids **620** (442)
ant, Texas leaf-cutting **256** (417, 419)
Anthocephalus chinensis **429** (828)
anthracnose **155** (481)
antiattractants **231** (403, 415)
apartment construction **639** (1176)
aphid, balsam woolly **260** (399)
Appalachian hardwoods (853)
Arasan **442** (855)
Araucaria araucana **201** (494)
Arceuthobium **147-149, 228, 230** (501, 502, 520, 538, 575, 582)
  vaginatum **489** (929)
Archips semiferanus **167** (387)
Armillaria mellea **200** (562-564)
artificial regeneration **360, 399-431, 532** (725, 740, 781-837, 994, 1037)
ash, green **351, 412, 440** (720, 805, 852)
  tropical (832)
  white **415** (809)
ash dieback **119** (461-463)
aspen, **165, 301, 329, 362** (110, 746, 747, 1032, 1110-1112)
  quaking (trembling) **314, 357** (365-368, 683-685)
aspen diseases **221** (506)
  ecology **221** (506)
  insects **221** (506)
  management **220** (473)
associations, timber producers **310** (1135)
attractants **134, 135, 191, 208, 231, 234, 246, 256** (334, 351, 376, 383, 384, 396, 403, 410, 415, 417, 419)
Australian toon **587** (829, 830, 1296)
automatic data processing **370** (428-431)
auxins (866)
avalanche **345** (131)

B

Bacillus thuringiensis **252** (421)
bacteria **137, 159** (359, 370)
bacterial infection **590** (1304)
bag papers **607** (1365)
bags **607** (1365)
baldcypress **389** (766, 767)
balloon logging **3, 316** (4, 1207)
balsam woolly aphid **260** (399)
bark **582, 595, 636** (682, 698, 1145, 1284, 1313, 1326)
  beetles **120, 142, 146, 226** (312, 323, 324, 374, 680, 681)
  characteristics **582** (1284)
  chip separation **313** (1194-1196, 1202-1205, 1217, 1218)
  thickness (903, 922)
  utilization **582** (1284)
barking **221** (506)
basal area, stand **481** (915)
basswood, American (1033)
bed material movement **330** (79)
beech **211** (375)
beech bark disease **151** (548)
beetle, bark, elm **183, 237** (348, 393)
  bark, smaller European elm **184, 185, 236, 237** (342, 362, 393, 404)
  Douglas-fir **231** (403, 415)
  fir engraver **185** (362)
  mountain pine **140, 233, 234, 276** (316, 408, 410, 422)
  southern pine **136-138** (321, 359, 360)
  spruce **141, 143, 232** (349, 354, 416)
beetles, bark **120, 142, 146, 226** (312, 323, 324, 374, 680, 681)
  powder-post **631** (448)
benomyl **235, 236, 238** (404, 574, 585, 600)
behavior patterns, small owners **304** (1109)
Betula alleghaniensis **175** (498)
  papyrifera **356** (733)
bibliography, **65, 649** (289, 1189)
  fish conservation **65** (289)
  wildlife conservation **65** (289)
bibliography questionnaires **72** (290)
big game **221** (506)
billberry **75** (700, 701)
bioassays **208, 237** (376, 393)
biochemical methods **450** (867)
biological control **228, 263, 266, 267, 621** (575, 586-588, 601-603, 608)
biological control (insects) **136, 137, 140, 159, 162, 163, 185, 213, 248, 249, 253, 254, 620** (316, 331, 345, 359, 360, 362, 370, 379, 395, 398, 406, 411, 442)
biological evaluations **130** (305)
biology **133, 144, 146, 163, 167, 168, 188, 192, 211, 258, 631** (313, 323, 324, 333, 356, 371, 373, 375, 379, 387, 405, 448)
biology, seed **190** (326)
biomass **366** (755)
birch, **552** (1022)
  Alaska paper **357** (683-685)
  paper **386** (761, 762, 786, 787)
  river **531** (993)
  yellow **391** (771)
black root rot **199** (529)
blight **150, 262** (522, 593)
blister rust **239, 520, 521** (571, 952, 981, 982)
blueberry **75** (700, 701)
board stabilization **570** (1262)
bog **346** (127)
bogs, lake-filled **325** (57)
bond **562** (1235)

Bordeaux fungicide **152** (516, 517, 550)
borer, cottonwood clearwing **210** (315)
mahogany shoot **227** (714)
poplar **210** (315)
red oak **213** (345)
Botrytis **238** (600)
breeding, mutation (1012)
bridge decks, wood **614** (1390)
design **614** (1390)
bridges, wood **614** (1390)
brown spot **152, 153** (497, 516, 517, 550)
brown-spot needle blight **522, 528** (983, 990)
browse management **32** (179, 206)
plants **35** (166)
ranges **40, 41, 43** (190, 202, 215)
brush control **13** (21, 801)
bud activation **88** (623)
growth (859)
budworm, spruce **129** (300)
western **17, 128, 130, 131, 160, 161, 251** (299, 302, 305, 310, 318, 353, 412)
building codes **651** (1178-1181)
burning **14** (19)
butt sweep **272** (581)

C

cable insulating materials **627** (444)
insulation, nontoxic **628** (447)
calibration **372** (426)
California, forest inventory **281** (1052)
cambial growth (1038)
xylem derivatives (1038)
campground management **59, 64** (283, 284)
camping, family **58** (280, 281)
campsite facilities **62** (272)
Canadian forest products **638** (1152)
canker, *Fusarium* **171** (484)
canker development **174** (474)
disease **175** (498)
cankers **221** (506)
canoeing use **63** (292)
carbamates **626** (443)
carbohydrates **600** (1324, 1343-1346)
carpenterworm **208, 209** (376, 377)
Carya **206** (479)
casebearer, larch **249** (398)
casuarina **410** (797)
cattle feed **597** (1315, 1327, 1335)
cedar, Spanish **517** (976)
Cedrela **517** (976)
Cedrus **223** (569)
cellulose **113, 600** (677, 1324, 1343-1346)
fibers **565** (1253)
cemetery ecology **52** (258)
Cenococcum graniforme **267** (588)
census **33, 34** (160, 171)
Ceratocystis **186** (528)
coerulescens **189** (499, 500)
fagacearum **187, 235** (515, 574, 585)
ulmi **235** (574, 585)
Chamaecyparis lawsoniana **277** (606)
chaparral **5** (3)
management **307** (1130)
chemical control **235, 238, 245, 255, 262, 268** (401, 402, 574, 585, 593, 594, 600)
chemical control (insects) **17, 163, 232, 233, 237, 250, 252, 259, 276, 625-627, 631** (299, 379, 393, 397, 407-409, 413, 414, 416, 421, 422, 443, 444, 446, 448)
chemical inhibitors **246** (396)
chemicals **318, 349** (42, 145)
in wood **591** (1309)
chemistry **600** (1324, 1343-1346)
chemotherapy **235** (574, 585)
cherry, black **355, 411** (732, 803)
children, fire prevention lessons **81** (632, 634)
chip storage **629** (618)
chipping, field **312** (1206)
chipping head rigs **309** (1139)
chlorinated hydrocarbons **625** (446)
chlorogenic acid **444** (858)
Christmas tree diseases **125** (468, 469)
trees **152, 649** (516, 517, 550, 1189)
chromatography, gas **571** (1263)
CL-215 airplane **98** (661)
classification **499, 548** (950, 1015)
of forest trees **498-500** (948-951)
cleaning **434** (843)
clearcut **346** (127)
clearcutting **14, 361-363** (19, 744-748)
climate **368, 413** (757, 806)
Clitocybe tabescens **198** (533)
codes, building **651** (1178-1181)
colchicine **552** (1022)
cold-hardiness (728)
Coleosporium crowellii **157** (523)
color films **371** (425, 427)
infrared photography **121, 373** (304, 423)
inheritance (958)
Columnocystis **622** (607, 611, 612)
combustion **113** (677)
intensity **114** (668, 673)
commercial thinning **470** (899)
compaction **3** (4)
competition **362** (725, 746, 747)
composition boards **570** (1262)
compression wood **555** (1221)
computer graphs **104** (658)
model **115, 116, 637** (672, 674, 1169)
models, for wilderness management **54** (261)
programs **71, 103, 559** (294, 654, 657, 930, 931, 1231)
systems **488** (928)
technique **113** (677)
condensation **605** (1353)
cone and seed diseases **258** (405)
cone and seed insects **133, 190, 192, 258, 259** (313, 326, 333, 397, 405)
cone crops (722)
crops, rating **423** (821)
cones **416** (810)
conifers **272** (581)
construction, **562, 615** (1235, 1391-1393)
apartment **639** (1176)
house **605, 613** (1353, 1387)
road **7** (1)
container planting **418** (812)
containers, corrugated **604** (1334)
shipping **604** (1365)
contour trenches **342** (66)
control, biological; *see* biological control
brush **13** (21, 801)
chemical; *see* chemical control
cultural **257, 258** (400, 405)
disease; *see* disease control
hardwood **439** (713, 848-851)
silvicultural; *see* silvicultural control
copper sulphate **262** (593)
coppicing **441** (854)
corrugated containers **604** (1334)
fiberboard **599** (1322)
cost savings, skidding **643** (1170)
cottonwood, **387, 419, 454, 457, 511, 594** (763, 813, 874, 875, 969, 1026, 1027)
black **357** (683-685)
cottonwood clearwing borer **210** (315)
epidemiology **210** (315)
photosynthesis **456** (877)
rooting **536** (1011)
counter, electric eye **69** (277)
creep **612** (1385)
creosote **571** (1263)
Cronartium fusiforme **177, 178, 239, 242-245** (488, 510, 511, 525, 554, 560, 571, 573, 578, 589, 590, 594)
himalayense **181** (476)
ribicola **180, 219, 239-241** (489, 492, 530, 570, 571, 576, 577, 583, 584)
crop-tree release (853)
crown fuels **91** (647)
cruising, presale **484** (920)
timber **569** (1260)
Cryptococcus fagi **151** (548)
cultivar (967)
cultural control **257, 258** (400, 405)
practices **276** (422)
culture, short rotation (826)
Cupressus arizonica **262** (593)
cutting, shelterwood **354** (727)
cutting systems, forest **358** (734, 735)
Cylindrocladium clavatum **201** (494)

D

DDT **11, 12** (16, 26)
Dalbergia **223** (569)
data analysis **371** (425, 427)
processing, automatic **370** (428-431)
dawn redwood **537** (1000)
decay, **132, 205, 206, 271, 621** (460, 470-472, 479, 480, 491, 513, 597, 599, 608)
wood **203, 269, 622-624, 629, 630** (531, 541, 543, 545, 547, 572, 607, 610-612, 614-618)
decay detection **132** (460, 470-472)
prevention **624** (614, 615)
process **203** (531, 541, 543, 545, 547)
decline **216, 218** (509, 555)
deer, **32, 43** (179, 206, 215)
black-tailed **443** (857)
defect **132, 207, 212, 272, 274** (460, 470-472, 475, 546, 581, 604)
defoliation **122** (301)
effects **129** (300)
defoliators **250** (407, 409, 413, 414)
DeHaviland aircraft **99** (656)
Delaware, forest inventory **279** (1039)
demand and price situation **640** (1158)

Dendrosoter protuberans **183** (348)
desert grasslands **328** (120)
design criteria **611** (1384)
detection, disease **119, 126** (461-463, 466)
diapause **160** (353)
Diaporthe **175** (498)
dieback, **200** (562-564)
  ash **119** (461-463)
dimension cuttings **576** (1272)
  stock **559** (1231)
  yield **586** (1293)
Dioryctria abietella **191** (334)
direct seeding **408, 411** (796, 803, 804)
disease, **212, 333** (117, 475)
  beech bark **151** (548)
  canker **175** (498)
  Dutch elm **119, 186, 235, 236, 603** (404, 461-463, 528, 574, 585, 1342, 1352)
  physiogenic **217, 218** (518, 519, 555)
  sapstreak **189** (499, 500)
disease control **152, 173, 236, 261, 275** (404, 516, 517, 549-551, 595, 605)
  detection **119, 126** (461-463, 466)
  impact **119, 220** (461-463, 473)
  physiology **186, 187, 200, 203, 217, 219** (492, 515, 518, 519, 528, 531, 541, 543, 545, 547, 562-564)
  resistance **239** (471)
  survey **119, 223** (461-463, 527, 569)
diseases, **225** (544)
  aspen **221** (506)
  Christmas tree **125** (468, 469)
  cone and seed **258** (405)
  hardwood **155** (481)
  insect **124, 248, 253** (308, 395, 406)
  nursery; *see* nursery diseases
  plantation **198, 201, 244, 267** (494, 533, 588, 590)
  rust **526, 527** (988, 989)
  shade tree **235, 271** (574, 585, 597, 599)
  shrub **224** (341)
  tree; *see* tree diseases
disking **408, 475** (796, 908)
distribution and classification of forest trees **498-500** (948-951)
dog leg **272** (581)
dormancy, **551** (1019)
  seed **400** (782)
Douglas-fir, **348, 407, 418, 428, 442-444, 465, 473, 504, 513, 572, 587** (142, 795, 812, 827, 855, 857, 858, 890, 892, 905, 956, 958, 971, 1264, 1296)
  coast **569** (1260)
  Rocky Mountain **588** (1300)
Douglas-fir beetle **231** (403, 415)
  thinning **299** (1101)
  tussock moth **124, 164, 247** (308, 386, 420)
drainage **325, 352, 389** (57, 723, 766, 767)
drought **217** (518, 519)
Dutch elm disease **119, 186, 235, 236, 603** (404, 461-463, 528, 574, 585, 1342, 1352)
dwarf mistletoe **147-149, 228-230, 489** (501, 502, 520, 538, 575, 582, 596, 929)
  infection **149** (538)
  spread **148** (520)

E

eastern spruce budworm **139** (300)
ecological relationships **364-369** (686, 749-757)
ecology, **169, 258** (332, 405)
  aspen **221** (506)
  cemetery **52** (258)
  plant **28, 38, 39** (170, 210, 211)
  range **27** (162, 163)
  systems (753)
economics, **47** (249)
  timber growing **301-305** (1109-1112, 1115, 1117, 1122)
El Dorado pine (967)
Elaphomyces **196** (559)
elm **603** (1342)
  bark beetle **183, 237** (348, 393)
  spanworm **169** (332)
end and edge gluing **559** (1231)
Endogone **127, 196** (464, 558)
energy balance **346** (127)
  exchange **346** (127)
environment **80** (708-712)
environmental influences **134, 136, 139, 169, 197, 618** (332, 352, 360, 383-385, 437, 439)
enzymes **450, 533, 597** (867, 995, 1315, 1327, 1335)
epicormic sprouting **435** (844)
epidemiology, **143, 166, 167, 247** (354, 358, 387, 420)
  cottonwood **210** (315)
Epilobium **156** (507)
equipment **374** (424)
erodibility indices **6** (15)
erosion, **3, 5, 6, 9, 14, 15, 344** (3, 4, 7, 15, 18, 19, 137)
  logging **4** (9)
erosion control **324** (47)
ethylene dibromide **232** (416)
Eucalyptus **201, 212, 225, 410** (475, 494, 544, 797)
  robusta **587** (1296)
eucalyptus, saligna (830, 831)
Eucosma cocana **192** (333)
European pine shoot moth **170, 246** (330, 396)
evaluating and improving wildlife habitat **36** (699)
evaluation methods **128** (302)
evaporation, weather control **321** (41)
evaporation supression **318** (42)
evapotranspiration **320, 346** (38, 127)
exposure times **372** (426)
extractives **603** (1342, 1352)

F

Fagus grandifolia **151** (548)
family camping **58** (280, 281)
fatty acids **619** (433-435)
feed, animal **597** (1315, 1327, 1335)
  cattle **597** (1315, 1327, 1335)
fertilization, **1, 37, 195, 383, 384, 386, 389, 425, 458** (5, 194, 512, 758, 759, 761, 762, 766, 767, 823, 878)
  nitrogen **425** (823)
fertilizer (770)
fiberboard, corrugated **599** (1322)
fibers, **595** (1313, 1326)
  cellulose **565** (1253)
fiberwood prices **634** (1142)
field chipping **312** (1206)
financial evaluation and planning **297-300** (1098, 1100-1102)
  maturity **302** (1115)
fine wood residues **633** (1140)
fir, balsam **129** (300, 726)
  noble **418** (812)
  red **319** (36)
  silver **399** (781)
  white **164** (386)
fir engraver beetle **185** (362)
fire, **4** (9)
  effect in wilderness **56** (259, 266)
fire behavior **114, 118** (668, 669, 671, 673, 675, 679)
  danger indexes **103** (654, 657)
  danger rating **105** (653)
  effects **205, 206** (479, 480, 491, 513)
  environment **117** (667)
  fuel relations **85** (641)
  hazard reduction **87** (642)
  igniter **95** (639)
  management **94** (628, 636, 640, 646)
  planning **84** (622, 624)
  prevention **81, 83, 86** (631, 632, 634, 635, 644, 713)
  protection **93** (630)
  retardants **100, 101, 111, 112** (655, 662, 670, 676)
  spread **115, 116** (672, 674)
  weather **118** (669, 671, 675, 679)
  weather data processing **103** (654, 659)
  weather observer's handbook **106** (652)
  whirls **117** (666, 667)
  zones, urban **651** (1178-1181)
fires, man-caused **82** (633)
firs, true **499** (950)
fish conservation bibliography **65** (289)
  habitat **13** (21)
fisherman preferences **66** (287)
fishing management **66** (287)
flame suppressants **378** (693)
flammability **109, 111** (663, 664, 670, 678)
flooding **352** (723)
floods **335, 339** (72, 93)
flow, peak **326** (125)
flowering **503, 537, 539, 546** (955, 1000, 1002, 1013)
fluoride **19** (450)
foliage analysis **383, 395** (758, 776)
Fomes annosus **119, 264, 264, 265** (461-463, 579, 592)
food reserves **460** (883)
forage plants **35, 37-39** (166, 194, 210, 211)
  production **42** (222)
forest appraisal **569** (1260)
  cutting **335** (72)
  cutting systems **358** (734, 735)
  defoliators **250** (407, 409, 413, 414)
  fire prevention programs **83** (631, 644)
  fire protection **93** (630)

fires, man-caused **82** (633)
floor **11, 12, 349** (16, 26, 145)
fuels **84, 91** (622, 624, 647)
industry, secondary **641** (1156)
industry marketing report **645** (1153, 1154, 1159, 1160)
inventory **278-382, 482** (717, 916, 1039, 1042-1045, 1049, 1050, 1052, 1053)
inventory, data retrieval **282** (1042)
management **333** (117)
measurements **276-487** (909-925)
products **554, 579** (1220, 1279)
products, Canadian **638** (1152)
ranges **30, 45** (169, 231)
seed production **356** (733)
soil **11, 341** (16, 77)
survey **481** (915)
wildlife relations **31, 32** (179, 183, 206)
forests **339** (93)
Formosan termite **616, 617** (436, 438)
freezing resistance **217** (518, 519)
fuel arrangement **85** (641)
break **86** (635)
break management **88, 89** (623, 643)
consumption **114** (668, 673)
inventories **92** (645)
moisture **105, 107** (651, 653)
types **91** (647)
fuels, crown **91** (647)
fungi **136, 622** (360, 607, 611, 612)
fungicides, **155, 235, 245, 262** (481, 574, 585, 593, 594)
Bordeaux **152** (516, 517, 550)
fungus antagonism **621** (608)
variation **153, 178** (497, 525, 554)
Fusarium canker **171** (484)
fusiform rust **177, 178, 524** (488, 510, 511, 525, 554, 560, 716, 985)

G

game, big **221** (506)
gas chromatography **571** (1263)
genetic markers **533** (995)
selection **513** (971)
terms **548** (1015)
genetics, (1028)
shrub **40** (202)
tree **532** (994)
geographic variation **135, 501, 502, 504** (351, 953, 954, 956)
geometry, hydraulic **330** (79)
germination **352, 415** (723, 809, 836)
gibberellic acid (859, 866, 870, 880, 885)
glossary, **548** (1015)
geographic information systems **296** (718)
glucose metabolism **161** (318)
gluing, **564** (1245)
end and edge **559** (1231)
Goes pulverulentus **211** (375)
grafting **534, 538** (996, 1001)
grafts, root (822)
grain, spiral (1025)
wavy **508** (965)
Grand Canyon **230** (582)
grasslands, desert **328** (120)
gray mold **238** (600)
grazing **28, 44** (170, 218, 234)
greenhouses **420** (814)
growing stock levels (890)
growth, **368, 519,** (757,978)
bud (859)
cambial (1038)
radial **466** (893)
root **475** (879, 908)
seedling **452** (871)
shoot (879)
tree **122, 129** (300, 301)
tree diameter **480** (109)
growth and yield **463-475, 483, 487,** (860, 889-908, 918, 919, 924)
effects **147** (501, 502)
requirements **445-462** (859-888)
response **470** (899)
variation **514** (972)
grubs, white **268** (402)
gummosis **212** (475)
gypsy moth **122, 123, 158, 159, 200, 252, 254** (301, 306, 307, 357, 370, 411, 421, 562-564)

H

habitats **32, 46,** (179, 206, 246)
classification **452** (871)
handcrafts **648** (1187)
hardboard uses **611** (1384)
hardwood **340** (113)
brush **433** (842)
control **439** (713, 849-851)
diseases **155** (481)
insects **122, 165, 167** (301, 322, 387)
lumber **650** (1185)
planting **360** (740)
plywood stock panels **647** (1167)
sawing system **635** (1143)
hardwoods, **398, 541, 583, 602** (780, 817, 1006, 1286, 1329-1331)
Appalachian (853)
northern **36** (699)
southern **208, 209, 211** (375-377)
tropical **227, 409, 410, 429, 517** (714, 797, 800, 828, 976)
harvesting methods **359** (736-739, 742)
hauling distances, pulpwood **287** (1059)
Hawaiian Islands **6, 215, 216** (15, 363, 509)
hazard control **274** (604)
reduction **274** (604, 713)
headrigs, chipping **309** (1139)
heart rot **203, 205, 206, 220, 270, 271** (473, 479, 480, 491, 513, 531, 541, 543, 545, 547, 597-599)
Hebeloma **448** (865)
height growth (870)
helicopters **102** (659, 660)
hemicellulose **112** (676)
herbicides **13, 86, 89, 360, 433, 439** (21, 635, 643, 740, 832, 848-851)
heritability (952)
Hibiscus elatus **410** (797)
high walls **337** (101)
honeycomb **590** (1304)
host specificity **162** (331)
hosts **215** (363)
house construction **605, 613** (1353, 1387)
maintenance **605** (1353)
houses, low-cost **613** (1387)
housing **615** (1391-1393)
demand **637** (1169)
surveys **613** (1387)
huckleberries **75** (700,701)
hybrid mahogany **409** (800)
poplar **171, 455, 547** (484, 833, 876, 1014)
Hydnochaete **611** (607, 611, 612)
hydraulic conductivity **325** (57)
geometry **330** (79)
seeder **101** (662)
hydrocarbons, chlorinated **625** (446)
hydrogen chloride **19** (450)
hydrologic behavior **324** (47)
properties **341** (77)
Hypoxylon **220, 314** (365-368, 473)
mammatum **174** (474)

I

identification **124, 145, 146** (308,323, 324, 391, 392)
ignitability **109** (663, 664, 678)
impact tests **607** (1365)
imports **638** (1152)
India **222** (477)
indices **6** (15)
indoleacetic acid **451** (869, 879)
indolebutyric acid (859)
infection, bacterial **590** (1304)
infestation assessment **120** (312)
infiltration **5, 341** (3, 77)
infrared **293-295** (1078, 1085, 1091, 1096, 1097)
inherent variation **501-519** (901, 944, 952-980)
inheritance **535** (997)
inhibitors, chemical **246** (396)
insect behavior **133, 134, 144, 146, 158, 167, 168, 188, 209, 256** (313, 323, 324, 356, 357, 371, 373, 377, 383, 384, 387, 417, 419)
biochemistry **161, 619** (318, 433-435)
control **16, 17, 129, 130, 135, 161, 190, 231, 234, 237, 246, 247, 252, 257, 268, 276, 628** (298-300, 305, 318, 326, 351, 393, 396, 400, 402, 403, 410, 415, 420-422, 447)
development **617** (438)
disease resistance **520-528** (981-990)
diseases **124, 248, 253** (308, 395, 406)
dispersal **123** (306,307)
growth **251** (412)
impacts **122, 165** (301, 322)
morphology **191** (334)
nutrition **184** (342)
parasites **131, 162, 183, 249, 254** (310, 331, 348, 398, 411)
populations **620** (442)
predators **140, 158, 260** (316, 357, 399)
rearing **254** (411)
survival **251** (412)
insecticide synergists **17** (299)
insects, **225** (544)
aspen **221** (506)

cone and seed **133, 190, 192, 258, 259,** (313, 326, 333, 397, 405)
hardwood **122, 165, 167** (301, 322, 387)
mechanical control (715)
shelterbelt **168** (356)
shoot- and tip-feeding **227** (714)
shrub **224** (341)
silvicultural control (715)
insulation **605** (1353)
insurance companies **305** (1122)
intensive culture **475** (908)
intensive forest culture **225** (544)
interception **340** (113)
interflow **9** (7)
inventory, forest **278-283, 482** (717, 916, 1039, 1042-1045, 1049, 1050, 1052, 1053)
inventorying landscapes **48** (257)
Ips calligraphus **145** (391, 392)
cribricollis **145** (391, 392)
grandicollis **134** (383, 384)
pini **135, 144** (351, 373)
irradiation, pollen **531, 553** (993, 1026)

J

jet stream effect **381** (697)
Juglans **155** (481)
Juniperus **150, 223** (522, 527, 569)
juvenile hormones **251** (412)

K

kadam **410, 429** (797, 828)
kiln drying **567, 572, 650** (1185, 1257, 1264)

L

lake-filled bogs **325** (57)
laminated lumber **609** (1381)
products **585** (1291)
land classification **373** (423)
use classification **370** (428-431)
use planning **84** (622, 624)
landscape resource analysis **48, 49, 51** (253, 256, 257)
resource management **48, 49** (256, 257)
resources **48, 49, 51** (253, 256, 257)
landslides **2, 344** (11, 137)
larch, Japanese **78** (705)
western **249** (398)
larch casebearer **249** (398)
Larcobius erichsonii **260** (399)
leaf retention **454** (878)
legume **333** (117)
Lenzites trabea **619** (433-435)
life tables **130** (305)
light traps **133** (313)
lightning **375-377** (687, 691, 694)
lignin **597** (1315, 1327, 1335)
limb rust **179** (524)
line intersect method **90** (638)
linear expansion **570** (1262)
programming **306** (1126)
literature retrieval **219** (492, 530)
litter control **67** (269-271)
livestock management **45** (231)
loan security **305** (1122)
lobelia cultivation **632** (1034, 1035)
harvest **632** (1034, 1035)
lodgepole needle miner **163, 166** (358, 379)
log centers of gravity **314** (1212, 1216)
grade(s) **556, 563, 569** (1222, 1243, 1260)
grading **583** (1286)
storage **557** (1225)
weights **314** (1212, 1216)
yield **568** (1259)
logging, aerial **316** (1207, 1211)
balloon **3, 316** (4, 1207)
mechanized **359** (736-739)
roadless **316** (1207, 1211)
skyline **316** (1211)
logging erosion **4** (9)
plans **315** (1200, 1201, 1213)
residue(s) **90, 290, 312, 313, 561** (638, 1070, 1073, 1194-1196, 1202-1205, 1217, 1218, 1233)
roads **315** (1200, 1201, 1213)
system **3** (4)
systems, advanced **316** (1207, 1211)
systems, comparing **317** (1198, 1199, 1208, 1209)
Lophodermella morbida **154** (556)
low-cost houses **613** (1387)
lumber, **560** (1232)
hardwood **650** (1185)
laminated **609** (1381)
lumber finishing **579** (1279)
grading **608** (1371)

M

Macrophomina **126** (466)
phaseoli **199** (529)
Magdalis gentilis **257** (400)
mahoe **410** (797)
mahogany, hybrid **409** (800)
mahogany shoot borer **227** (714)
malathion aerial spraying **255** (401)
management, **335** (72)
aspen **220** (473)
browse **32** (179, 206)
campground **59, 64** (283, 284)
chaparral **307** (1130)
fire **94** (628, 636, 640, 646)
forest **333** (117)
landscape resource **48, 49** (256, 257)
range **34, 42, 44, 47, 306-308, 331** (50, 160, 218, 222, 234, 249, 1126, 1130, 1132)
recreation **68** (278)
river **63** (292)
watershed **343** (107)
wilderness **53-55** (260, 261, 265)
wildland **347** (133)
wildlife **33, 46, 47** (171, 246, 249)
management planning **488-489** (926-933)
systems **347** (133)
maple, bigleaf **494** (937, 938)
sugar **302, 490-493, 495, 496, 505** (934, 935, 939-942, 957, 961, 1115)
maple sap **270** (598)
syrup production **490-496** (934-942)
syrup products **646** (1173, 1174)
marketing **633-636** (1140-1145)
markets, timber product **640** (1158)
mass wasting **4, 344** (9, 137)
mathematical model(s) **322, 607,** (46, 1365)
measurements, timber **487** (924)
mechanical properties **606, 612** (1355, 1356, 1385)
mechanized logging **359** (736-739)
medicinal plants **80** (708-712, 837)
Melampsora pinitorqua **239** (571)
meristem, shoot apical **507** (963, 964)
mesosystem weather **380** (689)
mesquite **328** (120)
metabolism, glucose **161** (318)
metrication **482** (916)
Metrosideros collina **216** (509)
Mexico **179** (524)
microorganisms **138** (321)
mineral weathering **348** (142)
mining, surface **15, 336-338** (14, 18, 101, 102)
mistletoe, dwarf **147-149, 228-230, 489** (501, 502, 520, 538, 575, 582, 596, 929)
mobile homes **642** (1162, 1164)
model, forest economy **311** (1138)
mathematical **322, 607** (46, 1365)
projection **297** (1098)
modeling **123, 141, 317, 347** (133, 306, 307, 349, 1198, 1199, 1208, 1209)
moisture content, **109** (663)
seedling (728)
wood **557, 572, 574** (1225, 1264, 1270)
moisture movement **612** (1385)
relations **570** (1262)
mold, gray **238** (600)
Monodontomerus dentipes **162** (331)
monoterpenes **518** (977)
morphological characteristics **578** (1275)
morphology, **601** (1325, 1348, 1349, 1351)
insect **191** (334)
mortality, tree **122, 129, 215** (300, 301, 363)
Morus **223** (569)
moth, pine shoot **170, 246** (330, 396)
tussock **124, 164, 247** (308, 386, 420)
mountain pine beetle **140, 233, 234, 275** (316, 408, 410, 422)
mountain streams **330** (79)
multiple use **332** (97)
multiple-use economics **306-308** (1125, 1130, 1132)
multispectral scanning **121** (304)
mutation breeding (1012)
mycangium **138** (321)
mycoplasma **182** (542)
mycorrhizae **127, 195, 196, 266, 267, 399, 426, 448, 459** (464, 512, 552, 558, 559, 586-588, 781, 824, 865, 882)
mycorrhizae regeneration **449** (864)

N

National Fire -Danger Rating System **115** (674)

Natural Areas, Research **364** (751, 1030)
natural environments **364** (751)
regeneration **350-357, 362, 421, 522** (683-685, 719-733, 746, 747, 815, 816, 983, 1037)
scientific preserves **364** (751)
nature scenery preference **50, 51** (253, 255)
naval stores **497** (943-947)
Nectria coccinea **151** (548)
needle blight, brown spot **522, 528** (983, 990)
cast **153, 154** (497, 556)
miner, lodgepole **163, 166** (358, 379)
nematodes **193, 194** (535-537)
net radiation **346** (127)
Nicaragua **145** (391, 392)
nitrate reductase **195** (512)
nitrogen **333** (117)
fertilization **425** (823)
fixation **263** (601-603)
oxides **19** (450)
noise abatement **76** (702, 703)
noncommercial thinning (840)
nondestructive testing **608** (1371)
nongame **31** (183)
nontoxic cable insulation **628** (447)
North Carolina **260** (399)
northern hardwoods **36** (699)
nurseries **420** (814)
nursery diseases **126, 150, 176, 194, 199, 238, 245, 261, 262, 266** (466, 521, 522, 529, 535, 586, 587, 593-595, 600)
management **426** (824)
methods (829)
practices **424** (788, 799, 820)
production (817)
nutrient cycling **348, 398** (142, 780)
movements **457** (874)
nutrients **14, 328** (19, 120)
nutrition, **616-619** (433-439)
animal **43, 45** (215, 231)
insect **184** (342)
Nyssa aquatica **389** (766, 767)
silvatica var. *biflora* **352** (723)

O

oak, California black **396** (778)
northern red **417** (811)
red **211, 384** (375, 759)
oak chaparral **116** (672)
wilt **119, 187, 188, 235** (371, 461-463, 515, 574, 585)
oaks, **367, 549** (730, 756, 1020)
red **167** (387)
upland **468** (764, 897)
Oberea schaumii **214** (365-368)
oleoresin **506** (959)
organic matter 8 (6)
soils **325** (57)
organophosphates **626** (443)
owners, woodland **304** (1109)
oxidant air pollution **20-22, 25** (455, 456, 458, 459, 619-621)
ozone **19-23, 25** (450, 454-456, 458, 459, 619-621)

P

paint finishes **587** (1296)
Pakistan **223** (527, 569)
paper **601, 602** (1325, 1329-1331, 1348, 1349, 1351)
testing **598, 599** (1317, 1322, 1328)
paraformaldehyde **270** (598)
parasites, insect **131, 183, 249, 254** (310, 331, 348, 398, 411)
particle size 8 (6)
particle size (snow) **322** (46)
particleboard **570** (1262)
particulate air pollution **24** (451)
patents **567** (1257)
pathogenicity **136, 137** (359, 360)
peak flow **326** (125)
peatland **325** (57)
percolation **320** (38)
performance evaluation **607** (1365)
Peridermium filamentosum **179** (524)
harknessii **176** (521)
pini **239** (571)
permeability **341, 584** (77, 1288)
peroxidase **204** (504)
pesticide related research **18, 237** (297, 393)
screening **250** (407, 409, 413, 414)
pesticides, **12, 438** (26, 830, 847)
systemic **18** (297)
Phaeocollybia **196** (552)
Phellinus tomentella **622** (607, 611, 612)
torulosus **202** (485)
phenolic resin **564** (1245)
phenolics **603** (1342, 1352)
pheromones, trail-marking **256** (417, 419)
Phomopsis **150, 175** (498, 522)
phorate **259** (397)
phosphorus, **458** (878)
soil **385** (760)
photochemical smog **20, 25** (456, 619-621)
photography, **292-295** (1078, 1085, 1090, 1091, 1095-1097)
aerial **119, 121, 372, 374** (312, 424, 426, 461-463)
color **121, 373** (304, 423)
infrared **121, 373** (304, 423)
photoscale **120** (312)
photosynthesis, **456** (877, 887)
cottonwood **456** (877)
physiogenic disease **217, 218** (518, 519, 555)
physiology, **23** (454)
disease; *see* disease physiology
Phytophthora cinnamomi **198, 267** (533, 588)
lateralis **277**(606)
phytotoxin **187** (515)
Picea **225** (544)
engelmannii **421, 460** (815, 816, 883)
glauca **420** (814)
mariana **346** (127)
pine, **340** (115)
Aleppo **497** (946, 947)
Austrian **553** (1023)
Caribbean **145** (391, 392)
Choctawhatchee (860)
El Dorado (967)
Honduras **410** (797-799, 885)
jack **408, 420, 461, 509** (796, 814, 884, 966)
Jeffrey **472** (904)
loblolly **139, 197, 324, 382, 398, 400, 401, 406, 413, 414, 426, 427, 466, 506, 510** (47, 352, 385, 692, 721, 780, 782, 783, 793, 802, 806, 807, 824, 825, 893, 901, 959, 968)
lodgepole **163, 166, 234, 257, 276, 319, 358, 445, 446, 480,** (36, 109, 358, 379, 400, 410, 422, 734, 735, 861, 862, 896)
longleaf **522, 528** (983, 990)
ponderosa **394, 432, 436, 445, 446, 464, 472, 485, 489, 572, 579, 593** (775, 777, 838, 845, 861, 862, 891, 904, 906, 921, 929, 1024, 1264, 1279)
ponderosa, Black Hills (731)
red **359, 476** (736-739, 909, 910)
sand **385, 406** (760, 793)
Scotch (Scots) **508, 512, 523, 553** (887, 888, 962, 965, 970, 979, 980, 984, 1023)
shortleaf **390, 403, 404, 506** (768, 769, 790, 791, 959)
slash **406, 425, 426, 458, 469, 475, 477, 478, 515, 524, 535** (793, 822-824, 829, 878, 898, 902, 908, 911, 912, 973, 974, 985, 997, 1018)
spruce **578** (1275)
sugar **563** (1243)
Virginia **427, 503** (825, 955)
white **500, 521, 525, 526, 556, 603** (870, 951, 981, 986-988, 1222, 1352)
white, western **434, 470, 520, 551** (843, 899, 952, 982, 1019)
pine shoot moth **170, 246** (330,396)
pines, **527, 540** (989, 1003)
southern **192, 259, 515, 557, 566, 577, 582, 589** (333, 397, 974, 1225, 1254, 1273, 1284, 1302)
western **142** (374)
Pinus **176, 179, 201, 223, 225, 239, 242, 243** (494, 521, 524, 544, 569, 571, 573, 588, 589)
attenuata **154** (556)
ayacahuite **157** (523)
banksiana **420** (814)
canariensis **181** (476)
caribaea **555** (1221)
clausa **193, 197** (533, 536, 537)
contorta **358, 445** (734, 735, 861)
coulteri **23, 172** (454, 553)
elliottii **177, 244, 264, 267** (488, 510, 511, 560, 588, 590, 592)
jeffreyi **21, 23** (454, 455, 458)
lambertiana **23** (454)
merkusii **462** (886)
monticola **240, 241, 273** (570, 576, 577, 580, 583, 584)
palustris **153, 193** (497, 536, 537)
ponderosa **21, 23, 147, 154, 218, 230, 446, 464, 485, 489, 579** (454, 455, 458, 501, 502, 555, 556, 582, 861, 891, 921, 929, 1279)
resinosa **152, 173** (516, 517, 549-551)
sabiniana **149** (538)

strobiformis **202** (485)
sylvestris **125, 152, 153** (468, 469, 497, 516, 517, 550)
taeda **245, 264, 265** (579, 592, 594)
pinyon-juniper **554** (1220)
Pisolithus tinctorius **267** (588)
planning **347** (133)
plant biochemistry **139** (385)
control **27, 30** (162, 163, 169)
development **35, 37-39** (166, 194, 210, 211)
ecology **28, 38, 39** (170, 210, 211)
hormones **539, 543** (1002, 1007, 1008)
insect relations **190** (326)
plantation diseases **198, 201, 244, 267** (494, 533, 588, 590)
plantations **477, 555** (912, 1221)
planting, **421** (815, 816)
container **418** (812)
hardwood **360** (740)
replacement (819)
planting black walnut (772)
failure **198** (533)
plantings, protection **76-79** (702-707)
plants, browse **35** (166)
forage **35, 37-39** (166, 194, 210, 211)
medicinal **80** (708-712, 837)
Platanus **155** (481)
plywood, southern pine **564** (1245)
plywood stock panels **647** (1167)
pole blight **273** (580)
pollen **402** (789)
irradiation **531, 553** (993, 1023)
pollination **413** (806)
pollution;*see also* air pollution **596** (1314, 1332, 1333, 1337, 1338)
polyembryony **505** (957)
polyploidy **552** (1022)
polyvinyl chloride **628** (447)
poplar, **539** (1002)
balsam **357** (683-685)
hybrid **171, 455, 547** (484, 833, 876, 1014)
poplar borer **210** (315)
population dynamics **143, 158, 159, 165, 166, 170, 210, 214, 231, 247, 544** (315, 322, 330, 354, 357, 358, 365-368, 370, 403, 415, 420, 1009)
Populus **171, 225** (484, 544)
tremuloides **174, 220, 221** (473, 474, 506)
Poria weirii **119, 204, 253, 273** (461-463, 504, 580, 601-603)
powder-post beetles **631** (448)
preattack guide **93** (630)
precommercial thinning **414** (807)
predators, **185** (362)
insect **140, 158, 260** (316, 357, 399)
vertebrate **141,213** (345,349)
prediction **122, 123, 128, 141** (301, 302, 305, 307, 349)
prefreezing **558** (1227)
preplanning timber sales **315** (1200, 1201, 1213)
presale cruising **484** (920)
prescribed burning **94, 173** (549, 551, 628, 636, 640, 646, 713)
fire **95, 97** (625-627, 629, 637, 639)
preservatives, wood **630** (616, 617)
Press-Lam **609** (1381)
prices, pulpwood **286** (1063, 1072)
primary plant residues **288** (1062, 1065, 1067, 1068, 1069)
probability distributions **379** (688, 690, 695, 696)
procurement strategies **644** (1148)
production function **332** (97)
progeny testing **535** (997)
programming, **299, 559,** (1101, 1231)
linear **306** (1126)
projection model **297** (1098)
properties, mechanical **606, 612** (1355, 1356, 1385)
protection plantings **76-79** (702-707)
protozoa **616-618** (436-439)
provenance trials **462, 517** (886, 976)
provenience (962)
pruning (841)
Pseudopityophthorus minutissimus **188** (371)
pruinosis **188** (371)
Pseudotrichonympha grassi **616** (436)
Pseudotsuga **225** (544)
menziesii **195, 263, 273, 465, 587** (512, 580, 601-603, 892, 1296)
psocids **620** (442)
publications list (1028)
Pucciniastrum **156** (507)
pulp **600, 602** (1324, 1329-1331, 1343-1346)
chips **629** (618)
properties **601** (1325, 1348, 1349, 1351)
pulping **595, 600** (1313, 1324, 1326, 1343-1346)
pulpwood **602** (1329-1331)
hauling distances **287** (1059)
prices **286** (1063, 1072)
production **285** (1058, 1064)
purification techniques **248** (395)
Pyemotes scolyti **185** (362)

Q

Quercus **155, 187, 205, 235, 242** (480, 481, 491, 513, 515, 573, 574, 585, 589)
questionnaires, bibliography **72** (290)

R

radial growth **466** (893)
radiation, net **346** (127)
radiation injury **461** (884)
shield **108** (650)
radiography **131, 132** (310, 460)
rainfall, simulated **6** (15)
range ecology **27** (162, 163)
improvements **29** (152)
management **34, 42, 44, 47, 306-308, 331** (50, 160, 218, 222, 234, 249, 1126, 1130, 1132)
surveys **26** (157, 158)
ranges, browse **40, 41, 43** (190, 202, 215)
rating cone crops **423** (821)
ray shake **207** (546)
rays, vascular **447** (863, 872)
rearing **160, 183, 184** (342, 348, 353)
reclamation **171, 336** (14, 484)
recovery ratios **560** (1232)
recreation, in cemeteries **52** (258)
recreation area vegetation **61** (286)
facilities **62** (272)
management **68** (278)
planning **73, 74** (268, 285)
policy **74** (285)
site maintenance **274** (604)
recreational demands **73, 74** (268, 285)
use **69-71** (276, 277, 294)
recycling **596, 604** (1314, 1332-1334, 1337, 1338)
red-headed pine sawfly **255** (401)
red oak borer **213** (345)
redwood, **587** (1296)
dawn **537** (1000)
reforestation **406** (793)
regeneration, **443, 452, 459** (742, 857, 871, 882)
advance (730)
artificial **360, 399-431, 532** (725, 740, 781-837, 994, 1037)
natural **350-357, 362, 421, 522** (683-685, 719-733, 746, 747, 815, 816, 983, 1037)
remote sensing **26, 119-121, 370-374** (157, 158, 304, 312, 423-431, 461-463)
replacement planting (819)
research, pesticide-related **18, 237** (297, 393)
Research Natural Areas **364** (751, 1030)
research needs (926)
reservoirs **8** (6)
residue, logging **90, 290, 312, 313, 561** (638, 1070, 1073, 1194-1196, 1202-1205, 1217, 1218, 1233)
residues, **269, 288, 290, 623, 636** (572, 610, 1062, 1065, 1067, 1068-1070, 1073, 1145)
fine wood **633** (1140)
primary plant **288** (1062, 1065, 1067, 1068, 1069)
resin, phenolic **564** (1245)
xylem **142** (374)
resin ducts **497** (946, 947)
resistance, **239, 265** (571, 579)
disease **239** (571)
resistance mechanism **240, 243** (576-578, 584)
screening **241, 242** (570, 573, 583, 589)
resources, landscape **48, 49, 51** (253, 256, 257)
visual **48, 49** (256, 257)
respiration (887)
retardants **112** (676)
Reticulitermes flavipes **619** (433-435)
revegetation **1, 10** (5,8)
rhododendron **388** (765)
Ribes **180** (489)
ring failure **590** (1304)
shake **207, 591** (546, 1309)
rings, tree **483** (917-919)
river management **63** (292)
road construction **7** (1)
roadbuilding, accelerated **303** (1117)
roadless logging **316** (1207, 1211)
roads, **9, 10** (7, 8)

logging **315** (1200, 1201, 1213)
Robinia **182** (542)
robusta eucalyptus **587** (1296)
root development **197** (352, 866)
effects **147** (501, 502)
grafts (822)
growth **475** (879, 908)
rot **119, 126, 198, 200-202, 216, 264-266** (461-463, 466, 485, 494, 509, 533, 562-564, 579, 586, 587, 592)
rot, black **199** (529)
suckers **329** (110)
rooting, **542, 543, 545** (1005-1008, 1010)
cottonwood **536** (1011)
rooting cuttings (802)
hormones **545** (1010)
roots, **366** (755)
adventitious (880)
rootwood **577** (1273)
rot, heart **203, 205, 206, 220, 270, 271** (473, 479, 480, 491, 513, 531, 541, 543, 545, 547, 597-599)
root **119, 126, 198, 200-202, 216, 264-266** (461-463, 466, 485, 494, 509, 533, 562-564, 579, 586, 587, 592)
white **204, 623** (504, 610)
runoff **9, 14, 331, 339** (7, 19, 50, 93)
rust, **156, 157, 176, 177, 179-181, 239-244** (476, 488, 489, 507, 510, 511, 521, 523, 524, 560, 570, 571, 573, 576-578, 583, 584, 589, 590)
fusiform **177, 178, 524** (488, 510, 511, 525, 554, 560, 716, 985)
limb **179** (524)
rust diseases **526, 527** (988, 989)
inoculation **241** (570, 583)
resistance **178, 510** (525, 554, 968)

S

sage, creeping **87** (642)
saligna eucalyptus (830, 831)
sampling, **90** (638, 788)
stratified (925)
3P **291** (1093, 1094)
trail use **69** (277)
visitor use **70, 71** (276, 294)
sandhills **406** (793)
Saperda inornata **214** (365-368)
sapstreak disease **189** (499, 500)
sawflies **162** (331)
sawfly, red-headed pine **255** (401)
sawing **576** (1272)
system, hardwood **635** (1143)
yields **576** (1272)
sawmill equipment **560** (1232)
saw-palmetto **30** (169)
scanning electron microscopy **565** (1253)
Scirrhia acicola **152, 153** (497, 516, 517, 550)
Sclerroderris canker **173** (549, 551)
scrub oak regrowth **89** (643)
seasonal changes **139** (385)
secondary forest industry **641** (1156)
secretion **497** (946, 947)
sediment **10, 348** (8, 142)
deposition **8** (6)
yield **324** (47)
seed **350-352, 415** (719, 720, 723, 809)
and cone handling **424** (820)
biology **190** (326)
coatings **442** (855)
collection, tree **423** (821)
dormancy **400** (782)
extraction **429** (828)
orchards **402, 403, 529, 534, 546** (789, 790, 991, 996, 1013)
production **382, 404, 413, 425** (692, 721, 791, 806, 823)
ripeness (731)
set **530** (992)
storage **401** (782, 784)
tree **363** (748)
viability **355** (732)
yield **509** (966)
zones **423** (821)
seeder, hydraulic **101** (662)
seeding, direct **408, 411** (796, 803, 804)
seedling growth **452** (871)
moisture content (728)
storage **238** (600)
survival **420, 452** (814, 871)
selection **363** (748)
gain (952)
self pollination **530** (992)
Sequoia gigantea **238** (600)
sempervirens **587** (1296)
shade tree diseases **235, 271** (574, 585, 597, 599)
trees **275** (605)
shake, **207** (546)
ray **207** (546)
ring **207, 591** (546, 1309)
shelterbelt insects **168** (356)
shelterbelts **76-79** (702-707)
shelterwood **363** (748)
cutting **354** (727)
shipping containers **607** (1365)
shoot- and tip-feeding insects **227** (714)
shoot apical meristem **507** (963, 964)
growth (879)
short rotation culture (826)
shrub diseases **224** (341)
genetics **40** (202)
insects **224** (341)
regrowth **88** (623)
silvicultural control **148, 175, 222, 229, 230, 244, 264, 272, 273, 277** (477, 498, 520, 580-582, 590, 592, 596, 606)
methods **358-363** (734-748)
treatments **357** (683-685)
simulated rainfall **6** (15)
simulation **317, 323, 489** (33, 929, 933, 1198, 1199, 1208, 1209)
Sirococcus **172** (553)
site evaluation and soil improvement **383-398** (758-780)
index **390, 394, 476** (764, 768, 769, 775, 777, 909, 910)
preparation **244, 406** (590, 785-787, 793, 794, 801)
protection **338** (102)
quality **385, 397** (760, 779)
Sitka spruce-hemlock type **354** (727)
ski area analysis **60** (279)
area development **60** (279)
areas **1** (5)
skidder operator **643** (1170)
skidding cost savings **643** (1170)
skyline logging **316** (1211)
SLAR **121** (304)
slash fuel loading **92** (645)
slope stability **2, 7** (1, 11)
smaller European elm bark beetle **184, 185, 236, 237** (342, 362, 393, 404)
smog, photochemical **20, 25** (456, 619-621)
smog resistance **523** (984)
smoke, **96, 97** (625-627, 629, 637, 648)
wildfire **96** (648)
smoke management **84** (622, 624)
snow **335, 345** (72, 131)
cover **334** (76)
density **321** (41)
management **318** (42)
movement **272** (581)
samples **321** (41)
snowmelt **323, 335** (33, 72)
softwood cut-up **559** (1231)
soil bank **478** (911)
chemistry **391** (771)
fertility **446** (861)
freezing **341** (77)
improvement **383-398** (758-780)
management **398** (780)
moisture **389** (766, 767)
phosphorus **385, 476** (760, 909, 910)
plant animal relations **27-30** (152, 162, 163, 169, 170)
ripping **331** (50)
salinity **218** (555)
water **342** (66)
soils, organic **325** (57)
SOLAR **219** (492, 530)
solid wastes **596** (1314, 1332, 1333, 1337, 1338)
sonic boom **345** (131)
southern hardwoods **208, 209, 211** (375-377)
pine (s) **192, 259, 515, 557, 566, 577, 582, 589** (333, 397, 974, 1225, 1254, 1273, 1284, 1302)
pine beetle **136-138** (321, 359, 360)
pine plywood **564** (1245)
spacing **474** (822, 868, 907)
Spanish cedar **517** (976)
spanworm, elm **169** (332)
species adaptability **410** (797)
composition **434** (843)
specific gravity, wood **575, 580, 589** (1271, 1282, 1302)
spiral grain (1025)
spoil banks **171, 338, 471** (102, 484, 900)
stabilization **336** (14)
spray deposit assessment **16** (298)
spraying, aerial **16** (298)
spraying equipment **253** (406)
sprouting, **441** (854)
epicormic **435** (844)
spruce, black **346** (127, 804)
Engelmann **421, 460, 486** (728, 729, 815, 816, 883, 903, 922, 923)
Norway **507** (722, 963, 964)
red **518** (977)
Sitka **353, 568** (724, 1259)

white **146, 226, 357, 366, 420, 437, 514** (323, 324, 680, 681, 683-685, 755, 814, 846, 972)
spruce beetle **141, 143, 232** (349, 354, 416)
budworm **129** (300)
pine **578** (1275)
stabilization **10** (8)
stand basal area **481** (915)
condition **358** (734, 735)
conversion (743)
density **481** (915)
improvement **432-441, 465** (838-854, 892)
rehabilitation (787)
simulation **365** (752, 754)
yield tables **489** (929)
statistical procedures (932)
storage treatment **629** (618)
stratification **415** (809)
stratified sampling (925)
streamflow **335** (72)
timing **323** (33)
streams, **349** (145)
mountain **330** (79)
Streptomyces **621** (608)
stress grading **606** (1355, 1356)
structural members **585** (1291)
sublimation **321, 322** (41, 46)
subterranean termites **618, 625-628** (437, 439, 443, 444, 446, 447)
succession (754)
sulfur dioxide **19, 216** (450, 509)
surface albedo **346** (127)
form **326** (125)
mining **15, 336-338** (14, 18, 101, 102)
waters **349** (145)
survey, disease **119, 123** (461-463, 527, 569)
survey methods **128** (302)
techniques **291-296** (718, 1078, 1085, 1090, 1091, 1093-1097)
surveys, range **26** (157, 158)
sweep, butt **272** (581)
sweetgum **350, 351, 519** (719, 720, 978)
swelling **570** (1262)
Swertia **181** (476)
Swietenia **409** (800)
sycamore **350, 351, 412, 474** (719, 720, 808, 826, 907)
systemic pesticides **18** (297)
systemics **259** (397)
systems ecology (753)

T

tap holes **270** (598)
taper curves **477** (912)
tapping (936)
taxonomy **499** (950)
teliospore induction **180** (489)
temperature **446** (861)
histories **110** (665)
Tennessee, forest inventory **280** (1049, 1050)
tensile strength **608** (1371)
termite, Formosan **616, 617** (436, 438)
termite metabolism **619** (433-435)
proof polymers **628** (447)
termites, subterranean **618, 625-628** (437, 439, 443, 444, 446, 447)
testing, nondestructive **608** (1371)
Texas leaf-cutting ant **256** (417, 419)
Thanasimus undatulus **140** (316)
thinning, **362, 432, 435, 437, 469, 472, 473, 560** (746, 747, 838, 839, 844, 846, 895, 896, 898, 902, 904-906, 1232)
commercial **460** (899)
Douglas-fir **299** (1101)
noncommercial (840)
precommercial **414** (807)
Three P sampling **291** (1093, 1094)
Thuja plicata **273** (580)
tillage (772)
timber cruising **569** (1260)
growing economics **301-305** (1109-1112, 1115, 1117, 1122)
harvest **289** (1074-1077)
management planning **283** (717)
measurements **487** (924)
producers associations **310** (1135)
product markets **640** (1158)
sales, preplanning **315** (1200, 1201, 1213)
utilization **284** (1062, 1065, 1067, 1068)
tip blight **172** (553)
tissue analysis (1031)
culture **177, 540** (488, 510, 511, 560, 1003)
toon, Australian **587** (829, 830, 1296)
Toona australis **587** (829, 830, 1296)
tracers **319** (36)
tracheid dimensions **577** (1273)
trail counter, wilderness **69** (277)
trail-marking pheromones **256** (417, 419)
trail use sampling **69** (277)
Trechispora **622** (607, 611, 612)
tree breeding **239-242** (570, 571, 573, 576, 577, 583, 584, 589)
breeding methodology **529-553** (991-1023)
crown width **487** (924)
deterioration **164** (386)
diameter growth **480** (109)
diameter measurements (917)
diseases **19-22, 24, 25, 194, 222** (450, 451, 455, 456, 458, 459, 477, 535, 619-621)
distribution **445** (862)
genetics **532** (994)
grades **556** (1222)
growth **122, 129** (300, 301)
hazard **274** (604)
improvement **541, 547, 549, 550** (1006, 1014, 1020, 1021)
mortality **122, 129, 215** (300, 301, 363)
resistance **142** (374)
rings **483** (917-919)
seed collection **423** (821)
stress **217** (518, 519)
survival and growth **445** (862)
susceptibility **197** (352)
value projections **298** (1100)
volume tables **485, 486** (921, 923)
trees, urban **275** (605)
trees to enhance the environment **76-80** (702-712)
tricresyl phosphate **627** (444)
tropical ash (832)
hardwoods **227, 409, 410, 429, 517** (714, 797, 800, 828, 976)
true firs **499** (950)
Tsuga **229** (596)
tupelo, (1029)
swamp **352** (723)
water **389** (766, 767)
tussock moth **124, 164, 247** (308, 386, 420)

U

Ulmus **235** (574, 585)
americana **271** (597, 599)
ultraviolet degradation **592** (1312)
upland oaks **468** (764, 897)
urban fire zones **651** (1178-1181)
trees **275** (605)
utilization, timber **284** (1062, 1065, 1067, 1068)
waste **597** (1315, 1327, 1335)
wood **566** (1254)

V

Vaccinium **75, 156** (507, 700, 701)
vacuum pumping **492, 493, 495** (939-941)
vapor barriers **605** (1353)
variation, **269** (572)
inherent **501-519** (901, 944, 952-980)
vascular rays **447** (863, 872)
wilt **186, 187, 189** (499, 500, 515, 528)
vegetation **349** (145)
vegetation for recreation areas **61** (286)
veneer **573, 585** (1267, 1291)
bolts **477** (912)
cutting **581** (1283)
quality **581** (1283)
slicing **581** (1283)
vertebrate predators **141, 213** (345, 349)
virulence **137** (359)
virus(es) **124, 247, 248, 253** (308, 395, 406, 420)
visitor impact **57** (263)
information **68** (278)
preferences **62, 64** (272, 284)
use sampling **70, 71** (276, 294)
visual resources **48, 49** (256, 257)
volume tables, tree **485, 486** (921, 923)

W

walnut, **550** (1021)
black **392, 393, 395, 430, 431, 453, 501, 502, 529, 538, 558** (741, 773, 774, 776, 834, 835, 841, 873, 953, 954, 991, 1001, 1036, 1227)
warp **567, 576** (1257, 1272)
waste utilization **597** (1315, 1327, 1335)
wastepaper **596** (1314, 1332, 1333, 1337, 1338)
wastes, solid **596** (1314, 1332, 1333, 1337, 1338)
woods **595** (1313, 1326)
water, soil **342** (66)

water depletion **320** (38)
level tolerance **445** (862)
movement **319** (36)
quality **11, 15, 343** (16, 18, 107, 682)
repellency **5** (3)
storage **320** (38)
table **325** (57)
temperature **327** (80)
thickening **98** (661)
tupelo **389** (766, 767)
yield **326, 327, 332, 340, 342, 343** (66, 80, 97, 107, 113, 125)
waterfowl management **34** (160)
watershed behavior **326** (125)
management **343** (107)
wavy grain **508** (965)
weather, mesosystem **380** (689)
weather control evaporation **321** (41)
modification and effects **375-382** (687-697)
station **108** (649)
weathering, **7, 15** (1, 18)
mineral **348** (142)
weed control **13** (21)
weeding (798, 840)
weevil, white-pine **525** (986, 987)
weevils **168** (356)
weight scaling **557** (1225)
West Pakistan **80** (708-712)
western budworm **17, 128, 130, 131, 160, 161, 251** (299, 392, 305, 310, 318, 353, 412)
distribution **170** (330)
pines **142** (374)
wetlands **346** (127)
white grubs **268** (402)
pine blister rust **239** (571)
pine weevil **525** (986, 987)
pines **500, 521, 526** (951, 981, 988)
rot(s) **204, 623** (504, 610)
whortleberry **75** (700, 701)
wilderness, visitor impact **57** (263)
wilderness fire effect **56** (259, 266)
management **53-55** (260, 261, 265)
trail counter **69** (277)
wildfire **5, 118, 327** (3, 80, 669, 671, 675, 679)
smoke **96** (648)
wildland management **347** (133)
wildlife conservation bibliography **65** (289)
food plants **31, 40, 41, 43** (183, 190, 202, 215)
habitat **221, 224** (341, 506)
habitat, evaluating and improving **36** (699)
management **33, 46, 47** (171, 246, 249)
willow rooting **536** (999)
wilt, oak **119, 187, 188, 235** (371, 461-463, 515, 574, 585)
vascular **186, 187, 189** (499, 500, 515, 528)
wind **322, 466** (46, 893)
flows **379** (688, 690, 695, 696)
Wisconsin, forest inventory **278** (1053)
witches'-broom **182** (542)
wood anatomy **594** (1026, 1027)
and timber quality **593, 594** (1024-1027)
boring **574** (1270)
bridge decks **614** (1390)
bridges **614** (1390)
characteristics **573** (1267)
combustion **110** (665)
decay **203, 269, 622-624, 629, 630** (531, 541, 543, 545, 547, 572, 607, 610-612, 614-618)
density **436, 575, 580, 589, 593** (845, 1024, 1271, 1282, 1302)
development **447** (863, 872)
drying **572, 590** (1264, 1304)
engineering **606** (1355, 1356)
fibers **601, 604** (1325, 1334, 1348, 1349, 1351)
finishing **592** (1312)
framing **610** (1383)
floors **615** (1391-1393)
machining **574** (1270)
moisture **610** (1383)
moisture content **557, 572, 574** (1225, 1264, 1270)
moisture fasteners **615** (1391-1393)
moisture relations **558** (1227)
preservation **588** (1300)
preservatives **630** (616, 617)
processing **566** (741, 1254)
production **455** (876)
products (741)
products in I-beams **611** (1384)
properties **464, 516, 554, 555, 558, 566, 573, 578** (891, 975, 1220, 1221, 1227, 1254, 1267, 1275)
quality **464, 466, 575, 580** (891, 893, 1271, 1282, 1302)
shrinkage **584** (1288)
specific gravity **575, 580, 589** (1272, 1282, 1302)
structures **610** (1383)
swelling **584** (1288)
utilization **566** (1254)
woods wastes **595** (1313, 1326)
woodland owners **304** (1109)
species **554** (1220)
woodpeckers **141, 213** (345, 349)
wound dressings **271** (597, 599)
wounds **203, 207** (531, 541, 543, 545-547)

X

x-ray(s) **132, 512** (460, 470-472, 970, 979)
xylem derivatives, cambial (1038)
resin **142** (374)
Xylosandrus compactus **215** (363)

Y

yellow-poplar **422, 427, 435, 463, 516, 542** (818, 825, 844, 889, 975, 1004, 1005
yield tables **467, 468, 477, 478, 489** (894, 897, 911, 912, 929)

U. S. GOVERNMENT PRINTING OFFICE : 1974 O - 547-679